

Prior Authorization Criteria

BCBS-SC PDP

Effective: 07/01/2021

## **ACTEMRA SC (S)**

---

### **Products Affected**

- Actemra INJ 162MG/0.9ML
- Actemra Actpen

Last Updated: June 2021

Formulary ID: 21381, Version 14

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. One of the following: a) Either a trial and failure, contraindication, or intolerance (TF/C/I) to two of the following: Enbrel (etanercept), Humira (adalimumab), Rinvoq (upadacitinib), Xeljanz/Xeljanz XR (tofacitinib) or attestation demonstrating a trial may be inappropriate, OR b) For continuation of prior therapy. Giant Cell Arteritis (GCA) (Initial): Diagnosis of GCA. Trial and failure, contraindication, or intolerance to a glucocorticoid. Systemic Juvenile Idiopathic Arthritis (SJIA) (Initial): Diagnosis of active SJIA. Trial and failure, contraindication, or intolerance to one of the following: NSAID, methotrexate, or systemic glucocorticoid. Polyarticular Juvenile Idiopathic Arthritis (PJIA) (Initial): Diagnosis of active PJIA. One of the following: Trial and failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior therapy. Systemic sclerosis-associated interstitial lung disease (SSc-ILD) (Initial): Diagnosis of SSc-ILD as documented by the following: a) Exclusion of other known causes of ILD (eg, domestic and occupational environmental exposures, connective tissue disease, drug toxicity) AND b) One of the following: i) In patients not subjected to surgical lung biopsy, the presence of idiopathic interstitial pneumonia (eg, fibrotic nonspecific interstitial pneumonia [NSIP], usual interstitial pneumonia [UIP] and centrilobular fibrosis) pattern on high-resolution computed tomography (HRCT) revealing SSc-ILD or probable SSc-ILD, OR ii) In patients subjected to a lung biopsy, both HRCT and surgical lung biopsy pattern revealing SSc-ILD or probable SSc-ILD.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	RA, GC, SJIA, PJIA (initial): Prescribed by or in consultation with a rheumatologist. SSc-ILD (initial): Prescribed by or in consultation with a pulmonologist or rheumatologist.
<b>Coverage Duration</b>	RA, GC, SJIA, PJIA, SSc-ILD (initial, reauth): 12 months

Last Updated: June 2021

<b>Other Criteria</b>	RA, GC, SJIA, PJIA, SSc-ILD (Reauth): Documentation of positive clinical response to therapy.
-----------------------	-----------------------------------------------------------------------------------------------

# ACTIMMUNE (S)

---

## Products Affected

- Actimmune

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of one of the following: 1) Chronic granulomatous disease (CGD), or 2) severe malignant osteopetrosis (SMO).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

## ADCIRCA (S)

---

### Products Affected

- Alyq
- Tadalafil TABS 20MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist.
<b>Coverage Duration</b>	PAH: Initial: 6 months. Reauth: 12 months.
<b>Other Criteria</b>	PAH (Reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

## ADEMPAS (S)

---

### Products Affected

- Adempas

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH AND PAH is symptomatic AND One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH. Chronic thromboembolic pulmonary hypertension (CTEPH) (Initial): One of the following: A) Both of the following: 1) Diagnosis of inoperable or persistent/recurrent CTEPH and 2) CTEPH is symptomatic OR B) Patient is currently on any therapy for the diagnosis of CTEPH.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	PAH, CTEPH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist.
<b>Coverage Duration</b>	PAH, CTEPH: Initial: 6 months. Reauth: 12 months.
<b>Other Criteria</b>	PAH, CTEPH (Reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

# AFINITOR (S)

---

## **Products Affected**

- Afinitor TABS 10MG
- Everolimus TABS 2.5MG, 5MG, 7.5MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Subependymal Giant Cell Astrocytoma (SEGA) associated with tuberous sclerosis complex (TSC): Diagnosis of SEGA associated with TSC that requires therapeutic intervention but patient is not a candidate for curative surgical resection. Renal cell carcinoma: Diagnosis of advanced or metastatic renal cell carcinoma AND trial and failure, contraindication, or intolerance to SUTENT (sunitinib) or NEXAVAR (sorafenib). Neuroendocrine tumors of pancreatic origin (pNET): Diagnosis of progressive pNET that are unresectable, locally advanced, or metastatic. Renal angiomyolipoma: Diagnosis of renal angiomyolipoma and TSC AND Patient does not require immediate surgery. Breast Cancer: Patient is a postmenopausal woman AND Diagnosis of advanced hormone receptor-positive, HER2-negative breast cancer AND trial and failure, contraindication, or intolerance to FEMARA (letrozole) or ARIMIDEX (anastrozole) AND used in combination with AROMASIN (exemestane). Neuroendocrine tumors of gastrointestinal (GI) or lung origin: Diagnosis of progressive, well-differentiated, non-functional NET of GI or lung origin AND patient has unresectable, locally advanced or metastatic disease.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses: Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	All uses: 12 months
<b>Other Criteria</b>	All Indications: Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14



# AFINITOR DISPERZ (S)

---

## Products Affected

- Afinitor Disperz

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Subependymal Giant Cell Astrocytoma (SEGA): Diagnosis of SEGA associated with tuberous sclerosis complex (TSC) that requires therapeutic intervention but patient is not a candidate for curative surgical resection. TSC-associated partial-onset seizures: Diagnosis of TSC-associated partial-onset seizures. Used as adjunctive therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	SEGA: Prescribed by or in consultation with an oncologist. TSC-associated partial-onset seizures: Prescribed by or in consultation with a neurologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

# AIMOVIG (S)

---

## **Products Affected**

- Aimovig

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	<p>Episodic Migraines (EM) (initial): Diagnosis of EM. Patient has 4 to 14 migraine days per month, but no more than 14 headache days per month.</p> <p>Chronic Migraines (CM) (initial): Diagnosis of CM. Medication overuse headache has been considered and potentially offending medication(s) have been discontinued. Patient has greater than or equal to 15 headache days per month, of which at least 8 must be migraine days for at least 3 months.</p> <p>All Indications (initial): Two of the following: a) History of failure (after at least a two month trial) or intolerance to Elavil (amitriptyline) or Effexor (venlafaxine), OR patient has a contraindication to both Elavil (amitriptyline) and Effexor (venlafaxine), b) History of failure (after at least a two month trial) or intolerance to Depakote/Depakote ER (divalproex sodium) or Topamax (topiramate), OR patient has a contraindication to both Depakote/Depakote ER (divalproex sodium) and Topamax (topiramate), or c) History of failure (after at least a two month trial) or intolerance to one of the following beta blockers: atenolol, propranolol, nadolol, timolol, or metoprolol, OR patient has a contraindication to all of the following beta blockers: atenolol, propranolol, nadolol, timolol, or metoprolol. Medication will not be used in combination with another injectable CGRP inhibitor.</p>
<b>Age Restrictions</b>	EM, CM (initial): 18 years of age or older.
<b>Prescriber Restrictions</b>	EM, CM (initial, reauth): Prescribed by or in consultation with a neurologist, headache specialist, or pain specialist.
<b>Coverage Duration</b>	EM, CM (initial): 6 months. EM, CM (reauth): 12 months.
<b>Other Criteria</b>	<p>EM, CM (reauth): Patient has experienced a positive response to therapy, demonstrated by a reduction in headache frequency and/or intensity. Use of acute migraine medications (e.g., non-steroidal anti-inflammatory drugs [NSAIDs], triptans) has decreased since the start of CGRP therapy. Medication will not be used in combination with another injectable CGRP inhibitor.</p> <p>CM (reauth): Patient continues to be monitored for medication overuse headache.</p>

Last Updated: June 2021

## ALDURAZYME (S)

---

### Products Affected

- Aldurazyme

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Mucopolysaccharidosis I: confirmed diagnosis of Hurler and Hurler-Scheie forms of Mucopolysaccharidosis I (MPS I), OR confirmed diagnosis of Scheie form of Mucopolysaccharidosis I (MPS I) who have moderate to severe symptoms.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

12

## ALECENSA (S)

---

### Products Affected

- Alecensa

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Non-small cell lung cancer (NSCLC): A) Diagnosis of metastatic NSCLC AND B) Patient has anaplastic lymphoma kinase (ALK)-positive disease as detected with a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

13

## ALPHA-1 PROTEINASE INHIBITOR, NON-PREFERRED (S)

### Products Affected

- Aralast Np INJ 1000MG, 500MG
- Zemaira

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Alpha-1 antitrypsin (AAT) deficiency: Diagnosis of congenital AAT deficiency. Diagnosis of emphysema. Continued optimal conventional treatment for emphysema (e.g., bronchodilators). One of the following: 1) PiZZ, PiZ(null), or Pi(null)(null) protein phenotypes (homozygous) OR 2) other rare AAT disease-causing alleles associated with serum AAT level less than 11 $\mu$ M/L [e.g., Pi(Malton, Malton)]. One of the following: Circulating serum concentration of AAT level less than 11 $\mu$ M/L (which corresponds to less than 80 mg/dL if measured by radial immunodiffusion or less than 57 mg/dL if measured by nephelometry) OR the patient has a concomitant diagnosis of necrotizing panniculitis. One of the following: FEV1 level is between 30% and 65% of predicted OR the patient has experienced a rapid decline in lung function (i.e., reduction of FEV1 more than 120 mL/year) that warrants treatment OR the patient has a concomitant diagnosis of necrotizing panniculitis. Trial and failure, or intolerance to Prolastin.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

14

# ALPHA-1 PROTEINASE INHIBITOR, PROLASTIN (S)

## Products Affected

- Prolastin-c INJ 1000MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Alpha-1 antitrypsin (AAT) deficiency: Diagnosis of congenital AAT deficiency. Diagnosis of emphysema. Continued optimal conventional treatment for emphysema (e.g., bronchodilators). One of the following: 1) PiZZ, PiZ(null), or Pi(null)(null) protein phenotypes (homozygous) OR 2) other rare AAT disease-causing alleles associated with serum AAT level less than 11 $\mu$ M/L [e.g., Pi(Malton, Malton)]. Circulating serum concentration of AAT level less than 11 $\mu$ M/L (which corresponds to less than 80 mg/dL if measured by radial immunodiffusion or less than 57 mg/dL if measured by nephelometry), unless the patient has a concomitant diagnosis of necrotizing panniculitis. One of the following: FEV1 level is between 30% and 65% of predicted OR the patient has experienced a rapid decline in lung function (i.e., reduction of FEV1 more than 120 mL/year) that warrants treatment OR the patient has a concomitant diagnosis of necrotizing panniculitis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

15

## ALUNBRIG (S)

---

### Products Affected

- Alunbrig

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Non-small cell lung cancer (NSCLC): Diagnosis of metastatic NSCLC. Patient has an anaplastic lymphoma kinase (ALK)-positive tumor as detected with a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

16



## AMPYRA (S)

---

### Products Affected

- Dalfampridine Er

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple Sclerosis (MS) (initial): Diagnosis of MS. Physician confirmation that patient has difficulty walking (eg, timed 25 foot walk test). One of the following: expanded disability status scale (EDSS) score less than or equal to 7, or not restricted to using a wheelchair (if EDSS is not measured).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	MS (initial): Prescribed by or in consultation with a neurologist.
<b>Coverage Duration</b>	MS (Initial): 6 months. (Reauth): 12 months.
<b>Other Criteria</b>	MS (Reauth): Physician confirmation that the patient's walking improved with therapy. One of the following: EDSS score less than or equal to 7, or not restricted to using a wheelchair (if EDSS is not measured).

Last Updated: June 2021

Formulary ID: 21381, Version 14

17

# ANADROL-50 (S)

---

## Products Affected

- Anadrol-50

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Anemia (initial): Diagnosis of anemia caused by deficient red cell production AND trial and failure or intolerance to two standard therapies for anemia (i.e., erythropoiesis-stimulating agents, immunosuppressants) AND Treatment will not replace other supportive measures (e.g., transfusion, correction of iron, folic acid, vitamin B12 or pyridoxine deficiency, antibacterial therapy, corticosteroids).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Initial, reauth: 12 months
<b>Other Criteria</b>	Anemia (reauth): Documentation of a positive clinical response to therapy as evidenced by an improvement in anemia (e.g., increased hemoglobin, increased reticulocyte count, reduction/elimination for need of blood transfusions).

Last Updated: June 2021

Formulary ID: 21381, Version 14

18

## APOKYN (S)

---

### Products Affected

- Apokyn INJ 30MG/3ML

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	PD (Initial): Used with any 5-HT3 antagonist (eg, ondansetron, granisetron, dolasetron, palonosetron, alosetron)
<b>Required Medical Information</b>	Parkinson's disease (PD) (Initial): Diagnosis of PD. Patient is experiencing acute intermittent hypomobility (defined as “off” episodes characterized by muscle stiffness, slow movements, or difficulty starting movements). Used in combination with other medications for the treatment of PD (e.g., carbidopa/levodopa, pramipexole, ropinirole, etc.).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	PD (Initial): Prescribed by or in consultation with a neurologist.
<b>Coverage Duration</b>	PD (Initial, reauth): 12 months
<b>Other Criteria</b>	PD (Reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

19

# ARANESP (S)

---

## Products Affected

- Aranesp Albumin Free INJ  
100MCG/0.5ML, 100MCG/ML,  
10MCG/0.4ML, 150MCG/0.3ML,  
200MCG/0.4ML, 200MCG/ML,  
25MCG/0.42ML, 25MCG/ML,  
300MCG/0.6ML, 300MCG/ML,  
40MCG/0.4ML, 40MCG/ML,  
500MCG/ML, 60MCG/0.3ML,  
60MCG/ML

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Anemia with Chronic Kidney Disease (CKD) (Initial): Diagnosis (Dx) of CKD. Anemia by lab values (Hct less than 30% or Hgb less than 10 g/dL) collected within 30 days of request. One of the following: a) both of the following: Patient is on dialysis, patient is without ESRD OR b) all of the following: patient is not on dialysis, the rate of hemoglobin decline indicates the likelihood of requiring a red blood cell (RBC) transfusion, and reducing the risk of alloimmunization and/or other RBC transfusion-related risks is a goal. Anemia with chemo (Initial): Other causes of anemia have been ruled out. Anemia by lab values (Hct less than 30%, Hgb less than 10 g/dL) collected within the prior 2 weeks of request. Cancer is a non-myeloid malignancy. Patient is receiving chemo. CKD (init, reauth), Chemo (init), MDS (init): Verify iron evaluation for adequate iron stores. Anemia in Myelodysplastic Syndrome (MDS) (Initial): Dx of MDS. Serum erythropoietin level is 500 mU/mL or less, or dx of transfusion-dependent MDS.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	CKD (Init): 6 mo. CKD (reauth): 12 mo. Chemo(init, reauth): 3 mo. MDS: (init) 3 mo,(reauth) 12 mo.
<b>Other Criteria</b>	Subject to ESRD review. CKD (Reauth): Dx of CKD. One of the following: 1) Most recent or average (avg) Hct over 3 mo is 33% or less (Hgb 11 g/dL or less) for patients on dialysis, without ESRD, 2) Most recent or avg Hct over 3 mo is 30% or less (Hgb 10 g/dL or less) for patients not on dialysis, OR 3) Most recent or avg Hct over 3 mo is 36% or less (Hgb 12 g/dL or less) for pediatric patients. Decrease in the need for blood transfusion or Hgb increased by 1 g/dL or more from pre-treatment level. Chemo (Reauth): Anemia by lab values (Hgb less than 10 g/dl or Hct less than 30%) collected within the prior 2 weeks of request. Decrease in the need for blood transfusion or Hgb increased by 1 g/dL or more from pre-treatment level. Patient is receiving chemo. MDS (Reauth): Most recent or avg Hct over 3 months is 36% or less, OR most

Last Updated: June 2021

	recent or avg Hgb over 3 months is 12 g/dl or less. Decrease in the need for blood transfusion or Hgb increased by 1 g/dL or more from pre-treatment level. Off-label uses (except MDS, HCV): Will not be approved if patient has Hgb greater than 10 g/dL or Hct greater than 30%.
--	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

# ARCALYST (S)

---

## **Products Affected**

- Arcalyst

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cryopyrin-Associated Periodic Syndromes (CAPS) (Initial): Diagnosis of CAPS, including Familial Cold Auto-inflammatory Syndrome (FCAS) and/or Muckle-Wells Syndrome (MWS). The medication will not be used in combination with another biologic. Deficiency of Interleukin-1 Receptor Antagonist (DIRA): Diagnosis of DIRA. Patient weighs at least 10 kg. Patient is currently in remission (e.g., no fever, skin rash, and bone pain/no radiological evidence of active bone lesions/C-reactive protein [CRP] less than 5 mg/L). Recurrent Pericarditis (Initial): Diagnosis of recurrent pericarditis as evidenced by at least 2 episodes that occur a minimum of 4 to 6 weeks apart. Trial and failure, contraindication, or intolerance (TF/C/I) to at least one of the following: nonsteroidal anti-inflammatory drugs (e.g., ibuprofen, naproxen), colchicine, or corticosteroids (e.g., prednisone).
<b>Age Restrictions</b>	CAPS (Initial): 12 years of age or older
<b>Prescriber Restrictions</b>	CAPS (initial): Prescribed by or in consultation with an immunologist, allergist, dermatologist, rheumatologist, neurologist or other medical specialist. Recurrent Pericarditis (initial): Prescribed by or in consultation with a cardiologist.
<b>Coverage Duration</b>	CAPS, Recurrent Pericarditis (initial, reauth): 12 months. DIRA: 12 months.
<b>Other Criteria</b>	CAPS (Reauth): Patient has experienced disease stability or improvement in clinical symptoms while on therapy as evidence by one of the following: A) improvement in rash, fever, joint pain, headache, conjunctivitis, B) decreased number of disease flare days, C) normalization of inflammatory markers (CRP, ESR, SAA), D) corticosteroid dose reduction, OR E) improvement in MD global score or active joint count. Recurrent Pericarditis (Reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

24



## AURYXIA (S)

---

### Products Affected

- Auryxia

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	Exclude if used for iron deficiency anemia in chronic kidney disease (CKD) not on dialysis.
<b>Required Medical Information</b>	Hyperphosphatemia in chronic kidney disease: Diagnosis of hyperphosphatemia. Patient has chronic kidney disease (CKD). Patient is on dialysis.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

25

# AUSTEDO (S)

---

## Products Affected

- Austedo

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chorea associated with Huntington's disease (initial): Diagnosis of Chorea associated with Huntington's disease. Tardive dyskinesia (initial): Diagnosis of moderate to severe tardive dyskinesia. One of the following: 1) Patient has persistent symptoms of tardive dyskinesia despite a trial of dose reduction, tapering, or discontinuation of the offending medication or 2) Patient is not a candidate for a trial of dose reduction, tapering, or discontinuation of the offending medication.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Huntington's disease chorea (initial): Prescribed by a neurologist. Tardive dyskinesia (initial): Prescribed by or in consultation with a neurologist or psychiatrist.
<b>Coverage Duration</b>	Initial: 3 months. Reauth: 12 months
<b>Other Criteria</b>	All indications (Reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

26

# AVASTIN (S)

---

## Products Affected

- Avastin

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	Non-Small Cell Lung Cancer: Excluded if squamous cell histology.
<b>Required Medical Information</b>	<p>Colorectal Cancer: Diagnosis of metastatic colorectal cancer. One of the following: 1) Both of the following: a) used as first- or second-line treatment and b) used in combination with an intravenous 5-fluorouracil-based chemotherapy, OR 2) All of the following: a) used as second-line treatment, b) used in combination with fluoropyrimidine-irinotecan-based chemotherapy or fluoropyrimidine-oxaliplatin-based chemotherapy, and c) patient has progressed on a first-line bevacizumab-containing regimen.</p> <p>Non-Small Cell Lung Cancer (NSCLC): Diagnosis of NSCLC. Disease is unresectable, locally advanced, recurrent, or metastatic. Used as first-line treatment. Used in combination with paclitaxel and carboplatin.</p> <p>Renal Cell Cancer: Diagnosis of metastatic renal cell cancer. Used in combination with interferon-alpha.</p> <p>Cervical Cancer: Diagnosis of carcinoma of the cervix. Disease is persistent, recurrent, or metastatic. Used in combination with one of the following: a) paclitaxel and cisplatin or b) paclitaxel and topotecan.</p> <p>Glioblastoma: Diagnosis of recurrent glioblastoma.</p> <p>Epithelial Ovarian, Fallopian Tube, or Primary Peritoneal Cancer: Diagnosis of epithelial ovarian cancer, fallopian tube cancer, or primary peritoneal cancer. One of the following: 1) All of the following: a) disease is stage 3 or 4, b) patient has been treated with bevacizumab as a single agent, c) treatment is following surgical resection, and d) used in combination with carboplatin and paclitaxel, OR 2) All of the following: a) disease is platinum-resistant recurrent, b) patient has received no more than 2 prior chemotherapy regimens, and c) used in combination with paclitaxel, pegylated liposomal doxorubicin, or topotecan, OR 3) All of the following: a) disease is platinum-sensitive recurrent, b) patient has been treated with bevacizumab as a single agent, and c) used in combination with one of the following: i) carboplatin and paclitaxel or ii) carboplatin and gemcitabine.</p>
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months

Last Updated: June 2021

<b>Other Criteria</b>	Approve for continuation of prior therapy.
-----------------------	--------------------------------------------

## AYVAKIT (S)

---

### Products Affected

- Ayvakit

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Gastrointestinal stromal tumor (GIST): Diagnosis of GIST. Disease is one of the following: unresectable or metastatic. Presence of platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation, including PDGFRA D842V mutations.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

30

# BAFIERTAM (S)

---

## Products Affected

- Bafiertam

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple Sclerosis (MS): Diagnosis of a relapsing form of MS (e.g., clinically isolated syndrome, relapsing-remitting disease, secondary progressive disease, including active disease with new brain lesions). One of the following: a) Failure after a trial of at least 4 weeks, contraindication, or intolerance to two of the following disease-modifying therapies for MS: 1) Aubagio (teriflunomide), 2) Gilenya (fingolimod), or 3) Brand Tecfidera/generic dimethyl fumarate, OR b) for continuation of prior therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

## BALVERSA (S)

---

### Products Affected

- Balversa

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Urothelial Carcinoma: Diagnosis of urothelial carcinoma (UC). One of the following: Locally advanced or Metastatic AND Patient has fibroblast growth factor receptor (FGFR) 3 or FGFR2 genetic alterations as detected by an U.S. Food and Drug Administration (FDA)-approved test (therascreen FGFR RGQ RT-PCR Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). One of the following: 1) Patient has progressed during or following at least one line of prior chemotherapy or immunotherapy OR 2) Patient has progressed within 12 months of neoadjuvant or adjuvant platinum-containing chemotherapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

32



# BAVENCIO (S)

---

## Products Affected

- Bavencio

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Merkel Cell Carcinoma (MCC): Diagnosis of metastatic Merkel cell carcinoma. Urothelial Carcinoma (UC): Diagnosis of locally advanced or metastatic urothelial carcinoma. One of the following: 1) Patient has disease progression during or following platinum-containing chemotherapy, OR 2) Patient has disease progression within 12 months of neoadjuvant or adjuvant treatment with platinum-containing chemotherapy. Renal Cell Carcinoma (RCC): Diagnosis of advanced renal cell carcinoma. Used as first-line treatment in combination with Inlyta (axitinib).
Age Restrictions	MCC: Patient is 12 years of age or older.
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

## BENLYSTA (S)

---

### Products Affected

- Benlysta

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Systemic lupus erythematosus (SLE) (init): Diagnosis of active SLE. Autoantibody positive (ie, anti-nuclear antibody [ANA] titer greater than or equal to 1:80 or anti-dsDNA level greater than or equal to 30 IU/mL). Currently receiving at least one standard of care treatment for active SLE (eg, antimalarials [eg, Plaquenil (hydroxychloroquine)], corticosteroids [eg, prednisone], or immunosuppressants [eg, methotrexate, Imuran (azathioprine)]). Lupus Nephritis (init): Diagnosis of active lupus nephritis. Currently receiving standard of care treatment for active lupus nephritis (e.g., corticosteroids [e.g., prednisone] with mycophenolate or cyclophosphamide).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	SLE (init): Prescribed by or in consultation with a rheumatologist. Lupus Nephritis (init): Prescribed by or in consultation with a nephrologist or rheumatologist.
<b>Coverage Duration</b>	SLE, Lupus Nephritis (init, reauth): 6 months
<b>Other Criteria</b>	SLE, Lupus Nephritis (reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

34

## BERINERT (S)

---

### Products Affected

- Berinert

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Treatment of hereditary angioedema (HAE) attacks: Diagnosis of HAE. For the treatment of acute HAE attacks. Not used in combination with other approved treatments for acute HAE attacks.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	HAE: Prescribed by or in consultation with an immunologist, allergist, or rheumatologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

35

## BOSULIF (S)

---

### Products Affected

- Bosulif

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chronic myelogenous/myeloid leukemia (CML): Diagnosis of Philadelphia chromosome-positive (Ph+) CML.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist or hematologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

36

# BRAFTOVI (S)

---

## Products Affected

- Braftovi

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Melanoma: Diagnosis of unresectable melanoma or metastatic melanoma. Cancer is BRAF V600E or V600K mutant type (MT) as detected by a U.S. Food and Drug Administration (FDA)-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Used in combination with Mektovi (binimetinib). Colorectal Cancer: One of the following diagnoses: Colon Cancer or Rectal Cancer. One of the following: 1) Unresectable or advanced disease or 2) Metastatic disease. Patient has received prior therapy. Cancer is BRAF V600E mutant type as detected by a U.S. Food and Drug Administration (FDA)-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Used in combination with Erbitux (cetuximab).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy

Last Updated: June 2021

## BRIVIACT (S)

---

### Products Affected

- Briviact SOLN
- Briviact TABS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Partial-onset seizures: Diagnosis of partial-onset seizures.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

38

# BRONCHITOL (S)

---

## Products Affected

- Bronchitol
- Bronchitol Tolerance Test

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cystic Fibrosis (CF) (initial): Diagnosis of CF. Patient has passed the Bronchitol Tolerance Test (BTT). One of the following: 1) Patient is currently receiving Pulmozyme (dornase alfa), OR 2) Patient has a contraindication, intolerance, or is not a candidate for continued Pulmozyme therapy. Trial and failure, contraindication, or intolerance to inhaled hypertonic saline.
<b>Age Restrictions</b>	CF (initial): Patient is 18 years of age or older.
<b>Prescriber Restrictions</b>	CF (initial): Prescribed by or in consultation with a pulmonologist or specialist affiliated with a CF care center.
<b>Coverage Duration</b>	CF (initial): 6 months. CF (reauth): 12 months.
<b>Other Criteria</b>	CF (reauth): Documentation of positive clinical response to therapy (e.g., improvement in lung function [forced expiratory volume in one second {FEV1}]).

Last Updated: June 2021

Formulary ID: 21381, Version 14

39

## BRUKINSA (S)

---

### Products Affected

- Brukinsa

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of relapsed or refractory mantle cell lymphoma (MCL). Trial and failure, contraindication, or intolerance to at least ONE combination treatment of rituximab and chemotherapy (e.g., BR, R-CHOP, R-CVP, R-FCM).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist/hematologist.
<b>Coverage Duration</b>	12 months.
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14



## BYNFEZIA (S)

---

### Products Affected

- Bynfezia Pen

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acromegaly (initial): Diagnosis of acromegaly. One of the following: A) Inadequate response to surgical resection and/or pituitary irradiation OR B) Patient is not a candidate for surgical resection or pituitary irradiation. Trial and failure, contraindication or intolerance to a dopamine agonist (e.g., bromocriptine or cabergoline) at maximally tolerated doses. Carcinoid tumor (initial): Diagnosis of metastatic carcinoid tumor requiring symptomatic treatment of severe diarrhea or flushing episodes. Vasoactive intestinal peptide tumor (initial): Diagnosis of vasoactive intestinal peptide tumor requiring treatment of profuse watery diarrhea. All indications (initial): Trial and failure, or intolerance to generic octreotide.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	All uses (initial, reauth): 12 months
Other Criteria	Acromegaly (reauth): Documentation of positive clinical response to therapy (e.g., reduction or normalization of IGF-1/GH level for same age and sex, reduction in tumor size). Carcinoid tumor (reauth): Patient has improvement in number of diarrhea or flushing episodes. Vasoactive intestinal peptide tumor (reauth): Patient has improvement in number of diarrhea episodes.

Last Updated: June 2021

Formulary ID: 21381, Version 14

41

## CABLIVI (S)

---

### Products Affected

- Cablivi

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Acquired thrombotic thrombocytopenic purpura (aTTP): Diagnosis of aTTP. First dose was/will be administered by a healthcare provider as a bolus intravenous injection. Used in combination with immunosuppressive therapy (e.g. rituximab, glucocorticoids). One of the following: 1) Used in combination with plasma exchange or 2) both of the following: patient has completed plasma exchange and less than 59 days have or will have elapsed beyond the last plasma exchange.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist or oncologist.
<b>Coverage Duration</b>	3 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

## CABOMETYX (S)

---

### Products Affected

- Cabometyx

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Renal cell carcinoma (RCC): Diagnosis of RCC. RCC is advanced. Hepatocellular Carcinoma (HCC): Diagnosis of HCC. One of the following: a) Trial and failure or intolerance to Nexavar (sorafenib tosylate), or b) Patient has metastatic disease, or c) Patient has extensive liver tumor burden, or d) Patient is inoperable by performance status or comorbidity (local disease or local disease with minimal extrahepatic disease only), or e) Disease is unresectable.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	RCC: Prescribed by or in consultation with one of the following: an oncologist or nephrologist. HCC: Prescribed by or in consultation with one of the following: oncologist, hepatologist, or gastroenterologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

43

## CALQUENCE (S)

---

### Products Affected

- Calquence

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Mantle Cell Lymphoma: Diagnosis of mantle cell lymphoma (MCL) AND patient has received at least one prior therapy for MCL. Chronic Lymphocytic Leukemia (CLL) or Small Lymphocytic Lymphoma (SLL): Diagnosis of CLL or SLL.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist/oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

## CAPRELSA (S)

---

### Products Affected

- Caprelsa

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Thyroid Cancer: Diagnosis of medullary thyroid cancer (MTC).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with oncologist or endocrinologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

45

## CAYSTON (S)

---

### Products Affected

- Cayston

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cystic fibrosis (CF) (Initial, Reauth): Diagnosis of CF AND Patient has evidence of Pseudomonas aeruginosa in the lungs.
<b>Age Restrictions</b>	CF (Initial): 7 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	CF (Initial, reauth): 12 months
<b>Other Criteria</b>	CF (Reauth): Patient is benefiting from treatment (i.e. improvement in lung function [forced expiratory volume in one second (FEV1)], decreased number of pulmonary exacerbations).

Last Updated: June 2021

Formulary ID: 21381, Version 14

## CERDELGA (S)

---

### Products Affected

- Cerdelga

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Gaucher disease (Initial): Diagnosis of Gaucher disease type 1. Patient is an extensive metabolizer (EM), intermediate metabolizer (IM), or poor metabolizer (PM) of cytochrome P450 enzyme (CYP) 2D6 as detected by an FDA-cleared test.
<b>Age Restrictions</b>	Gaucher disease (initial): 18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Gaucher disease (initial, reauth): 12 months
<b>Other Criteria</b>	Gaucher disease (Reauth): Patient's condition has not progressed, as defined by ALL of the following: A) Hemoglobin level decreased greater than 1.5 g/dL from baseline, AND B) Platelet count decreased greater than 25% from baseline, AND C) Spleen volume increased greater than 25% from baseline, AND D) Liver volume increased greater than 20% from baseline.

Last Updated: June 2021

Formulary ID: 21381, Version 14

47

## CEREZYME (S)

---

### Products Affected

- Cerezyme

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Gaucher disease: Diagnosis of type 1 Gaucher disease. Patient has evidence of symptomatic disease (e.g., moderate to severe anemia, thrombocytopenia, bone disease, hepatomegaly, or splenomegaly).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Gaucher disease: 12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14



## CHENODAL (S)

---

### Products Affected

- Chenodal

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Radiolucent stones (RS) (initial): Diagnosis of radiolucent stones. Patient has a well-opacifying gallbladder visualized by oral cholecystography. Trial and failure, contraindication or intolerance to ursodiol. Patient is not a candidate for surgery. Stones are not calcified (radiopaque) or radiolucent bile pigment stones.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	RS (initial, reauth): 12 months.
<b>Other Criteria</b>	RS (reauth): Patient's disease status has been re-evaluated since the last authorization to confirm the patient's condition warrants continued treatment as evidenced by oral cholecystograms or ultrasonograms.

Last Updated: June 2021

Formulary ID: 21381, Version 14

## CHOLBAM (S)

---

### Products Affected

- Cholbam

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Bile acid synthesis disorders due to single enzyme defects (BAS) (initial): diagnosis of a bile acid synthesis disorder due to a single enzyme defect. Peroxisomal disorders (PD) (initial): All of the following: 1) diagnosis of peroxisomal disorder, 2) patient exhibits at least one of the following: a) liver disease (eg, jaundice, elevated serum transaminases), OR b) steatorrhea, OR c) complications from decreased fat-soluble vitamin absorption (eg, poor growth), AND 3) will be used as an adjunctive treatment.
Age Restrictions	N/A
Prescriber Restrictions	All uses (initial): Prescribed by a hepatologist, medical geneticist, pediatric gastroenterologist, OR other specialist that treats inborn errors of metabolism.
Coverage Duration	All uses (initial, reauth): 12 months
Other Criteria	All uses (reauth): documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

50

# CICLOPIROX (S)

---

## Products Affected

- Ciclodan SOLN
- Ciclopirox Nail Lacquer

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	All of the following: 1) Patient does not have lunula (matrix) involvement, 2) one of the following: a) Diagnosis of onychomycosis of the toenails, OR b) Diagnosis of onychomycosis of the fingernails, 3) Diagnosis of onychomycosis has been confirmed by one of the following: a) positive potassium hydroxide (KOH) preparation, OR b) culture, OR c) histology, 4) If toenail onychomycosis, patient has mild to moderate disease involving at least 1 great toenail, AND 5) Trial and failure, contraindication, or intolerance to oral terbinafine.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	48 weeks.
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

51

## CINRYZE (S)

---

### Products Affected

- Cinryze

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Prophylaxis of hereditary angioedema (HAE) attacks: Diagnosis of HAE. For prophylaxis against HAE attacks.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	HAE (prophylaxis): Prescribed by or in consultation with an immunologist, allergist, or rheumatologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

52

## COMETRIQ (S)

---

### Products Affected

- Cometriq

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Medullary thyroid cancer (MTC): Diagnosis of Metastatic MTC.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	MTC: Prescribed by or in consultation with an oncologist/hematologist or endocrinologist.
<b>Coverage Duration</b>	All uses: 12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

53

## COPIKTRA (S)

---

### Products Affected

- Copiktra

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chronic Lymphocytic Leukemia (CLL)/Small Lymphocytic Lymphoma (SLL): Diagnosis of CLL or SLL. Disease is relapsed or refractory. Trial and failure, contraindication, or intolerance to at least two prior therapies for CLL/SLL (e.g., Leukeran [chlorambucil], Gazyva [obinutuzumab], Arzerra [ofatumumab], Bendeka [bendamustine], Imbruvica [ibrutinib], Rituxan [rituximab], etc.). Follicular Lymphoma: Diagnosis of follicular lymphoma. Disease is relapsed or refractory. Trial and failure, contraindication, or intolerance to at least two prior systemic therapies for follicular lymphoma (e.g., Leukeran [chlorambucil], Gazyva [obinutuzumab], Arzerra [ofatumumab], Bendeka [bendamustine], Imbruvica [ibrutinib], Rituxan [rituximab], etc.).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist/oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

54

# CORLANOR (S)

## Products Affected

- Corlanor TABS

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chronic heart failure (CHF) (initial): Diagnosis of CHF. Patient has NYHA Class II, III, or IV symptoms. Patient has a left ventricular ejection fraction less than or equal to 35%. Patient is in sinus rhythm. Patient has a resting heart rate of greater than or equal to 70 beats per minute. One of the following: patient is on a beta-blocker (e.g., bisoprolol, carvedilol, metoprolol succinate extended release) at a maximally tolerated dose, or patient has a contraindication or intolerance to beta-blocker therapy. Patient has been hospitalized for worsening HF in the previous 12 months. Trial and failure, contraindication, or intolerance to maximally tolerated doses of an ACE inhibitor (e.g., captopril, enalapril, lisinopril) or ARB (e.g., candesartan, losartan, valsartan). Dilated Cardiomyopathy (DCM) (initial): Diagnosis of heart failure due to DCM. Patient has NYHA Class II, III, or IV symptoms. Patient is in sinus rhythm. Patient has an elevated heart rate. Trial and failure, contraindication or intolerance to one of the following: 1) Beta blocker (e.g., bisoprolol, metoprolol succinate extended release), 2) Angiotensin-converting enzyme (ACE) inhibitor (e.g., captopril, enalapril), or 3) Diuretic Agent (e.g., spironolactone, furosemide).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	CHF, DCM (initial): Prescribed by or in consultation with a cardiologist
<b>Coverage Duration</b>	CHF, DCM (initial, reauth): 12 months
<b>Other Criteria</b>	CHF, DCM (reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Last Updated: June 2021

Formulary ID: 21381, Version 14



# COSENTYX (S)

---

## **Products Affected**

- Cosentyx INJ 150MG/ML
- Cosentyx Sensoready Pen

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Plaque psoriasis (Initial): Diagnosis of moderate to severe plaque psoriasis. One of the following: Trial and failure, contraindication, or intolerance to one of the following: Enbrel (etanercept), Humira (adalimumab), Skyrizi (risankizumab), OR for continuation of prior therapy. Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. One of the following: a) Either trial and failure, contraindication, or intolerance to Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR b) for continuation of prior therapy. Ankylosing Spondylitis (AS) (Initial): Diagnosis of active AS. One of the following: Trial and failure, contraindication, or intolerance to Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior therapy. Non-radiographic axial spondyloarthritis (nr-axSpA, initial): Dx of active nr-axSpA with objective signs of inflammation (eg, C-reactive protein [CRP] levels above the upper limit of normal and/or sacroiliitis on magnetic resonance imaging [MRI], indicative of inflammatory disease, but without definitive radiographic evidence of structural damage on sacroiliac joints.) TF/C/I to two non-steroidal anti-inflammatory drugs (NSAIDs) (eg, ibuprofen, meloxicam, naproxen).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Plaque psoriasis (initial): Prescribed by or in consultation with a dermatologist. PsA (initial): Prescribed by or in consultation with a rheumatologist or dermatologist. AS, nr-axSpA (initial): Prescribed by or in consultation with a rheumatologist.
<b>Coverage Duration</b>	All uses (initial, reauth): 12 months
<b>Other Criteria</b>	PsA, AS, nr-axSpA (Reauth): Documentation of positive clinical response to therapy. Psoriasis (Reauth): Documentation of positive clinical response to therapy as evidenced by one of the following: reduction in the body surface area (BSA) involvement from baseline, OR improvement in symptoms (eg, pruritus, inflammation) from baseline.

Last Updated: June 2021

Last Updated: June 2021

Formulary ID: 21381, Version 14

## COTELLIC (S)

---

### Products Affected

- Cotellic

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Melanoma: Diagnosis of unresectable or metastatic melanoma. Patient has a BRAF V600E or V600K mutation as detected by a U.S. Food and Drug Administration (FDA)-approved test (e.g., cobas 4800 BRAF V600 Mutation Test) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Used in combination with vemurafenib.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

60

## CYSTADROPS (S)

---

### Products Affected

- Cystadrops

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cystinosis: Diagnosis of cystinosis, confirmed by elevated leukocyte cystine levels (LCL), genetic analysis of the CTNS gene or corneal cystine crystal accumulation.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

## CYSTARAN (S)

---

### Products Affected

- Cystaran

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cystinosis: Diagnosis of cystinosis, confirmed by elevated leukocyte cystine levels (LCL), genetic analysis of the CTNS gene or corneal cystine crystal accumulation.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

## D.H.E. 45 (S)

---

### Products Affected

- Dihydroergotamine Mesylate INJ

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Migraines (initial): Diagnosis of migraine headaches with or without aura. Will be used for the acute treatment of migraine. One of the following: Trial and failure or intolerance to one triptan (e.g., eletriptan, rizatriptan, sumatriptan) or contraindication to all triptans. If patient has 4 or more headache days per month, patient must meet one of the following: a) currently being treated with Elavil (amitriptyline) or Effexor (venlafaxine) unless there is a contraindication or intolerance to these medications, OR b) currently being treated with Depakote/Depakote ER (divalproex sodium) or Topamax (topiramate) unless there is a contraindication or intolerance to these medications, OR c) currently being treated with a beta blocker (i.e., atenolol, propranolol, nadolol, timolol, or metoprolol) unless there is a contraindication or intolerance to these medications. Cluster Headaches (CH) (initial): Diagnosis of cluster headache. Trial and failure, contraindication, or intolerance to sumatriptan injection.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Migraines, CH (initial, reauth): Prescribed by or in consultation with a neurologist or pain specialist.
<b>Coverage Duration</b>	Migraines, CH (initial): 3 months. Migraines, CH (reauth): 12 months.
<b>Other Criteria</b>	Migraines (reauth): Patient has experienced a positive response to therapy (e.g., reduction in pain, photophobia, phonophobia, nausea). CH (reauth): Patient has experienced a positive response to therapy, demonstrated by a reduction in headache frequency and/or intensity.

Last Updated: June 2021

## DALIRESP (S)

---

### Products Affected

- Daliresp

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chronic Obstructive Pulmonary Disease (COPD) (initial): Diagnosis of COPD. History of COPD exacerbations which required the use of systemic corticosteroids, antibiotics, or hospital admission. Trial and failure, intolerance, or contraindication to two prior therapies for COPD (e.g., Combivent, Spiriva).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	COPD (init, reauth): 12 months
<b>Other Criteria</b>	COPD (reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

64



# DARAPRIM (S)

---

## Products Affected

- Pyrimethamine TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Toxoplasmosis: 1) Patient is using pyrimethamine for the treatment of toxoplasmic encephalitis, secondary prophylaxis of toxoplasmic encephalitis, or treatment of congenital toxoplasmosis OR 2) Patient is using pyrimethamine for the primary prophylaxis of toxoplasmic encephalitis, patient has experienced intolerance to prior prophylaxis with trimethoprim-sulfamethoxazole (TMP-SMX), and one of the following: patient has been re-challenged with TMP-SMX using a desensitization protocol and is still unable to tolerate, or evidence of life-threatening reaction to TMP-SMX in the past (eg, toxic epidermal necrolysis, Stevens-Johnson syndrome). Malaria: Patient is using pyrimethamine for the treatment of acute malaria or chemoprophylaxis of malaria. Patient does not have megaloblastic anemia due to folate deficiency. The provider acknowledges that pyrimethamine is not recommended by the Centers for Disease Control and Prevention (CDC) for the treatment and/or prophylaxis of malaria.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist
Coverage Duration	12 months
Other Criteria	Toxoplasmosis only: Approve for continuation of prior therapy.

Last Updated: June 2021

## DAURISMO (S)

---

### Products Affected

- Daurismo

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Acute myeloid leukemia (AML): Diagnosis of newly-diagnosed acute myeloid leukemia (AML) AND Used in combination with low-dose cytarabine AND One of the following: 1) Patient is greater than or equal to 75 years old, or 2) Patient has comorbidities that preclude the use of intensive induction chemotherapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist/oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

66

# DEFERASIROX (S)

## Products Affected

- Deferasirox
- Jadenu Sprinkle

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chronic Iron Overload Due to Blood Transfusions (Initial): Diagnosis of chronic iron overload due to blood transfusions (transfusional hemosiderosis). Patient has a baseline ferritin level more than 1,000 mcg/L. Patient has required the transfusion of at least 100 mL/kg packed red blood cells. Myelodysplastic Syndrome (MDS) (Initial): Diagnosis of MDS. Patient has Low or Intermediate-1 disease or is a potential transplant patient. Patient has received more than 20 red blood cell transfusions. Chronic iron overload due to non-transfusion-dependent thalassemia (NTDT) (Initial): Diagnosis of chronic iron overload due to NTDT. Liver iron concentration (LIC) 5 milligrams of iron per gram of liver dry weight (mg Fe/g dw) or higher. Serum ferritin level greater than 300 mcg/L.
<b>Age Restrictions</b>	Iron Overload Due to Blood Transfusions (initial): 2 years of age or older. NTDT (initial): 10 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Iron Overload Due to Blood Transfusions, MDS (initial, reauth):12 mo. NTDT (initial, reauth): 6mo.
<b>Other Criteria</b>	Iron Overload Due to Blood Transfusions, MDS (Reauth): Patient experienced a reduction from baseline in serum ferritin level or LIC. NTDT (Reauth): Patient has LIC 3 mg Fe/g dw or higher. Patient experienced a reduction from baseline in serum ferritin level or LIC.

Last Updated: June 2021

## DIACOMIT (S)

---

### Products Affected

- Diacomit

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of seizures associated with Dravet syndrome (DS). Used in combination with clobazam.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

68

## DOJOLVI (S)

---

### Products Affected

- Dojolvi

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	Not used with any other medium-chain triglyceride (MCT) product.
<b>Required Medical Information</b>	Long-Chain Fatty Acid Oxidation Disorder (LC-FAOD) (initial): Diagnosis of a LC-FAOD. Disease has been molecularly confirmed (i.e., genetic testing).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a clinical specialist knowledgeable in appropriate disease-related dietary management (e.g., geneticist, cardiologist, gastroenterologist, etc.).
<b>Coverage Duration</b>	LC-FAOD (initial, reauth): 12 months
<b>Other Criteria</b>	LC-FAOD (reauth): Prescriber attests to continued need of therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

## **DUPIXENT (S)**

---

### **Products Affected**

- Dupixent

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Atopic dermatitis (AD) (init): Diagnosis (dx) of mod to severe AD. Trial and failure, contraindication, or intolerance (TF/C/I) to one med to high potency topical corticosteroid (eg, betamethasone, triamcinolone). One of the following: A) TF/C/I to pimecrolimus topical cream, unless patient is not a candidate for pimecrolimus therapy (e.g., immunocompromised, severe atopic dermatitis), B) TF/C/I to tacrolimus topical ointment, unless patient is not a candidate for tacrolimus ointment therapy (e.g., immunocompromised), or C) Eucrisa (crisaborole). Eosinophilic Asthma (EA) (init): Dx of mod to severe asthma. Asthma is an eosinophilic phenotype as defined by a baseline (pre-treatment) peripheral blood eosinophil level greater than or equal to 150 cells/microliter. One of the following: 1) Patient has had at least one or more asthma exacerbations requiring systemic corticosteroids within the past 12 mo, 2) Any prior intubation for an asthma exacerbation, or 3) Prior asthma-related hospitalization within the past 12 mo. One of the following: a) TF/C/I to Fasentra (benralizumab), Nucala (mepolizumab), or Cinqair (reslizumab) or b) For continuation of prior therapy. Corticosteroid Dependent Asthma (CDA) (init): Dx of mod to severe asthma. Patient is currently dependent on oral corticosteroids for the treatment of asthma. EA, CDA (init): Patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications: a) Both of the following: i) High-dose inhaled corticosteroid (ICS) [e.g., greater than 500 mcg fluticasone propionate equivalent/day] and ii) additional asthma controller medication [e.g., leukotriene receptor antagonist, long-acting beta-2 agonist (LABA), tiotropium], OR b) One max-dosed combination ICS/LABA product [e.g., Advair (fluticasone propionate/salmeterol), Dulera (mometasone/formoterol), Symbicort (budesonide/formoterol), Breo Ellipta (fluticasone/vilanterol)].
<b>Age Restrictions</b>	Asthma (initial): Age greater than or equal to 12 years. Atopic dermatitis/CRSwNP: no age restriction.
<b>Prescriber Restrictions</b>	Atopic dermatitis (Initial): Prescribed by or in consultation with one of the following: dermatologist, allergist/immunologist. Asthma (initial, reauth): Prescribed by or in consultation with a pulmonologist or allergist/immunologist. CRSwNP (initial, reauth): Prescribed by or in

Last Updated: June 2021

	consultation with an otolaryngologist, allergist/immunologist, or pulmonologist.
<b>Coverage Duration</b>	Atopic Dermatitis, CRSwNP (Init/Reauth): 12 months. Asthma (Init): 6 mo. Asthma (reauth): 12 mo.
<b>Other Criteria</b>	Chronic rhinosinusitis with nasal polyposis (CRSwNP) (initial): Diagnosis of CRSwNP. Unless contraindicated, the patient has had an inadequate response to 2 months of treatment with an intranasal corticosteroid (eg, fluticasone, mometasone). Used in combination with another agent for CRSwNP (eg, intranasal corticosteroid). Atopic dermatitis (reauth): Documentation of a positive clinical response to therapy (e.g., reduction in body surface area involvement, reduction in pruritus severity). EA (reauth): Documentation of a positive clinical response to therapy (e.g., reduction in exacerbations, improvement in forced expiratory volume in 1 second [FEV1], decreased use of rescue medications). CDA (reauth): Documentation of a positive clinical response to therapy (e.g., reduction in exacerbations, improvement in FEV1, reduction in oral corticosteroid dose). EA, CDA (reauth): Patient is currently being treated with ONE of the following unless there is a contraindication or intolerance to these medications: 1) Inhaled corticosteroid (ICS) (e.g., fluticasone, budesonide) and additional asthma controller medication [e.g., leukotriene receptor antagonist, long-acting beta-2 agonist (LABA), tiotropium] OR 2) A combination ICS/LABA product (e.g., Advair [fluticasone propionate/salmeterol], Dulera [mometasone/formoterol], Symbicort [budesonide/formoterol], Breo Ellipta [fluticasone/vilanterol]). CRSwNP (reauth): Documentation of a positive clinical response to therapy (e.g., reduction in nasal polyps score [NPS, 0-8 scale], improvement in nasal congestion/obstruction score [NC, 0-3 scale]). Used in combination with another agent for CRSwNP (eg, intranasal corticosteroid).



## DYSPORT (S)

---

### Products Affected

- Dysport

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cervical Dystonia (CD) (init): Diagnosis of CD (also known as spasmodic torticollis). Upper limb spasticity (ULS) (init): Diagnosis of ULS. Lower limb spasticity (LLS) (init): Diagnosis of lower limb spasticity.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	CD: 3 months for a single dose (init: up to 500 units, reauth: up to 1000 units). ULS, LLS: 3 months
<b>Other Criteria</b>	CD, ULS, LLS (reauth): Documentation of positive clinical response to therapy. At least 3 months have elapsed since the last treatment.

Last Updated: June 2021

Formulary ID: 21381, Version 14

73

## ELAPRASE (S)

---

### Products Affected

- Elaprase

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of Hunter Syndrome (Mucopolysaccharidosis II (MPS II))
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

# EMGALITY (S)

---

## Products Affected

- Emgality

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	<p>Episodic Migraines (EM) (120 mg/mL strength only) (initial): Diagnosis of EM. Patient has 4 to 14 migraine days per month, but no more than 14 headache days per month. Chronic Migraines (CM) (120 mg strength/mL only) (initial): Diagnosis of CM. Medication overuse headache has been considered and potentially offending medication(s) have been discontinued. Patient has greater than or equal to 15 headache days per month, of which at least 8 must be migraine days for at least 3 months. Episodic Cluster Headache (ECH) (100 mg/mL strength only) (initial): Diagnosis of episodic cluster headache. Patient has experienced at least 2 cluster periods lasting from 7 days to 365 days, separated by pain-free periods lasting at least three months. EM, CM (120 mg/mL strength only) (initial): Two of the following: a) History of failure (after at least a two month trial) or intolerance to Elavil (amitriptyline) or Effexor (venlafaxine), OR patient has a contraindication to both Elavil (amitriptyline) and Effexor (venlafaxine), b) History of failure (after at least a two month trial) or intolerance to Depakote/Depakote ER (divalproex sodium) or Topamax (topiramate), OR patient has a contraindication to both Depakote/Depakote ER (divalproex sodium) and Topamax (topiramate), or c) History of failure (after at least a two month trial) or intolerance to one of the following beta blockers: atenolol, propranolol, nadolol, timolol, or metoprolol, OR patient has a contraindication to all of the following beta blockers: atenolol, propranolol, nadolol, timolol, or metoprolol. All Indications (initial): Medication will not be used in combination with another injectable CGRP inhibitor.</p>
<b>Age Restrictions</b>	EM, CM, ECH (initial): 18 years of age or older.
<b>Prescriber Restrictions</b>	EM, CM, ECH (initial, reauth): Prescribed by or in consultation with a neurologist, headache specialist, or pain specialist.
<b>Coverage Duration</b>	EM, CM (initial): 6 months. ECH (initial): 3 months. EM, CM, ECH (reauth): 12 months.
<b>Other Criteria</b>	EM, CM (120 mg/mL strength only) (reauth): Patient has experienced a positive response to therapy, demonstrated by a reduction in headache

Last Updated: June 2021

---

<p>frequency and/or intensity. Use of acute migraine medications (e.g., non-steroidal anti-inflammatory drugs [NSAIDs], triptans) has decreased since the start of CGRP therapy. CM (120 mg/mL strength only) (reauth): Patient continues to be monitored for medication overuse headache. ECH (100 mg/mL strength only) (reauth): Patient has experienced a positive response to therapy, demonstrated by a reduction in headache frequency and/or intensity. All Indications (initial): Medication will not be used in combination with another injectable CGRP inhibitor.</p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

---

# ENBREL (S)

---

## **Products Affected**

- Enbrel
- Enbrel Mini
- Enbrel Sureclick

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. Trial and failure, contraindication, or intolerance to one disease-modifying antirheumatic drug (DMARD) [eg, methotrexate (Rheumatrex/Trexall), Arava (leflunomide), Azulfidine (sulfasalazine)]. Polyarticular Juvenile Idiopathic Arthritis (PJIA) (Initial): Diagnosis of moderately to severely active PJIA. Trial and failure, contraindication, or intolerance to one of the following DMARDs: Arava (leflunomide) or methotrexate (Rheumatrex/Trexall). Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. Plaque psoriasis (Initial): Diagnosis of moderate to severe chronic plaque psoriasis. Ankylosing Spondylitis (AS) (Initial): Diagnosis of active AS. Trial and failure, contraindication, or intolerance to two NSAIDs (e.g., diclofenac, ibuprofen, meloxicam, naproxen).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	RA (initial), PJIA (initial), AS (initial): Prescribed by or in consultation with a rheumatologist. PsA (initial): Prescribed by or in consultation with a rheumatologist or dermatologist. Plaque Psoriasis (initial): Prescribed by or in consultation with a dermatologist.
<b>Coverage Duration</b>	All uses (initial, reauth): 12 months
<b>Other Criteria</b>	RA, PJIA, PsA, AS (Reauth): Documentation of positive clinical response to therapy. Plaque psoriasis (Reauth): Documentation of positive clinical response to therapy as evidenced by one of the following: reduction in the body surface area (BSA) involvement from baseline, OR improvement in symptoms (eg, pruritus, inflammation) from baseline.

Last Updated: June 2021

Formulary ID: 21381, Version 14

79

## ENDARI (S)

---

### Products Affected

- Endari

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Sickle cell disease (initial): Diagnosis of sickle cell disease. Used to reduce acute complications of sickle cell disease. Patient has had 2 or more painful sickle cell crises within the past 12 months.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Sickle cell disease (initial): Prescribed by or in consultation with a hematologist/oncologist
<b>Coverage Duration</b>	Sickle cell disease (initial, reauth): 12 months
<b>Other Criteria</b>	Sickle cell disease (reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

80



## ENSPRYNG (S)

---

### Products Affected

- Enspryng

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Neuromyelitis Optica Spectrum Disorder (NMOSD) (initial): Diagnosis of NMOSD. Patient is anti-aquaporin-4 (AQP4) antibody positive.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	NMOSD (initial): Prescribed by or in consultation with a neurologist or ophthalmologist.
<b>Coverage Duration</b>	NMOSD (initial, reauth): 12 months
<b>Other Criteria</b>	NMOSD (reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

81

# ENTYVIO (S)

## Products Affected

- Entyvio

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Ulcerative Colitis (UC) (init): Diagnosis (Dx) of moderately to severely active UC. Trial and failure, contraindication, or intolerance (F/C/I) to one of the following conventional therapies: 6-mercaptopurine (Purinethol), aminosalicylates [eg, mesalamine (Asacol, Pentasa, Rowasa), olsalazine (Dipentum), sulfasalazine (Azulfidine, Sulfazine)], azathioprine (Imuran), corticosteroids (eg, prednisone, methylprednisolone). F/C/I to one tumor necrosis factor (TNF) inhibitor [eg, Humira (adalimumab), infliximab]. Crohn's Disease (CD) (init): Dx of moderately to severely active CD. F/C/I to one of the following medications: 6-mercaptopurine (Purinethol), azathioprine (Imuran), corticosteroids (eg, prednisone, methylprednisolone), methotrexate (Rheumatrex, Trexall). F/C/I to one TNF inhibitor [eg, Humira (adalimumab), infliximab].
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	UC, CD (init): Prescribed by or in consultation with a gastroenterologist
<b>Coverage Duration</b>	UC, CD (init): 14 weeks. UC, CD (reauth): 12 months.
<b>Other Criteria</b>	UC, CD (reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

82

## EPCLUSA PREFERRED (S)

---

### Products Affected

- Sofosbuvir/velpatasvir

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Criteria will be applied consistent with current AASLD/IDSA guideline. Diagnosis of chronic hepatitis C. Not used in combination with another HCV direct acting antiviral agent [e.g., Sovaldi (sofosbuvir)].
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with one of the following: Hepatologist, Gastroenterologist, Infectious disease specialist, HIV specialist certified through the American Academy of HIV Medicine.
<b>Coverage Duration</b>	12 to 24 weeks. Criteria will be applied consistent with current AASLD/IDSA guideline.
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

## EPIDIOLEX (S)

---

### Products Affected

- Epidiolex

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Lennox-Gastaut syndrome (LGS): Diagnosis of seizures associated with LGS. Trial of, contraindication, or intolerance to two formulary anticonvulsants (e.g., topiramate, lamotrigine, valproate). Dravet syndrome (DS): Diagnosis of seizures associated with DS. Tuberous sclerosis complex (TSC): Diagnosis of seizures associated with TSC.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	LGS, DS, TSC: Prescribed by or in consultation with a neurologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

## ERIVEDGE (S)

---

### Products Affected

- Erivedge

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Basal cell carcinoma: One of the following: A) Diagnosis of metastatic basal cell carcinoma OR B) Both of the following: 1) Diagnosis of locally advanced basal cell carcinoma AND 2) One of the following: a) Disease recurred following surgery or b) Patient is not a candidate for surgery and radiation.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist or dermatologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

85

## ERLEADA (S)

---

### Products Affected

- Erleada

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Non-metastatic castration-resistant or castration-recurrent prostate cancer (NM-CRPC): Diagnosis of non-metastatic, castration-resistant (chemical or surgical) or recurrent prostate cancer. Metastatic castration-sensitive prostate cancer (M-CSPC): Diagnosis of metastatic, castration-sensitive prostate cancer.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	NM-CRPC, M-CSPC: Prescribed by or in consultation with an oncologist or urologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

86

## ESBRIET (S)

---

### Products Affected

- Esbriet

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Idiopathic pulmonary fibrosis (IPF) (initial): Diagnosis of IPF as documented by all of the following: a) exclusion of other known causes of interstitial lung disease (ILD) (eg, domestic and occupational environmental exposures, connective tissue disease, drug toxicity), AND b) one of the following: i) in patients not subjected to surgical lung biopsy, the presence of a usual interstitial pneumonia (UIP) pattern on high-resolution computed tomography (HRCT) revealing IPF or probable IPF, OR ii) in patients subjected to a lung biopsy, both HRCT and surgical lung biopsy pattern revealing IPF or probable IPF.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	IPF (initial): Prescribed by or in consultation with a pulmonologist
<b>Coverage Duration</b>	initial, reauth: 12 months
<b>Other Criteria</b>	IPF (reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

87

## EUCRISA (S)

---

### Products Affected

- Eucrisa

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Atopic dermatitis (initial): Diagnosis of mild to moderate atopic dermatitis. Trial and failure, contraindication, or intolerance to one prescription strength topical corticosteroid, unless the affected area is sensitive (i.e., face, axillae, groin).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Reauth: Documentation of a positive clinical response to therapy (e.g., reduction in body surface area involvement, reduction in pruritus severity).

Last Updated: June 2021

Formulary ID: 21381, Version 14

88



## **EVRYSDI (S)**

---

### **Products Affected**

- Evryski

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Spinal muscular atrophy (SMA) (initial): Diagnosis of spinal muscular atrophy (SMA) type I, II, or III. Both of the following: a) The mutation or deletion of genes in chromosome 5q resulting in one of the following: 1) Homozygous gene deletion or mutation (e.g., homozygous deletion of exon 7 at locus 5q13) or 2) Compound heterozygous mutation (e.g., deletion of SMN1 exon 7 [allele 1] and mutation of SMN1 [allele 2]) AND b) Patient has at least 2 copies of SMN2. Patient is not dependent on both of the following: 1) Invasive ventilation or tracheostomy and 2) Use of non-invasive ventilation beyond use for naps and nighttime sleep. At least one of the following exams (based on patient age and motor ability) has been conducted to establish baseline motor ability: Hammersmith Infant Neurological Exam (HINE) (infant to early childhood), Hammersmith Functional Motor Scale Expanded (HFMSE), Upper Limb Module (ULM) Test (Non ambulatory), or Children's Hospital of Philadelphia Infant Test of Neuromuscular Disorders (CHOP INTEND), or Motor Function Measure 32 (MFM-32) Scale. Patient is not to receive concomitant chronic survival motor neuron (SMN) modifying therapy for the treatment of SMA (e.g., Spinraza). One of the following: a) patient has not previously received gene replacement therapy for the treatment of SMA (e.g., Zolgensma) or b) patient has previously received gene therapy for the treatment of SMA (e.g., Zolgensma) AND provider attests that there has been an inadequate response to gene therapy (e.g., sustained decrease in at least one motor test score over a period of 6 months).
<b>Age Restrictions</b>	Initial: Patient is at least 2 months of age or older
<b>Prescriber Restrictions</b>	SMA (Initial, Reauth): Prescribed by or in consultation with a neurologist with expertise in the diagnosis and treatment of SMA
<b>Coverage Duration</b>	Initial, Reauth: 12 months
<b>Other Criteria</b>	SMA (Reauth): Documentation of positive clinical response to therapy from pretreatment baseline status as demonstrated by the most recent results from one of the following exams: A) One of the following HINE-2 milestones: 1) Improvement or maintenance of previous improvement of

Last Updated: June 2021

at least a 2 point (or maximal score) increase in ability to kick OR 2) Improvement or maintenance of a previous improvement of at least a 1 point increase in any other HINE-2 milestone (e.g., head control, rolling, sitting, crawling, etc.), excluding voluntary grasp OR 3) Patient exhibited improvement, or maintenance of previous improvement in more HINE motor milestones than worsening, from pretreatment baseline (net positive improvement) OR 4) Patient has achieved and maintained any new motor milestones when they would otherwise be unexpected to do so (e.g., sit unassisted, stand, walk) OR B) One of the following HFMSE milestones: 1) Improvement or maintenance of a previous improvement of at least a 3 point increase in score from pretreatment baseline OR 2) Patient has achieved and maintained any new motor milestone from pretreatment baseline when they would otherwise be unexpected to do so (e.g., sit unassisted, stand, walk) OR C) One of the following ULM test milestones: 1) Improvement or maintenance of a previous improvement of at least a 2 point increase in score from pretreatment baseline OR 2) Patient has achieved and maintained any new motor milestone from pretreatment baseline when they would otherwise be unexpected to do so (e.g., sit unassisted, stand, walk) OR D) One of the following CHOP INTEND milestones: 1) Improvement or maintenance of a previous improvement of at least a 4 point increase in score from pretreatment baseline OR 2) Patient has achieved and maintained any new motor milestone from pretreatment baseline when they would otherwise be unexpected to do so (e.g., sit unassisted, stand, walk) OR E) One of the following MFM-32 milestones: 1) Improvement or maintenance of a previous improvement of at least a 3 point increase in score from pretreatment baseline OR 2) Patient has achieved and maintained any new motor milestone from pretreatment baseline when they would otherwise be unexpected to do so (e.g., sit unassisted, stand, walk). Patient continues to not be dependent on both of the following: 1) Invasive ventilation or tracheostomy AND 2) use of non-invasive ventilation beyond use for naps and nighttime sleep. Patient is not to receive concomitant chronic survival motor neuron (SMN) modifying therapy for the treatment of SMA (e.g., Spinraza). One of the following: a) patient has not previously received gene replacement therapy for the treatment of SMA (e.g., Zolgensma) or b) patient has previously received gene therapy for the treatment of SMA (e.g., Zolgensma) AND provider attests that there has been an inadequate response to gene therapy (e.g., sustained decrease in at least one motor test score over a period of 6 months).

## FABRAZYME (S)

---

### Products Affected

- Fabrazyme INJ 35MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Fabry Disease: Diagnosis of Fabry disease. Fabrazyme will not be used in combination with Galafold (migalastat).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Fabry Disease: 12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

92

## FARYDAK (S)

---

### Products Affected

- Farydak

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple Myeloma (MM): Diagnosis of MM. Used in combination with both of the following: Velcade (bortezomib) and dexamethasone. Patient has received at least two prior treatment regimens which included both of the following: Velcade (bortezomib) and an immunomodulatory agent [eg, Revlimid (lenalidomide), Thalomid (thalidomide)].
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist/hematologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

93

# FASENRA (S)

---

## Products Affected

- Fasenra
- Fasenra Pen

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Asthma (initial): Diagnosis of severe asthma. Asthma is an eosinophilic phenotype as defined by a baseline (pre-treatment) peripheral blood eosinophil level greater than or equal to 150 cells per microliter. One of the following: 1) Patient has had at least one or more asthma exacerbations requiring systemic corticosteroids within the past 12 months, OR 2) Any prior intubation for an asthma exacerbation, OR 3) Prior asthma-related hospitalization within the past 12 months. Patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications: a) Both of the following: i) High-dose inhaled corticosteroid (ICS) [e.g., greater than 500 mcg fluticasone propionate equivalent/day] and ii) additional asthma controller medication [e.g., leukotriene receptor antagonist, long-acting beta-2 agonist (LABA), tiotropium], OR b) One maximally-dosed combination ICS/LABA product [e.g., Advair (fluticasone propionate/salmeterol), Dulera (mometasone/formoterol), Symbicort (budesonide/formoterol), Breo Ellipta (fluticasone/vilanterol)].
<b>Age Restrictions</b>	Asthma (Initial): Patient is 12 years of age or older
<b>Prescriber Restrictions</b>	Asthma (Initial/Reauth): Prescribed by or in consultation with a pulmonologist or allergist/immunologist
<b>Coverage Duration</b>	Asthma (init): 6 months. Asthma (reauth): 12 months
<b>Other Criteria</b>	Asthma (Reauth): Documentation of positive clinical response to therapy (e.g., reduction in exacerbations, improvement in forced expiratory volume in 1 second [FEV1], decreased use of rescue medications). Patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications: a) Both of the following: i) inhaled corticosteroid (ICS) (e.g., fluticasone, budesonide) and ii) additional asthma controller medication [e.g., leukotriene receptor antagonist, long-acting beta-2 agonist (LABA), tiotropium], OR b) A combination ICS/LABA product [e.g., Advair (fluticasone propionate/salmeterol), Dulera (mometasone/formoterol), Symbicort (budesonide/formoterol), Breo Ellipta (fluticasone/vilanterol)].

Last Updated: June 2021

Last Updated: June 2021

Formulary ID: 21381, Version 14



# FENTANYL (S)

---

## Products Affected

- Fentanyl Citrate Oral Transmucosal

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For the management of breakthrough cancer pain. Patient is currently taking a long-acting opioid around the clock for cancer pain. Patient must have at least a one week history of ONE of the following medications to demonstrate tolerance to opioids: Morphine sulfate at doses of greater than or equal to 60 mg/day, Fentanyl transdermal patch at doses greater than or equal to 25 µg/hr, Oxycodone at a dose of greater than or equal to 30 mg/day, Oral hydromorphone at a dose of greater than or equal to 8 mg/day, Oral oxymorphone at a dose of greater than or equal to 25 mg/day, or an alternative opioid at an equianalgesic dose (e.g., oral methadone greater than or equal to 20 mg/day).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with one of the following: Pain specialist, Oncologist, Hematologist, Hospice care specialist, or Palliative care specialist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

97

# FERRIPROX (S)

---

## Products Affected

- Deferiprone
- Ferriprox
- Ferriprox Twice-a-day

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Transfusional iron overload due to thalassemia syndromes (Initial): Diagnosis of transfusional iron overload due to thalassemia syndromes. Absolute neutrophil count (ANC) greater than $1.5 \times 10^9/L$ . One of the following: A) Trial and failure, defined as serum ferritin greater than 2,500 mcg/L, to one chelation therapy (i.e., deferoxamine, deferasirox) OR B) History of contraindication or intolerance to one chelation therapy (i.e., deferoxamine, deferasirox).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	All uses (reauth): Patient has experienced greater than or equal to 20% decline in serum ferritin levels from baseline. ANC greater than $1.5 \times 10^9/L$ .

Last Updated: June 2021

Formulary ID: 21381, Version 14

98

## FINTEPLA (S)

---

### Products Affected

- Fintepla

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of seizures associated with Dravet syndrome
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

## FIRAZYR (S)

---

### Products Affected

- Icatibant Acetate

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Treatment of hereditary angioedema (HAE) attacks: Diagnosis of HAE. For the treatment of acute HAE attacks. Not used in combination with other approved treatments for acute HAE attacks.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	HAE: Prescribed by or in consultation with an immunologist, allergist, or rheumatologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

100

## FIRDAPSE (S)

---

### Products Affected

- Firdapse

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Lambert-Eaton Myasthenic Syndrome (LEMS) (initial): Diagnosis of LEMS.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	LEMS (initial): Prescribed by or in consultation with a neurologist.
<b>Coverage Duration</b>	LEMS (initial): 3 months. LEMS (reauth): 12 months.
<b>Other Criteria</b>	LEMS (reauth): Documentation of positive clinical response to therapy (e.g., improvement in dynamometry, Timed 25-Foot Walk Test, Timed Up and Go Test).

Last Updated: June 2021

Formulary ID: 21381, Version 14

101

## FIRMAGON (S)

---

### Products Affected

- Firmagon INJ 120MG/VIAL, 80MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of advanced or metastatic prostate cancer.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

102

## FOTIVDA (S)

---

### Products Affected

- Fotivda

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of renal cell carcinoma. Disease is one of the following: relapsed or refractory. Patient has received two or more prior systemic therapies (e.g., cabozantinib + nivolumab, lenvatinib + pembrolizumab, etc.).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist/nephrologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

103

## GALAFOLD (S)

---

### Products Affected

- Galafold

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Fabry Disease (FD) (initial): Diagnosis of FD. Patient has an amenable galactosidase alpha gene (GLA) variant based on in vitro assay data. FD (initial, reauthorization): Will not be used in combination with Fabrazyme (agalsidase beta).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	FD (initial, reauth): 12 months.
<b>Other Criteria</b>	FD (reauthorization): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

104



## GAMASTAN S/D (S)

---

### Products Affected

- Gamastan

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	Contraindications to immune globulin therapy (i.e., IgA deficiency with antibodies to IgA and a history of hypersensitivity or product specific contraindication).
<b>Required Medical Information</b>	Immune globulin is being used intramuscularly. The immune globulin will be administered at the minimum effective dose and appropriate frequency for the prescribed diagnosis. Patient requires immunization for hepatitis A, measles, rubella, or varicella.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	3 months (Approve one dose only)
<b>Other Criteria</b>	Subject to Part B vs D review.

Last Updated: June 2021

Formulary ID: 21381, Version 14

105

## GAMIFANT (S)

---

### Products Affected

- Gamifant INJ 100MG/20ML

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Primary Hemophagocytic Lymphohistiocytosis (HLH) (initial): Diagnosis of HLH. One of the following: 1) Disease is refractory, recurrent, or progressive, or 2) Trial and failure, contraindication, or intolerance to conventional HLH therapy (e.g., etoposide, dexamethasone, cyclosporine A, intrathecal methotrexate). HLH (initial, reauth): Patient has not received hematopoietic stem cell transplantation (HSCT).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	HLH (initial): Prescribed by or in consultation with a hematologist/oncologist.
<b>Coverage Duration</b>	HLH (initial, reauth): 6 months.
<b>Other Criteria</b>	HLH (reauth): Documentation of positive clinical response to therapy (e.g., improvement in hemoglobin/lymphocyte/platelet counts, afebrile, normalization of inflammatory factors/markers).

Last Updated: June 2021

## GATTEX (S)

---

### Products Affected

- Gattex

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Short Bowel Syndrome (SBS) (Initial): Diagnosis of SBS. Submission of medical records (e.g., chart notes, laboratory values) documenting that the patient is dependent on parenteral nutrition/intravenous (PN/IV) support for at least 12 months.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	SBS (Init, reauth): Prescribed by or in consultation with a gastroenterologist.
<b>Coverage Duration</b>	SBS (Init): 6 months. SBS (Reauth): 12 months.
<b>Other Criteria</b>	SBS (Reauth): Submission of medical records (e.g., chart notes, laboratory values) documenting that the patient has had a reduction in weekly parenteral nutrition/intravenous (PN/IV) support from baseline while on therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

107

# GAVRETO (S)

---

## Products Affected

- Gavreto

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Non-small cell lung cancer (NSCLC): Diagnosis of NSCLC. Presence of rearranged during transfection (RET) gene fusion-positive tumor(s). Medullary Thyroid Cancer (MTC): Diagnosis of medullary thyroid cancer (MTC). Disease is one of the following: advanced or metastatic. Disease has presence of rearranged during transfection (RET) gene mutation tumor(s). Disease requires treatment with systemic therapy. Thyroid Cancer: Diagnosis of thyroid cancer. Disease is one of the following: advanced or metastatic. Disease has presence of rearranged during transfection (RET) gene fusion-positive tumor(s). Disease requires treatment with systemic therapy. One of the following: patient is radioactive iodine-refractory or radioactive iodine therapy is not appropriate.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	NSCLC, MTC: Prescribed by or in consultation with an oncologist. Thyroid Cancer: Prescribed by or in consultation with an endocrinologist or an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy

Last Updated: June 2021

## GILENYA (S)

---

### Products Affected

- Gilenya

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple Sclerosis (MS): Diagnosis of a relapsing form of MS (eg, clinically isolated syndrome, relapsing-remitting disease, secondary progressive disease, including active disease with new brain lesions).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

109

## GILOTRIF (S)

---

### Products Affected

- Gilotrif

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Non-small cell lung cancer (NSCLC): A) Diagnosis of advanced or metastatic (stage IIIB or IV) NSCLC AND B) One of the following: 1) Both of the following: a) Tumors have non-resistant epidermal growth factor (EGFR) mutations as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) AND b) GILOTRIF will be used as first-line treatment, OR 2) All of the following: a) disease progressed after platinum-based chemotherapy and b) squamous NSCLC.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

## GIVLAARI (S)

---

### Products Affected

- Givlaari

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Initial: Diagnosis of acute hepatic porphyria (i.e., acute intermittent porphyria, hereditary coproporphyria, variegate porphyria, ALA dehydrase deficient porphyria). Patient has active disease with at least two documented porphyria attacks within the past 6 months. Provider attestation documenting elevated urinary or plasma levels of one of the following within the past 12 months: 1) porphobilinogen (PBG) or 2) delta-aminolevulinic acid (ALA). Patient has not had a liver transplant.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Initial, Reauth: Prescribed by or in consultation with a gastroenterologist or a specialist with expertise in the diagnosis and management of acute hepatic porphyria.
<b>Coverage Duration</b>	Initial: 6 months. Reauth: 12 months.
<b>Other Criteria</b>	Reauth: Documentation of positive clinical response while on therapy as demonstrated by both of the following: 1) Reduction in hemin administration requirements and 2) Reduction in the rate or number of porphyria attacks. Patient has not had a liver transplant.

Last Updated: June 2021

Formulary ID: 21381, Version 14

111

## GLATIRAMER ACETATE (S)

---

### Products Affected

- Glatiramer Acetate
- Glatopa

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple Sclerosis (MS): Diagnosis of a relapsing form of MS (eg, clinically isolated syndrome, relapsing-remitting disease, secondary progressive disease, including active disease with new brain lesions).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

112



## GLEEVEC (S)

### Products Affected

- Imatinib Mesylate

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	One of the following: A) Diagnosis of Philadelphia chromosome positive (Ph+)/BCR ABL-positive chronic myelogenous leukemia (CML) OR B) Ph+/BCR ABL+ acute lymphoblastic leukemia (ALL) OR C) Gastrointestinal stromal tumor (GIST) AND one of the following: 1) Patient has documented c-KIT (CD117) positive unresectable or metastatic malignant GIST, OR 2) Patient had resection of c-KIT (CD117) positive GIST and imatinib will be used as adjuvant therapy OR D) Dermatofibrosarcoma protuberans that is unresectable, recurrent, or metastatic OR E) Hypereosinophilic syndrome or chronic eosinophilic leukemia OR F) Myelodysplastic syndrome (MDS) or myeloproliferative disease associated with platelet-derived growth factor receptor gene rearrangements OR G) Aggressive systemic mastocytosis without the D816V c-KIT mutation or with c-KIT mutational status unknown.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses: Prescribed by or in consultation with an oncologist or hematologist
<b>Coverage Duration</b>	All uses: 12 months
<b>Other Criteria</b>	All uses: Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

113

# **GROWTH HORMONE, PREFERRED (S)**

---

## **Products Affected**

- Genotropin
- Genotropin Miniquick

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	<p>PGHD(initial):less than 4mo w/suspected GD based on clinical presentation (eg, persistent neonatal hypoglycemia, persistent/prolonged neonatal jaundice/elev bilirubin, male infant with microgenitalia, midline anatomical defects, failure to thrive),OR hx neonatal hypoglycemia assoc w/pituitary dz,or panhypopituitarism dx,or all of the following: PGHD dx [confrmd by ht (utilizing age and gender grwth charts related to ht) documented(doc) by ht more than 2.0SD below midparental ht or more than 2.25SD below population(pop) mean (below 1.2 percentile for age and gender),or grwth velocity more than 2SD below mean for age and gender, or delayed skeletal maturation more than 2SD below mean for age and gender (eg,delayed more than 2yrs compared w/chronological age)].</p> <p>PWS(reauth):evidence of positive response to tx(eg,incr in total LBM, decr in fat mass) and expctd adult ht not attained and doc of expctd adult ht goal. GFSGA(initial):SGA dx based on catchup grwth failure in 1st 24mo of life using 0-36mo grwth chart confrmd by birth wt or length below 3rd percentile for gestational age(more than 2SD below pop mean) and ht remains at or below 3rd percentile (more than 2SD below pop mean). TS,NS(initial):ped grwth failure dx assoc w/TS w/doc female w/bone age less than 14yrs, or NS and ht below 5th percentile on grwth charts for age and gender. SHOX(initial):ped grwth failure dx w/SHOX gene deficiency confirmed by genetic testing. GFCRI(initial): ped grwth failure dx assoc w/CRI. ISS(initial):ISS dx, diagnostic eval excluded other causes assoc w/short stature(eg GHD, chronic renal insufficiency), doc ht at or below -2.25SD score below corresponding mean ht for age and gender assoc with growth rates unlikely to permit attainment of adult height in the normal range. PGHD,NS,SHOX,GFCRI,ISS (initial): doc male w/bone age less than 16yrs or female w/bone age less than 14yrs. PGHD,GFSGA,TS/NS,SHOX,GFCRI,ISS(reauth):expctd adult ht not attained and doc of expctd adult ht goal.</p>
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	PGHD, PWS, GFSGA, TS/NS, SHOX, AGHD, TPAP, IGHDA, ISS: prescribed by endocrinologist. GFCRI: prescribed by endocrinologist or nephrologist

Last Updated: June 2021

<b>Coverage Duration</b>	All uses (initial, reauth): 12 months
<b>Other Criteria</b>	<p>AGHD(initial):dx of AGHD with clin records supporting dx of childhood-onset GHD, or adult-onset GHD w/clin records doc hormone deficiency d/t hypothalamic-pituitary dz from organic or known causes (eg,damage from surgery, cranial irradiation, head trauma, subarachnoid hemorrhage) and pt has 1GH stim test (insulin tolerance test [ITT], arginine/GHRH,glucagon,arginine) to confirm adult GHD w/peak GH values ([ITT at or below 5mcg/L],[GHRH+ARG at or below 11mcg/L if BMI less than 25kg/m2, at or below 8mcg/L if BMI at or above 25 and below 30kg/m2, or at or below 4mcg/L if BMI at or above 30kg/m2],[glucagon at or below 3mcg/L],[Arg at or below 0.4mcg/L],[macimorelin less than 2.8 ng/mL 30, 45, 60 and 90 mins following macimorelin administration]) or doc deficiency of 3 anterior pituitary hormones (prolactin,ACTH,TSH,FSH/LH) and IGF-1/somatomedinC below age and gender adjstd nrml range as provided by physicians lab. AGHD,IGHDA(reauth):monitoring as demonstrated by doc w/in past 12mo of IGF-1/somatomedinC level.</p> <p>TransitionPhaseAdolescent Pts(TPAP)(initial): attained expctd adult ht or closed epiphyses on bone radiograph, and doc high risk of GHD d/t GHD in childhood (from embryopathic/congenital defects, genetic mutations, irreversible structural hypothalamic-pituitary dz, panhypopituitarism, or deficiency of 3 anterior pituitary hormones: ACTH,TSH,prolactin,FSH/LH), w/IGF-1/somatomedinC below age and gender adj nrml range as provided by physicians lab, or pt does not have low IGF-1/somatomedinC and d/c GH tx for at least 1mo, and pt has 1 GH stim test (ITT,GHRH+ARG,ARG,glucagon) after d/c of tx for at least 1mo w/peak GH value [ITT at or below 5mcg/L], [GHRH+ARG at or below 11mcg/L if BMI less than 25kg/m2, at or below 8mcg/L if BMI at or above 25 and below 30kg/m2, or at or below 4mcg/L if BMI at or above 30kg/m2], [glucagon at or below 3mcg/L], [Arg at or below 0.4mcg/L], or at low risk of severe GHD(eg d/t isolated and/or idiopathic GHD) and d/c GH tx for at least 1mo, and pt has 1 GH stim test (ITT, GHRH+ARG, ARG, glucagon) after d/c of tx for at least 1mo w/corresponding peak GH value [ITT at or below 5mcg/L], [GHRH+ARG at or below 11mcg/L if BMI less than 25kg/m2, at or below 8mcg/L if BMI at or above 25 and below 30kg/m2, or at or below 4mcg/L if BMI at or above 30kg/m2], [glucagon at or below 3mcg/L], [Arg at or below 0.4mcg/L]. TPAP(reauth): evidence of positive response to therapy (eg,incr in total lean body mass, exercise capacity or IGF-1 and IGFBP-3). IGHDA(initial):doc GHD by failure to produce peak serum GH greater than 5 mcg/L after 2 provocative pharmacol stim tests(insulin,L-ARG,glucagon).</p>

Last Updated: June 2021

## HAEGARDA (S)

---

### Products Affected

- Haegarda

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Prophylaxis of hereditary angioedema (HAE) attacks: Diagnosis of HAE. For prophylaxis against HAE attacks.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	HAE: Prescribed by or in consultation with an immunologist, allergist, or rheumatologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

117

# HERCEPTIN (S)

---

## Products Affected

- Herceptin INJ 150MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Breast cancer: Diagnosis of human epidermal growth factor receptor 2 (HER2)-overexpressing breast cancer. One of the following treatment regimens: a) As adjuvant treatment, b) metastatic disease and one of the following: 1) used in combination with a taxane (eg, docetaxel, paclitaxel), or 2) used as a single agent in a patient who has received one or more chemotherapy regimens for metastatic disease, or c) used in combination with Perjeta (pertuzumab). Gastric Cancer: Diagnosis of HER2-overexpressing gastric or gastroesophageal junction adenocarcinoma (locally advanced, recurrent, or metastatic). Used in combination with one of the following treatment regimens: a) Platinol (cisplatin) and Adrucil (5-fluorouracil), or b) Platinol (cisplatin) and Xeloda (capecitabine).
Age Restrictions	N/A
Prescriber Restrictions	All uses: Prescribed by or in consultation with an oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

Last Updated: June 2021

# HETLIOZ (S)

---

## Products Affected

- Hetlitz

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Non-24-Hour Sleep-Wake Disorder (Non-24) (initial): Both of the following: 1) Diagnosis of non-24-hour sleep-wake disorder (also known as free-running disorder, free-running or non-entrained type circadian rhythm sleep disorder, or hypernycthemeral syndrome), AND 2) patient is totally blind (has no light perception). Smith-Magenis Syndrome (SMS) (initial): Diagnosis of Smith-Magenis Syndrome (SMS). Patient is experiencing nighttime sleep disturbances (i.e., difficulty falling asleep, frequent nighttime waking and early waking).
<b>Age Restrictions</b>	SMS (initial): 16 years of age or older
<b>Prescriber Restrictions</b>	Non-24 (initial): Prescribed by or in consultation with a specialist in sleep disorders or neurologist. SMS (initial): Prescribed by or in consultation with a specialist in sleep disorders.
<b>Coverage Duration</b>	Non-24, SMS (initial): 6 mo. (reauth): 12 mo
<b>Other Criteria</b>	Non-24 (reauth): Documentation of positive clinical response to therapy. SMS (reauth): Documentation of positive clinical response to therapy (i.e., improvement in nighttime total sleep time, improvement in nighttime sleep quality)

Last Updated: June 2021

## HETLIOZ LQ (S)

---

### Products Affected

- Hetlioz Lq

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Smith-Magenis Syndrome (SMS) (initial): Diagnosis of Smith-Magenis Syndrome (SMS). Patient is experiencing nighttime sleep disturbances (i.e., difficulty falling asleep, frequent nighttime waking and early waking).
<b>Age Restrictions</b>	SMS (initial): Patient is 3 through 15 years of age
<b>Prescriber Restrictions</b>	SMS (initial): Prescribed by or in consultation with a specialist in sleep disorders or neurologist.
<b>Coverage Duration</b>	SMS (initial): 6 mo. (reauth): 12 mo
<b>Other Criteria</b>	SMS (reauth): Documentation of positive clinical response to therapy (i.e., improvement in nighttime total sleep time, improvement in nighttime sleep quality).

Last Updated: June 2021

Formulary ID: 21381, Version 14

120



## HRM - ANTIPSYCHOTICS

---

### Products Affected

- Thioridazine Hcl TABS 100MG, 10MG, 25MG, 50MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The drug is being used for an FDA-approved diagnosis AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication. Trial and failure, contraindication or intolerance to one of the following: haloperidol, fluphenazine, or an atypical antipsychotic.
<b>Age Restrictions</b>	PA applies to patients 65 years or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Applies to New Starts only.

Last Updated: June 2021

Formulary ID: 21381, Version 14

121

## HRM - ENDOCRINE

---

### Products Affected

- Megestrol Acetate SUSP  
40MG/ML, 625MG/5ML
- Megestrol Acetate TABS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication.
<b>Age Restrictions</b>	PA applies to patients 65 years or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Applies to New Starts only.

Last Updated: June 2021

Formulary ID: 21381, Version 14

122

# HRM - PHENOBARBITAL, PENTOBARBITAL

---

## Products Affected

- Phenobarbital ELIX 20MG/5ML
- Phenobarbital TABS 100MG, 15MG, 16.2MG, 30MG, 32.4MG, 60MG, 64.8MG, 97.2MG
- Phenobarbital Sodium INJ 130MG/ML, 65MG/ML

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The drug is being used for an FDA-approved diagnosis AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication.
<b>Age Restrictions</b>	PA applies to patients 65 years or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Applies to New Starts only.

Last Updated: June 2021

Formulary ID: 21381, Version 14

123

## HRM - SKELETAL MUSCLE RELAXANTS

---

### Products Affected

- Chlorzoxazone TABS 500MG
- Cyclobenzaprine Hydrochloride  
TABS 10MG, 5MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication.
<b>Age Restrictions</b>	PA applies to patients 65 years or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

124

# HRM - TCA

---

## Products Affected

- Amitriptyline Hcl TABS 100MG, 150MG, 25MG, 75MG
- Amitriptyline Hydrochloride TABS 10MG, 50MG
- Doxepin Hcl CAPS 100MG, 10MG, 150MG, 50MG, 75MG
- Doxepin Hcl CONC
- Doxepin Hydrochloride CAPS 25MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication .
Age Restrictions	PA applies to patients 65 years or older
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Applies to New Starts only.

Last Updated: June 2021

Formulary ID: 21381, Version 14

125

# HUMIRA (S)

---

## **Products Affected**

- Humira
- Humira Pediatric Crohns Disease Starter Pack
- Humira Pen
- Humira Pen-cd/uc/hs Starter
- Humira Pen-pediatric Uc Starter Pack
- Humira Pen-ps/uv Starter

Last Updated: June 2021

Formulary ID: 21381, Version 14

126

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. Trial and failure, contraindication, or intolerance (TF/C/I) to one disease modifying antirheumatic drug (DMARD) [eg, methotrexate (Rheumatrex/Trexall), Arava (leflunomide), Azulfidine (sulfasalazine)]. Juvenile Idiopathic Arthritis (JIA) (Initial): Diagnosis of moderately to severely active polyarticular JIA. TF/C/I to one of the following DMARDs: Arava (leflunomide) or methotrexate (Rheumatrex/Trexall). Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. Plaque psoriasis (Initial): Diagnosis of moderate to severe chronic plaque psoriasis. Ankylosing Spondylitis (AS) (Initial): Diagnosis of active AS. TF/C/I to two NSAIDs (e.g., diclofenac, ibuprofen, meloxicam, naproxen). Crohn's Disease (CD) (Initial): Diagnosis of moderately to severely active CD. TF/C/I to one of the following conventional therapies: 6-mercaptopurine, Imuran (azathioprine), corticosteroid (eg, prednisone, methylprednisolone), methotrexate (Rheumatrex/Trexall), or failure (ie, lost response) or intolerance to infliximab. Ulcerative Colitis (UC) (Initial): Diagnosis of moderately to severely active UC. TF/C/I to one of the following conventional therapies: 6-mercaptopurine, Imuran (azathioprine), corticosteroid (eg, prednisone, methylprednisolone), aminosalicylate [eg, mesalamine (Asacol, Pentasa, Rowasa), Dipentum (olsalazine), Azulfidine (sulfasalazine)]. Hidradenitis suppurativa (Initial): Diagnosis of moderate to severe hidradenitis suppurativa (ie, Hurley Stage II or III). Uveitis (initial): Diagnosis of non-infectious uveitis, classified as intermediate, posterior, or panuveitis.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	RA, AS, JIA (initial): Prescribed by or in consultation with a rheumatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist. Plaque Psoriasis, HS (initial): Prescribed by or in consultation with a dermatologist. CD, UC (initial): Prescribed by or in consultation with a gastroenterologist. Uveitis (initial): Prescribed by or in consultation with an ophthalmologist or rheumatologist.
<b>Coverage Duration</b>	UC (Initial): 12 wks. UC (reauth): 12 mo. All other indications (initial, reauth): 12 mo.

Last Updated: June 2021

<b>Other Criteria</b>	RA, JIA, PsA, AS, CD, Hidradenitis suppurativa (HS), Uveitis (Reauth): Documentation of positive clinical response to therapy. Plaque psoriasis (Reauth): Documentation of positive clinical response to therapy as evidenced by one of the following: reduction in the body surface area (BSA) involvement from baseline, OR improvement in symptoms (eg, pruritus, inflammation) from baseline. UC (Reauth): For patients who initiated therapy within the past 12 weeks: Documentation of clinical remission or significant clinical benefit by eight weeks (Day 57) of therapy OR For patients who have been maintained on therapy for longer than 12 weeks: Documentation of positive clinical response to therapy.
-----------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------



## IBRANCE (S)

---

### Products Affected

- Ibrance

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Breast Cancer: Diagnosis of advanced or metastatic breast cancer. Disease is a) hormone receptor (HR)-positive, and b) human epidermal growth factor receptor 2 (HER2)-negative. One of the following: 1) used in combination with an aromatase inhibitor (e.g., anastrozole, letrozole, exemestane) and one of the following: a) patient is a male, or b) patient is a postmenopausal woman, OR 2) both of the following: used in combination with Faslodex (fulvestrant) and disease has progressed following endocrine therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

129

## ICLUSIG (S)

---

### Products Affected

- Iclusig

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chronic myelogenous leukemia: Diagnosis of chronic myelogenous leukemia AND One of the following: A) Trial and failure, contraindication, or intolerance to at least TWO other tyrosine kinase inhibitors (e.g., Gleevec [imatinib], Sprycel, Tasigna, and Bosulif) or B) Patient has the T315I mutation. Acute Lymphoblastic Leukemia: Diagnosis of Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph+ ALL) AND One of the following: A) Trial and failure, resistance relapse, contraindication, or intolerance to at least TWO other FDA-approved tyrosine kinase inhibitors (e.g., Gleevec [imatinib], Sprycel), or B) Patient has the T315I mutation.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses: Prescribed by or in consultation with an oncologist or hematologist
<b>Coverage Duration</b>	All uses: 12 months
<b>Other Criteria</b>	All uses: Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

130

## IDHIFA (S)

---

### Products Affected

- Idhifa

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Acute Myeloid Leukemia (AML): Diagnosis of AML. Disease is relapsed or refractory. Patient has an isocitrate dehydrogenase-2 (IDH2) mutation as detected by a U.S. Food and Drug Administration (FDA)-approved test (e.g., Abbott RealTime IDH2 assay) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist/oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

131

## ILUMYA (S)

---

### Products Affected

- Ilumya

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Plaque Psoriasis (initial): Diagnosis of moderate-to-severe plaque psoriasis. One of the following: set A) Both of the following: 1) Trial and failure, contraindication, or intolerance to one of the following: Enbrel (etanercept), Humira (adalimumab), Skyrizi (risankizumab), AND 2) Trial and failure, contraindication, or intolerance to Cosentyx (secukinumab), or set B) For continuation of prior therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Plaque Psoriasis (initial): Prescribed by or in consultation with a dermatologist
<b>Coverage Duration</b>	Initial, reauth: 12 months
<b>Other Criteria</b>	Plaque Psoriasis (Reauth): Documentation of positive clinical response to therapy as evidenced by one of the following: reduction in the body surface area (BSA) involvement from baseline, OR improvement in symptoms (eg, pruritus, inflammation) from baseline.

Last Updated: June 2021

Formulary ID: 21381, Version 14

132

## IMBRUVICA (S)

---

### Products Affected

- Imbruvica

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Mantle cell lymphoma (MCL): Diagnosis of MCL AND patient has relapsed or is refractory to at least one prior therapy for the treatment of MCL. Chronic lymphocytic leukemia (CLL): Diagnosis of CLL. Waldenstrom's macroglobulinemia: Diagnosis of Waldenstrom's macroglobulinemia/lymphoplasmacytic lymphoma. Small lymphocytic lymphoma (SLL): Diagnosis of SLL. Marginal zone lymphoma (MZL): Diagnosis of MZL AND patient has received at least one prior anti-CD20-based therapy for MZL [e.g., Rituxan (rituximab), Zevalin (ibritumomab), Gazyva (obinutuzumab, etc.)]. Chronic graft versus host disease (cGVHD): Diagnosis of cGVHD AND trial and failure of one or more lines of systemic therapy (e.g., corticosteroids, mycophenolate).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses (except chronic graft versus host disease): Prescribed by or in consultation with an oncologist or hematologist. Chronic graft versus host disease: Prescribed by or in consultation with a hematologist, oncologist, or physician experienced in the management of transplant patients.
<b>Coverage Duration</b>	All Uses: 12 months
<b>Other Criteria</b>	All Uses: Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

133

## IMFINZI (S)

---

### Products Affected

- Imfinzi

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Urothelial carcinoma: 1) Diagnosis of locally advanced or metastatic urothelial carcinoma AND 2) One of the following: a) Patient has experienced disease progression during or following platinum-containing chemotherapy OR b) Patient has experienced disease progression within 12 months of neoadjuvant or adjuvant treatment with platinum-containing chemotherapy. Non-Small Cell Lung Cancer (NSCLC): 1) Diagnosis of NSCLC AND 2) Disease is stage III and unresectable AND 3) Disease has not progressed following concurrent platinum-based chemotherapy and radiation therapy. Small Cell Lung Cancer: 1) Diagnosis of extensive-stage small cell lung cancer (ES-SCLC) AND 2) Used as first line treatment AND 3) Both of the following: a) Used in combination with Etoposide and b) Used in combination with carboplatin or cisplatin.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

134

## INCRELEX (S)

---

### Products Affected

- Increlex

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Insulin-like Growth Factor-1 (IGF-1) deficiency (initial): Diagnosis of severe primary IGF-1 deficiency. Height standard deviation score of -3.0 or less. Basal IGF-1 standard deviation score of -3.0 or less. Normal or elevated growth hormone (GH). GH gene deletion (initial): Diagnosis of GH gene deletion in patients who have developed neutralizing antibodies to GH.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Initial: Prescribed by or in consultation with a pediatric endocrinologist
<b>Coverage Duration</b>	Initial, reauth: 12 months
<b>Other Criteria</b>	(Reauth): Evidence of positive response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

135

# INFLECTRA (S)

---

## Products Affected

- Inflectra



<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Crohn's Disease (CD) and Fistulizing Crohn's Disease (FCD) (initial): Dx of moderately to severely active CD or FCD. Ulcerative colitis (UC) (initial): Dx of moderately to severely active UC. Rheumatoid arthritis (RA) (initial): Diagnosis (Dx) of moderately to severely active RA. Ankylosing spondylitis (AS) (initial): Dx of active AS. Psoriatic arthritis (PsA) (initial): Dx of active PsA. Plaque psoriasis (initial): Dx of chronic severe (ie, extensive and/or disabling) plaque psoriasis. CD, FCD (initial): Trial and failure, contraindication or intolerance (TF/C/I) to one of the following conventional therapies: 6-mercaptopurine (Purinethol), azathioprine (Imuran), corticosteroids (eg, prednisone, methylprednisolone), methotrexate (Rheumatrex, Trexall). UC (initial): TF/C/I to one of the following conventional therapies: corticosteroids, aminosalicylate [eg, mesalamine (Asacol/Pentasa/Rowasa), olsalazine (Dipentum), sulfasalazine (Azulfidine/Sulfazine)], azathioprine (Imuran), 6-mercaptopurine (Purinethol). RA (initial): Receiving concurrent therapy with methotrexate (Rheumatrex/Trexall), or TF/C/I to methotrexate. AS (initial): TF/C/I to two or more NSAIDs (e.g., diclofenac, ibuprofen, meloxicam, naproxen). All indications (initial): One of the following: 1) Trial and failure or intolerance to Remicade or 2) For continuation of prior Inflectra therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	RA, AS: Prescribed or recommended by a rheumatologist. PsA: Prescribed or recommended by rheumatologist or dermatologist. Crohn's Disease, Fistulizing Crohn's Disease, UC: Prescribed or recommended by a gastroenterologist. Plaque Psoriasis: Prescribed or recommended by a dermatologist.
<b>Coverage Duration</b>	All uses (initial, reauth): 12 months
<b>Other Criteria</b>	Reauth (CD, UC, AS, PsA, RA): Documentation of positive clinical response to therapy. Reauth (plaque psoriasis): Documentation of positive clinical response to therapy as evidenced by one of the following:

Last Updated: June 2021

	reduction in the body surface area (BSA) involvement from baseline, OR improvement in symptoms (eg, pruritus, inflammation) from baseline.
--	--------------------------------------------------------------------------------------------------------------------------------------------

## INLYTA (S)

---

### Products Affected

- Inlyta

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Renal cell carcinoma (RCC): Diagnosis of RCC. One of the following: (1) disease has relapsed or (2) diagnosis of stage IV disease.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

139

## INQOVI (S)

---

### Products Affected

- Inqovi

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Myelodysplastic syndrome (MDS): Diagnosis of myelodysplastic syndrome. Patient has ONE of the following French- American-British subtypes: a) refractory anemia, b) refractory anemia with ringed sideroblasts, c) refractory anemia with excess blasts, or d) chronic myelomonocytic leukemia (CMML).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist/oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

140

## INREBIC (S)

---

### Products Affected

- Inrebic

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Myelofibrosis: Diagnosis of one of the following: primary myelofibrosis, post-polycythemia vera myelofibrosis, or post-essential thrombocythemia myelofibrosis.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist/oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

141

## INTRON A (S)

---

### Products Affected

- Intron A

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic hepatitis B: Diagnosis of chronic hepatitis B infection, and patient is without decompensated liver disease. Chronic Hepatitis C: Diagnosis of chronic hepatitis C, patient without decompensated liver disease, patient has not previously been treated with interferon, and one of the following - 1) used in combination with ribavirin or 2) contraindication or intolerance to ribavirin. Metastatic renal cell carcinoma (RCC): diagnosis of metastatic RCC, used in combination with Avastin (bevacizumab). Other: diagnosis of condylomata acuminata (genital or perianal), diagnosis of hairy cell leukemia, diagnosis of AIDS-related Kaposi's sarcoma, diagnosis of malignant melanoma, diagnosis of Stage III or IV follicular Non-Hodgkin's Lymphoma, as maintenance therapy for the treatment of multiple myeloma.
Age Restrictions	N/A
Prescriber Restrictions	RCC: Prescribed by or in consultation with an oncologist.
Coverage Duration	HepB, HepC: 48 wks. Condylomata acuminata (genital or perianal): 6 wks. Other: 12 months
Other Criteria	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

142

## IRESSA (S)

---

### Products Affected

- Iressa

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Non-small cell lung cancer (NSCLC): Diagnosis of metastatic NSCLC AND Patient has known active epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutations as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

143

# ISOTRETINOIN (S)

## Products Affected

- Accutane
- Amnesteem
- Claravis
- Isotretinoin CAPS 10MG, 20MG, 30MG, 40MG
- Myorisan
- Zenatane

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Acne (initial): Diagnosis of acne. One of the following: A) Prescribed by a dermatologist or, B) Trial and failure, contraindication or intolerance to an adequate trial (at least 6 weeks) on two of the following conventional therapy regimens: a) topical retinoid or retinoid-like agent [eg, Retin-A/Retin-A Micro (tretinoin)], b) oral antibiotic [eg, Ery-Tab (erythromycin), Minocin (minocycline)], c) topical antibiotic with or without benzoyl peroxide [eg, Cleocin-T (clindamycin), erythromycin, BenzaClin (benzoyl peroxide/clindamycin), Benzamycin (benzoyl peroxide/erythromycin)].
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Acne (initial): 5 months. Acne (reauth): Retreatment - 5 months, Dose Titration - 1 month
<b>Other Criteria</b>	Acne, Retreatment (reauth): After more than 2 months off therapy, persistent or recurring acne is still present. Acne, Dose Titration (reauth): Confirmation that the total cumulative dose is less than 150 mg/kg.

Last Updated: June 2021



## ISTODAX (S)

---

### Products Affected

- Romidepsin INJ 27.5MG/5.5ML

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cutaneous T-cell lymphoma (CTCL): Diagnosis of CTCL. Trial and failure, contraindication, or intolerance to at least one systemic therapy for the treatment of CTCL [e.g., Trexall (methotrexate), Targretin (bexarotene), cyclophosphamide] Peripheral T-cell lymphoma (PTCL): Diagnosis of PTCL. Trial and failure, contraindication, or intolerance to at least one therapy for the treatment of PTCL (e.g., conventional chemotherapy such as CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone), CHOEP (cyclophosphamide, doxorubicin, vincristine, etoposide, prednisone), Dose-adjusted EPOCH (etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin), HyperCVAD (cyclophosphamide, vincristine, doxorubicin, dexamethasone) alternating with high-dose methotrexate and cytarabine).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	CTCL, PTCL: Prescribed by or in consultation with an oncologist/hematologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

145

## ISTURISA (S)

---

### Products Affected

- Isturisa

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cushing's disease (initial): Diagnosis of Cushing's disease. One of the following: a) Patient is not a candidate for pituitary surgery, OR b) Pituitary surgery has not been curative for the patient.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Cushing's disease (initial): Prescribed by or in consultation with an endocrinologist.
<b>Coverage Duration</b>	Cushing's disease (initial, reauth): 12 months
<b>Other Criteria</b>	Cushing's disease (reauth): Documentation of positive clinical response to therapy (e.g., a clinically meaningful reduction in 24-hour urinary free cortisol levels, improvement in signs or symptoms of the disease).

Last Updated: June 2021

Formulary ID: 21381, Version 14

146

# IVIG (S)

---

## Products Affected

- Bivigam INJ 5GM/50ML
- Carimune Nanofiltered INJ 12GM, 6GM
- Flebogamma Dif
- Gammagard Liquid
- Gammagard S/d Iga Less Than 1mcg/ml
- Gammaked INJ 10GM/100ML, 1GM/10ML, 20GM/200ML, 5GM/50ML
- Gammaplex INJ 10GM/100ML, 10GM/200ML, 20GM/200ML, 20GM/400ML, 5GM/100ML, 5GM/50ML
- Gamunex-c
- Octagam INJ 10GM/100ML, 10GM/200ML, 1GM/20ML, 2.5GM/50ML, 20GM/200ML, 2GM/20ML, 30GM/300ML, 5GM/100ML, 5GM/50ML
- Panzyga
- Privigen

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	All uses (initial, reauth): Contraindications to immune globulin therapy (i.e., IgA deficiency with antibodies to IgA and a history of hypersensitivity or product specific contraindication). Priviligen only: Hyperprolinemia. Octagam only: Allergy to corn. Gammaplex only: Hereditary intolerance to fructose. Infants for whom sucrose or fructose tolerance has not been established.
<b>Required Medical Information</b>	Initial: Immune globulin (Ig) will be administered at the minimum effective dose and appropriate frequency for the prescribed diagnosis. For IVIG – Ig is being used intravenously (IV) AND One of the following diagnoses: [A] Primary Immunodeficiency 1) Common variable immunodeficiency. 2) Congenital agammaglobulinemia (X-linked or autosomal recessive). 3) Severe combined immunodeficiencies. 4) Wiskott-Aldrich syndrome. OR 5) Other PI with an immunologic evaluation including IgG levels below the normal laboratory value for the patient’s age at the time of diagnosis and the patient lacks an adequate response to protein and polysaccharide antigens (i.e., tetanus toxoid or diphtheria toxoid and pneumovax or HiB vaccine). [B] Secondary Acquired Antibody Deficiency 1) B-cell chronic lymphocytic leukemia with an Ig level less than 500 mg/dL OR history of recurrent bacterial infections. 2) HIV infection with an Ig level less than 400 mg/dL OR Patient has active bleeding or a platelet count less than $10 \times 10^9/L$ . 3) Multiple myeloma in plateau phase and patient has hypogammaglobulinemia. [C] Hematological Autoimmune Disorders 1) Acquired (pure) red cell aplasia (PRCA) that is immunologic and patient had a trial and failure, contraindication, or intolerance (TF/C/I) to a corticosteroid and an immunosuppressant (i.e., cyclophosphamide, cyclosporine) OR patient has viral PRCA caused by parvovirus B19. 2) Fetal alloimmune thrombocytopenia. 3) Hemolytic disease of the newborn and the patient has established hyperbilirubinemia. 4) Idiopathic thrombocytopenic purpura and patient had a TF/C/I to a corticosteroid OR a platelet count less than 30,000 cells/mm <sup>3</sup> . Continued in Other Criteria Section.
<b>Age Restrictions</b>	HIV (initial): patient is less than or equal to 12 years of age.
<b>Prescriber Restrictions</b>	All uses (initial, reauth): Prescribed by or in consultation with a physician who has specialized expertise in managing patients on immune globulin therapy (e.g., immunologist, hematologist, neurologist).

Last Updated: June 2021

<b>Coverage Duration</b>	4 months: Solid organ transplant. 12 months: all other diagnoses.
<b>Other Criteria</b>	<p>[D] Neuromuscular Autoimmune Disorders 1) Chronic inflammatory demyelinating polyneuropathy. 2) Guillain-Barré syndrome. 3) Inflammatory myopathies (dermatomyositis or polymyositis) AND Patient had a TF/C/I to a corticosteroid AND an immunosuppressant (i.e., azathioprine, methotrexate, cyclosporine A, cyclophosphamide, or tacrolimus). 4) Lambert-Eaton myasthenic syndrome AND Patient had a TF/C/I to a corticosteroid AND an immunosuppressant (e.g., azathioprine). 5) Multifocal motor neuropathy. 6) Myasthenia gravis with severe exacerbations or myasthenic crises AND Patient had a TF/C/I to a corticosteroid AND an immunosuppressant (i.e., azathioprine, cyclosporine, cyclophosphamide, or mycophenolate mofetil). 7) Stiff person syndrome AND Patient had a TF/C/I to at least 2 standard therapies (i.e., benzodiazepines, muscle relaxants, or anti-convulsants).</p> <p>[E] Other Disorders 1) Autoimmune blistering disease AND Patient had a TF/C/I to a corticosteroid AND an immunosuppressant (i.e., cyclophosphamide, dapson, methotrexate, azathioprine, or mycophenolate mofetil). 2) Kawasaki syndrome. 3) Solid organ transplant and IVIG is being used for CMV prophylaxis, or patient is a kidney transplant recipient and has donor specific antibodies, or patient has steroid-resistant rejection and had a TF/C/I to standard therapies. For SCIG (Gamunex-C, Gammagard Liquid, Gammaked only)- Immune globulin is being used subcutaneously AND One of the following PI diagnoses: 1) Common variable immunodeficiency. 2) Congenital agammaglobulinemia (X-linked or autosomal recessive). 3) Severe combined immunodeficiencies. 4) Wiskott-Aldrich syndrome. OR 5) Other PI with an immunologic evaluation including IgG levels below the normal laboratory value for the patient's age at the time of diagnosis and patient lacks an adequate response to protein and polysaccharide antigens (i.e., tetanus toxoid or diphtheria toxoid and pneumovax or HiB vaccine). All products: Subject to Part B vs. Part D review. For non-oncology renewal, the patient has experienced an objective improvement on immune globulin therapy and the immune globulin will be administered at the minimum effective dose (by decreasing the dose, increasing the frequency, or implementing both strategies) for maintenance therapy.</p>

## JAKAFI (S)

---

### Products Affected

- Jakafi

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Myelofibrosis: Diagnosis of primary myelofibrosis, OR post-polycythemia vera myelofibrosis, OR post-essential thrombocythemia myelofibrosis. Polycythemia vera: Diagnosis of polycythemia vera, AND trial and failure, contraindication, or intolerance to hydroxyurea. Acute graft versus host disease (aGVHD): Diagnosis of aGVHD. Disease is steroid-refractory.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Myelofibrosis, Polycythemia vera: Prescribed by or in consultation with a hematologist/oncologist. Acute graft versus host disease: Prescribed by or in consultation with one of the following: hematologist, oncologist, physician experienced in the management of transplant patients.
<b>Coverage Duration</b>	12 months.
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

## JUXTAPID (S)

---

### Products Affected

- Juxtapid

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Homozygous familial hypercholesterolemia (HoFH) (initial): Submission of medical records (eg, chart notes, laboratory values) documenting diagnosis of HoFH as confirmed by one of the following: a) genetic confirmation of 2 mutations in the LDL receptor, ApoB, PCSK9, or LDL receptor adaptor protein 1 (ie, LDLRAP1 or ARH), or b) both of the following: 1) either untreated LDL-C greater than 500 mg/dL or treated LDL-C greater than 300 mg/dL AND 2) either xanthoma before 10 years of age or evidence of heterozygous FH in both parents. Patient is receiving other lipid-lowering therapy. Trial and failure, contraindication, or intolerance to Repatha therapy. Not used in combination with a proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	HoFH (initial, reauth): Prescribed by a cardiologist, endocrinologist, or lipid specialist.
<b>Coverage Duration</b>	HoFH (initial): 6 months. (reauth): 12 months
<b>Other Criteria</b>	HoFH (reauthorization): Patient continues to receive other lipid-lowering therapy. Submission of medical records (eg, chart notes, laboratory values) documenting LDL-C reduction while on therapy. Not used in combination with a proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor.

Last Updated: June 2021

## KALBITOR (S)

---

### Products Affected

- Kalbitor

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Treatment of hereditary angioedema (HAE) attacks: Diagnosis of HAE. For the treatment of acute HAE attacks. Not used in combination with other approved treatments for acute HAE attacks.
<b>Age Restrictions</b>	12 years of age or older
<b>Prescriber Restrictions</b>	HAE: Prescribed by or in consultation with an immunologist, allergist, or rheumatologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

152



## KALYDECO (S)

---

### Products Affected

- Kalydeco

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cystic Fibrosis (CF) (Initial): Diagnosis of cystic fibrosis. Patient has at least one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to ivacaftor potentiation based on clinical and/or in vitro assay data as detected by a U.S. Food and Drug Administration (FDA)-cleared cystic fibrosis mutation test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
<b>Age Restrictions</b>	CF (Initial): 4 months of age or older
<b>Prescriber Restrictions</b>	CF (initial): Prescribed by or in consultation with a specialist affiliated with a CF care center or pulmonologist
<b>Coverage Duration</b>	CF (initial, reauth): 12 months
<b>Other Criteria</b>	CF (Reauth): Documentation of positive clinical response (i.e. improvement in lung function [percent predicted forced expiratory volume in one second (PPFEV1)], decreased number of pulmonary exacerbations) while on therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

153

## KANJINTI (S)

---

### Products Affected

- Kanjinti

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Breast cancer: Diagnosis of human epidermal growth factor receptor 2 (HER2)-overexpressing breast cancer. One of the following treatment regimens: a) As adjuvant treatment, b) metastatic disease and one of the following: 1) used in combination with a taxane (eg, docetaxel, paclitaxel), or 2) used as a single agent in a patient who has received one or more chemotherapy regimens for metastatic disease, or c) used in combination with Perjeta (pertuzumab). Gastric Cancer: Diagnosis of HER2-overexpressing gastric or gastroesophageal junction adenocarcinoma (locally advanced, recurrent, or metastatic). Used in combination with one of the following treatment regimens: a) Platinol (cisplatin) and Adrucil (5-fluorouracil), or b) Platinol (cisplatin) and Xeloda (capecitabine).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses: Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

## KANUMA (S)

---

### Products Affected

- Kanuma

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Lysosomal acid lipase deficiency: Diagnosis of lysosomal acid lipase deficiency (LAL-D). Diagnosis was confirmed by an enzymatic blood (e.g., dried blood spot test) or genetic test.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a specialist experienced in the treatment of inborn errors of metabolism, gastroenterologist, or lipidologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

155

## **KESIMPTA (S)**

---

### **Products Affected**

- Kesimpta

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Relapsing forms of multiple sclerosis (RRMS) (initial): Diagnosis of a relapsing form of multiple sclerosis (MS) (e.g., clinically isolated syndrome, relapsing-remitting disease, secondary progressive disease, including active disease with new brain lesions). One of the following: a) Failure after a trial of at least 4 weeks, contraindication, or intolerance to one of the following: Aubagio (teriflunomide), Avonex (interferon beta-1a), Bafiertam (monomethyl fumarate), Betaseron (interferon beta-1b), Copaxone/Glatopa (glatiramer acetate), Extavia (interferon beta-1b), Gilenya (fingolimod), Mavenclad (cladribine), Mayzent (siponimod), Plegridy (peginterferon beta-1a), Rebif (interferon beta-1a), Brand Tecfidera/generic dimethyl fumarate, Vumerity (diroximel fumarate), Zeposia (ozanimod), OR b) Patient is not a candidate for any of the drugs listed as prerequisites due to the severity of their multiple sclerosis, OR c) For continuation of prior therapy. Not used in combination with another disease-modifying therapy for MS. Not used in combination with another B-cell targeted therapy (e.g., rituximab [Rituxan], belimumab [Benlysta], ocrelizumab [Ocrevus]).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	RRMS (Initial, Reauth): Prescribed by or in consultation with a neurologist
<b>Coverage Duration</b>	Initial, Reauth: 12 months
<b>Other Criteria</b>	RRMS (Reauth): Documentation of positive clinical response to therapy (e.g., stability in radiologic disease activity, clinical relapse, or disease progression). Not used in combination with another disease-modifying therapy for MS. Not used in combination with another B-cell targeted therapy (e.g., rituximab [Rituxan], belimumab [Benlysta], ocrelizumab [Ocrevus]).

Last Updated: June 2021

## KEYTRUDA (S)

---

### Products Affected

- Keytruda INJ 100MG/4ML

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

158

## KISQALI (S)

---

### Products Affected

- Kisqali

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Breast cancer: Diagnosis of advanced or metastatic breast cancer. Disease is hormone receptor (HR)-positive and human epidermal growth factor receptor 2 (HER2)-negative. One of the following: A) Used in combination with an aromatase inhibitor [e.g., Femara (letrozole)] OR B) Used in combination with Faslodex (fulvestrant).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy

Last Updated: June 2021

Formulary ID: 21381, Version 14

159

## KISQALI-FEMARA PACK (S)

---

### Products Affected

- Kisqali Femara 200 Dose
- Kisqali Femara 400 Dose
- Kisqali Femara 600 Dose

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Breast cancer: Diagnosis of advanced or metastatic breast cancer. Disease is hormone receptor (HR)-positive and human epidermal growth factor receptor 2 (HER2)-negative. One of the following: A) Patient is postmenopausal OR B) Both of the following: a) Patient is pre/perimenopausal AND b) Treated with a Luteinizing Hormone-Releasing Hormone (LHRH) agonist (e.g. leuprolide).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

160



## KORLYM (S)

---

### Products Affected

- Korlym

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cushing's syndrome (Initial): Diagnosis of endogenous Cushing's syndrome (i.e., hypercortisolism is not a result of chronic administration of high dose glucocorticoids). Diagnosis of either type 2 diabetes mellitus or diagnosis of glucose intolerance. Patient has either failed surgery or patient is not a candidate for surgery. Patient is not pregnant.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Initial: Prescribed by or in consultation with an endocrinologist.
<b>Coverage Duration</b>	Initial, reauth: 6 months
<b>Other Criteria</b>	Reauth: Documentation of one of the following: patient has improved glucose tolerance while on therapy or patient has stable glucose tolerance while on therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

161

# KOSELUGO (S)

---

## Products Affected

- Koselugo

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Neurofibromatosis Type 1 (NF1): Diagnosis of NF1. Patient has plexiform neurofibromas that are both of the following: inoperable and causing significant morbidity (e.g., disfigurement, motor dysfunction, pain, airway dysfunction, visual impairment). Patient is able to swallow a capsule whole.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with one of the following: oncologist or neurologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

162

## KUVAN (S)

---

### Products Affected

- Kuvan
- Sapropterin Dihydrochloride

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Phenylketonuria (PKU) (initial): Diagnosis of PKU. Patient will have blood Phe levels measured after 1 week of therapy (new starts to therapy only) and periodically for up to 2 months of therapy to determine response.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	PKU (Init): 2 months (Reauth): 12 months
<b>Other Criteria</b>	PKU (reauth): Patient has had an objective response to therapy, defined as a 30% or greater reduction in phenylalanine (Phe) blood levels from baseline. Patient will continue to have blood Phe levels measured periodically during therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

163

## KYNMOBI (S)

---

### Products Affected

- Kynmobi
- Kynmobi Titration Kit

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	Parkinson's disease (PD) (Initial): Not used with any 5-HT <sub>3</sub> antagonist (e.g., ondansetron, granisetron, dolasetron, palonosetron, alosetron)
<b>Required Medical Information</b>	Parkinson's disease (PD) (Initial): Diagnosis of PD. Patient is experiencing acute intermittent hypomobility (defined as “off” episodes characterized by muscle stiffness, slow movements, or difficulty starting movements). Used in combination with other medications for the treatment of PD (e.g., carbidopa/levodopa, pramipexole, ropinirole, etc.).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	PD (Initial): Prescribed by or in consultation with a neurologist.
<b>Coverage Duration</b>	PD (Initial, reauth): 12 months
<b>Other Criteria</b>	PD (Reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

164

## LEMTRADA (S)

---

### Products Affected

- Lemtrada

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple Sclerosis (MS): Diagnosis of a relapsing form of MS (eg, relapsing-remitting MS, secondary-progressive MS with relapses). One of the following: 1) Patient has not been previously treated with alemtuzumab, and failure after a trial of at least 4 weeks, contraindication, or intolerance to two of the following disease-modifying therapies for MS: Aubagio (teriflunomide), Avonex (interferon beta-1a), Betaseron (interferon beta-1b), Copaxone/Glatopa (glatiramer acetate), Extavia (interferon beta-1b), Gilenya (fingolimod), Mavenclad (cladribine), Mayzent (siponimod), Ocrevus (ocrelizumab), Plegridy (peginterferon beta-1a), Rebif (interferon beta-1a), Tecfidera (dimethyl fumarate), Tysabri (natalizumab), or 2) Patient has previously received treatment with alemtuzumab, and at least 12 months have or will have elapsed since the most recent treatment course with alemtuzumab. Not used in combination with another disease-modifying therapy for MS.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	MS: 12 months.
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

165

## LENVIMA (S)

---

### **Products Affected**

- Lenvima 10 Mg Daily Dose
- Lenvima 12mg Daily Dose
- Lenvima 14 Mg Daily Dose
- Lenvima 18 Mg Daily Dose
- Lenvima 20 Mg Daily Dose
- Lenvima 24 Mg Daily Dose
- Lenvima 4 Mg Daily Dose
- Lenvima 8 Mg Daily Dose

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Differentiated thyroid cancer (DTC): Diagnosis of DTC. Disease is locally recurrent or metastatic. Patient has symptomatic or progressive disease. Disease is refractory to radioactive iodine treatment. Renal Cell Carcinoma (RCC): Diagnosis of advanced RCC. Treatment follows one prior anti-angiogenic therapy. Used in combination with everolimus. Hepatocellular Carcinoma (HCC): Diagnosis of HCC. One of the following: patient has metastatic disease, or patient has extensive liver tumor burden, or patient is inoperable by performance status or comorbidity (local disease or local disease with minimal extrahepatic disease only), or disease is unresectable. Endometrial Carcinoma (EC): Diagnosis of advanced endometrial carcinoma that is not microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR). Patient has disease progression following systemic therapy. Used in combination with Keytruda (pembrolizumab) therapy. Patient is not a candidate for curative surgery or radiation.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	DTC/RCC/EC: Prescribed by or in consultation with an oncologist. HCC: Prescribed by or in consultation with one of the following: oncologist, hepatologist, or gastroenterologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

## LETAIRIS (S)

---

### Products Affected

- Ambrisentan

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist.
<b>Coverage Duration</b>	PAH (Initial): 6 months. PAH (Reauth): 12 months
<b>Other Criteria</b>	PAH (Reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

168



## LIBTAYO (S)

---

### Products Affected

- Libtayo

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cutaneous Squamous Cell Carcinoma (CSCC): Diagnosis of CSCC. Disease is metastatic or locally advanced. Patient is not a candidate for curative surgery or curative radiation.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	CSCC: Prescribed by or in consultation with an oncologist or dermatologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

169

## LIDOCAINE TOPICAL (S)

---

### Products Affected

- 7t Lido Gel
- Glydo
- Lidocaine OINT 5%
- Lidocaine Hcl PRSY
- Lidocaine Hcl SOLN 4%
- Lidocaine Hcl Jelly
- Lidocaine/prilocaine CREA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	3 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

170

# LIDODERM (S)

---

## Products Affected

- Lidocaine PTCH 5%

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Post-herpetic neuralgia: Diagnosis of post-herpetic neuralgia.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

171

## LONSURF (S)

### Products Affected

- Lonsurf

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Colorectal Cancer: Diagnosis of metastatic colorectal cancer AND trial and failure, contraindication, or intolerance to at least one component in the following: fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy (e.g., FOLFOX, FOLFIRI, FOLFOXIRI) AND trial and failure, contraindication, or intolerance to at least one anti-VEGF therapy (e.g., Avastin) AND One of the following: A) patient has KRAS wild-type tumors and trial and failure, contraindication, or intolerance to at least one anti-EGFR therapy (e.g., Vectibix, Erbitux) OR Patient has KRAS mutant tumors. Gastric/Gastroesophageal Junction Adenocarcinoma: Diagnosis of metastatic gastric cancer or diagnosis of metastatic gastroesophageal junction adenocarcinoma. Trial and failure, contraindication or intolerance to at least two of the following: fluopyrimidine-based chemotherapy (e.g. fluorouracil), Platinum-based chemotherapy (e.g., carboplatin, cisplatin, oxaliplatin), Taxane (e.g., docetaxel, paclitaxel) or irinotecan-based chemotherapy, HER2/neu-targeted therapy (e.g., trastuzumab) (if HER2 overexpression).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

172

## LORBRENA (S)

---

### Products Affected

- Lorbrena

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Non-small cell lung cancer (NSCLC): Diagnosis of metastatic NSCLC. Patient has an anaplastic lymphoma kinase (ALK)-positive tumor.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

173

# LOTRONEX (S)

---

## Products Affected

- Alosetron Hydrochloride

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Severe Diarrhea-Predominant Irritable Bowel Syndrome (IBS) in Women (initial): All of the following: 1) diagnosis of severe diarrhea-predominant IBS, 2) symptoms for at least 6 months, 3) female patient, AND 4) trial and failure, contraindication, or intolerance to an antidiarrheal agent [eg, loperamide].
<b>Age Restrictions</b>	Initial: 18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	IBS (initial): 12 weeks. IBS (reauth): 6 mo.
<b>Other Criteria</b>	IBS (reauthorization): Symptoms of IBS continue to persist, AND documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

174

## LOVAZA (S)

---

### Products Affected

- Omega-3-acid Ethyl Esters

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Severe Hypertriglyceridemia (initial): Diagnosis of hypertriglyceridemia and patient has a pre-treatment triglyceride (TG) level greater than or equal to 500 mg/dL.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Initial/Reauth: 12 months
<b>Other Criteria</b>	Severe Hypertriglyceridemia (reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

175

## LUMIZYME (S)

---

### Products Affected

- Lumizyme

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Pompe disease: Diagnosis of Pompe disease [acid alpha-glucosidase (GAA) deficiency].
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

176



## LUPKYNIS (S)

---

### Products Affected

- Lupkynis

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Lupus Nephritis (initial): Diagnosis of active lupus nephritis. Used in combination with immunosuppressive therapy (e.g., mycophenolate mofetil, methylprednisolone).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Lupus Nephritis (initial): Prescribed by or in consultation with a nephrologist or rheumatologist
<b>Coverage Duration</b>	Lupus Nephritis (initial, reauth): 12 months
<b>Other Criteria</b>	Lupus Nephritis (reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

177

# LUPRON (S)

---

## Products Affected

- Leuprolide Acetate INJ 1MG/0.2ML

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Prostate Cancer: Diagnosis of advanced or metastatic prostate cancer.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Prostate CA: 12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

178

# LUPRON DEPOT (S)

---

## Products Affected

- Lupron Depot (1-month)
- Lupron Depot (3-month)
- Lupron Depot (4-month)
- Lupron Depot (6-month)

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Prostate Cancer (7.5 mg, 22.5 mg, 30 mg, 45 mg): Diagnosis of advanced or metastatic prostate cancer. Endometriosis (3.75 mg, 11.25 mg) (initial): Diagnosis of endometriosis. One of the following: Patient has had surgical ablation to prevent recurrence, or trial and failure, contraindication, or intolerance to one NSAID and one oral contraceptive. Uterine Leiomyomata (UL) (3.75 mg, 11.25 mg): a) For use prior to surgery to reduce size of fibroids to facilitate a surgical procedure (eg, myomectomy, hysterectomy) OR b) all of the following: treatment of anemia, anemia is caused by uterine leiomyomata (fibroids), and for use prior to surgery.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Prostate CA: 12 mo. Endomet (init, reauth):6mo. UL (anemia):3 mo (fibroids):4 mo
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

179

## LUPRON DEPOT PED (S)

---

### Products Affected

- Lupron Depot-ped (1-month)
- Lupron Depot-ped (3-month)

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Central Precocious Puberty (CPP) (initial): Diagnosis of CPP (idiopathic or neurogenic). Early onset of secondary sexual characteristics in females less than age 8 or males less than age 9. Advanced bone age of at least one year compared with chronologic age. One of the following: a) patient has undergone gonadotropin-releasing hormone agonist (GnRHa) testing AND Peak luteinizing hormone (LH) level above pre-pubertal range, or b) patient has a random LH level in the pubertal range.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	CPP (initial, reauth): Prescribed by or in consultation with a pediatric endocrinologist.
<b>Coverage Duration</b>	CPP (init, reauth): 12 months
<b>Other Criteria</b>	CPP (reauthorization): LH levels have been suppressed to pre-pubertal levels.

Last Updated: June 2021

Formulary ID: 21381, Version 14

180

# LYNPARZA TABLET (S)

---

## **Products Affected**

- Lynparza TABS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Ovarian cancer, advanced disease with known or suspected BRCA mutation with 3 or more prior lines of chemotherapy: Diagnosis of advanced ovarian cancer. Presence of deleterious or suspected deleterious germline BRCA-mutations as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Trial and failure, contraindication, or intolerance to three or more prior lines of chemotherapy (e.g., paclitaxel with cisplatin). Maintenance treatment of recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer: Diagnosis of one of the following: recurrent epithelial ovarian cancer, recurrent fallopian tube cancer, or recurrent primary peritoneal cancer. Used for maintenance treatment in patients who are in a complete or partial response to platinum-based chemotherapy (e.g., cisplatin, carboplatin). Breast cancer: Diagnosis of metastatic breast cancer. Presence of a deleterious or suspected deleterious germline BRCA-mutation as detected by an FDA-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Disease is human epidermal growth factor receptor 2 (HER2)-negative. Patient has been previously treated with chemotherapy (e.g., anthracycline, taxane) in the neoadjuvant, adjuvant, or metastatic setting. One of the following: a) Disease is hormone receptor (HR) negative, or b) Disease is hormone receptor (HR)-positive and one of the following: i) patient has been treated with prior endocrine therapy or ii) patient is considered an inappropriate candidate for endocrine therapy. See Other Criteria
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses (except prostate cancer): Prescribed by or in consultation with an oncologist. Prostate cancer: Prescribed by or in consultation with an oncologist or urologist.
<b>Coverage Duration</b>	12 months

Last Updated: June 2021

<p><b>Other Criteria</b></p>	<p>First-line maintenance treatment of BRCA-mutated advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer: Diagnosis of one of the following: advanced epithelial ovarian cancer, advanced fallopian tube cancer, or advanced primary peritoneal cancer. Presence of deleterious or suspected deleterious BRCA-mutation as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Patient has had a complete or partial response to first-line platinum-based chemotherapy (e.g., carboplatin, cisplatin). Will be used as first-line maintenance treatment. Pancreatic adenocarcinoma: Diagnosis of metastatic pancreatic adenocarcinoma. Presence of a deleterious or suspected deleterious germline BRCA-mutation as detected by an FDA-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Disease has not progressed while receiving at least 16 weeks of a first-line platinum-based chemotherapy regimen (e.g., FOLFIRINOX, FOLFOX, etc.). First-line maintenance treatment of HRD-positive advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer in combination with bevacizumab: Diagnosis of advanced epithelial ovarian cancer, advanced fallopian tube cancer, or advanced primary peritoneal cancer. Cancer is associated with homologous recombination deficiency (HRD)-positive status (defined by either: a deleterious or suspected deleterious BRCA mutation, and/or genomic instability) as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Patient has had a complete or partial response to first-line platinum-based chemotherapy (e.g., carboplatin, cisplatin). Used in combination with bevacizumab (e.g., Avastin, Mvasi). Will be used as first-line maintenance treatment. Prostate cancer: Diagnosis of metastatic castration-resistant prostate cancer. Presence of deleterious or suspected deleterious homologous recombination repair (HRR) gene mutation as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Disease has progressed following prior treatment with one of the following: a) enzalutamide (Xtandi) or b) abiraterone (e.g., Zytiga, Yonsa). All indications: Approve for continuation of prior therapy.</p>
------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

# MAKENA (S)

---

## Products Affected

- Hydroxyprogesterone Caproate INJ  
250MG/ML
- Makena INJ 275MG/1.1ML

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Preterm birth prophylaxis: Patient had a previous singleton (single offspring) spontaneous preterm birth. Patient is having a singleton pregnancy. Therapy will be started between 16 weeks, 0 days and 20 weeks, 6 days of gestation. Therapy will be continued until week 37 (through 36 weeks, 6 days) of gestation or delivery, whichever occurs first.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Preterm birth prophylaxis: Prescribed by or in consultation with a specialist in obstetrics and gynecology
<b>Coverage Duration</b>	Preterm birth prophylaxis: 21 weeks
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

184



# MARINOL (S)

---

## Products Affected

- Dronabinol

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Nausea and Vomiting Associated with Cancer Chemotherapy (CINV): Patient is receiving cancer chemotherapy. Trial and failure, contraindication, or intolerance to one 5HT-3 receptor antagonist (eg, Anzemet [dolasetron], Kytril [granisetron], or Zofran [ondansetron]). Trial and failure, contraindication, or intolerance to one of the following: Compazine (prochlorperazine), Decadron (dexamethasone), Haldol (haloperidol), Zyprexa (olanzapine). AIDS anorexia: Diagnosis of anorexia with weight loss in patients with AIDS. Patient is on antiretroviral therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	CINV: 6 months. AIDS anorexia: 3 months.
<b>Other Criteria</b>	Subject to Part B vs. Part D review. CINV: Approve for continuation of therapy for treatment covered under Part B when patient is receiving cancer chemotherapy.

Last Updated: June 2021

## MAVYRET (S)

---

### Products Affected

- Mavyret

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Criteria will be applied consistent with current AASLD/IDSA guideline. All patients: Diagnosis of chronic hepatitis C, patient is without decompensated liver disease (defined as Child-Pugh Class B or C), and not used in combination with another HCV direct acting antiviral agent [e.g., Harvoni, Zepatier].
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with one of the following: Hepatologist, Gastroenterologist, Infectious disease specialist, HIV specialist certified through the American Academy of HIV Medicine.
<b>Coverage Duration</b>	8 to 16 weeks. Criteria will be applied consistent with current AASLD/IDSA guideline.
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

186

## **MEKINIST (S)**

---

### **Products Affected**

- Mekinist

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Melanoma: Diagnosis of unresectable or metastatic melanoma AND cancer is BRAF V600E or V600K mutant type as detected by a U.S. Food and Drug Administration (FDA)-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Adjuvant Treatment for Melanoma: Diagnosis of melanoma. Cancer is BRAF V600E or V600K mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Involvement of lymph nodes following complete resection. Used as adjunctive therapy. Medication is used in combination with Tafenlar (dabrafenib). Non-small Cell Lung Cancer (NSCLC): All of the following: diagnosis of metastatic non-small cell lung cancer AND cancer is BRAF V600E mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) AND medication is used in combination with Tafenlar (dabrafenib). Anaplastic Thyroid Cancer (ATC): Diagnosis of locally advanced or metastatic ATC. Cancer is BRAF V600E mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Cancer may not be treated with standard locoregional treatment options. Medication is used in combination with Tafenlar (dabrafenib).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

## MEKTOVI (S)

---

### Products Affected

- Mektovi

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Melanoma: Diagnosis of unresectable melanoma or metastatic melanoma. Cancer is BRAF V600E or V600K mutant type (MT) as detected by a U.S. Food and Drug Administration (FDA)-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Used in combination with Braftovi (encorafenib).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy

Last Updated: June 2021

Formulary ID: 21381, Version 14

189

# METHOTREXATE INJECTION (S)

---

## Products Affected

- Rasuvo INJ 10MG/0.2ML, 12.5MG/0.25ML, 15MG/0.3ML, 17.5MG/0.35ML, 20MG/0.4ML, 22.5MG/0.45ML, 25MG/0.5ML, 30MG/0.6ML, 7.5MG/0.15ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Rheumatoid Arthritis (RA) (initial): Diagnosis of severe, active RA. Polyarticular juvenile idiopathic arthritis (PJIA) (initial): Diagnosis of active PJIA. Psoriasis (initial): Diagnosis of severe psoriasis. All Indications (initial): Trial and failure or intolerance to oral methotrexate.
Age Restrictions	N/A
Prescriber Restrictions	RA, PJIA (initial): Prescribed by or in consultation with a rheumatologist. Psoriasis (initial): Prescribed by or in consultation with a dermatologist.
Coverage Duration	All Indications (Initial, reauth): 12 months
Other Criteria	All Indications (reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

190

# MIGRANAL (S)

---

## Products Affected

- Dihydroergotamine Mesylate  
NASAL SOLN

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Initial: Diagnosis of migraine headaches with or without aura. Will be used for the acute treatment of migraine. One of the following: Trial and failure or intolerance to one triptan (e.g., eletriptan, rizatriptan, sumatriptan) or contraindication to all triptans. If patient has 4 or more headache days per month, patient must meet one of the following: a) currently being treated with Elavil (amitriptyline) or Effexor (venlafaxine) unless there is a contraindication or intolerance to these medications, OR b) currently being treated with Depakote/Depakote ER (divalproex sodium) or Topamax (topiramate) unless there is a contraindication or intolerance to these medications, OR c) currently being treated with a beta blocker (i.e., atenolol, propranolol, nadolol, timolol, or metoprolol) unless there is a contraindication or intolerance to these medications.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Initial, Reauth: Prescribed by or in consultation with a neurologist, headache specialist, or pain specialist.
<b>Coverage Duration</b>	Initial: 3 months. Reauth: 12 months.
<b>Other Criteria</b>	Reauth: Patient has experienced a positive response to therapy (e.g., reduction in pain, photophobia, phonophobia, nausea).

Last Updated: June 2021

## MOZOBIL (S)

---

### Products Affected

- Mozobil

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Hematopoietic Stem Cell (HSC) Mobilization: Patient with non-Hodgkin's lymphoma (NHL) or multiple myeloma (MM) who will be undergoing autologous HSC transplantation. Used in combination with granulocyte-colony stimulating factor (G-CSF) [e.g., Neupogen (filgrastim), Zarxio (filgrastim)].
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist/oncologist
<b>Coverage Duration</b>	One course of therapy up to 4 days
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

192



## MS INTERFERONS (NON-PREFERRED) (S)

---

### Products Affected

- Rebif
- Rebif Rebidose
- Rebif Rebidose Titration Pack
- Rebif Titration Pack

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Multiple Sclerosis (MS): Diagnosis of a relapsing form of MS (eg, clinically isolated syndrome, relapsing-remitting disease, secondary progressive disease, including active disease with new brain lesions). One of the following: 1) Trial and failure, contraindication, or intolerance (TF/C/I) to one of the following: Avonex (interferon beta-1a), Betaseron (interferon beta-1b), Extavia (interferon beta-1b), Plegridy (peginterferon beta-1a), or 2) for continuation of prior therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

193

## MS INTERFERONS (PREFERRED) (S)

---

### Products Affected

- Avonex
- Avonex Pen
- Betaseron
- Plegridy
- Plegridy Starter Pack

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple Sclerosis (MS): Diagnosis of a relapsing form of MS (eg, clinically isolated syndrome, relapsing-remitting disease, secondary progressive disease, including active disease with new brain lesions).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

194

## MVASI (S)

---

### Products Affected

- Mvasi

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	Non-Small Cell Lung Cancer: Excluded if squamous cell histology.
<b>Required Medical Information</b>	<p>Colorectal Cancer: Diagnosis of metastatic colorectal cancer. One of the following: 1) Both of the following: a) used as first- or second-line treatment and b) used in combination with an intravenous 5-fluorouracil-based chemotherapy, OR 2) All of the following: a) used as second-line treatment, b) used in combination with fluoropyrimidine-irinotecan-based chemotherapy or fluoropyrimidine-oxaliplatin-based chemotherapy, and c) patient has progressed on a first-line bevacizumab-containing regimen.</p> <p>Non-Small Cell Lung Cancer (NSCLC): Diagnosis of NSCLC. Disease is unresectable, locally advanced, recurrent, or metastatic. Used as first-line treatment. Used in combination with paclitaxel and carboplatin.</p> <p>Renal Cell Cancer: Diagnosis of metastatic renal cell cancer. Used in combination with interferon-alpha.</p> <p>Cervical Cancer: Diagnosis of carcinoma of the cervix. Disease is persistent, recurrent, or metastatic. Used in combination with one of the following: a) paclitaxel and cisplatin or b) paclitaxel and topotecan.</p> <p>Glioblastoma: Diagnosis of recurrent glioblastoma.</p> <p>Epithelial Ovarian, Fallopian Tube, or Primary Peritoneal Cancer: Diagnosis of epithelial ovarian cancer, fallopian tube cancer, or primary peritoneal cancer. One of the following: 1) All of the following: a) disease is stage 3 or 4, b) patient has been treated with bevacizumab as a single agent, c) treatment is following surgical resection, and d) used in combination with carboplatin and paclitaxel, OR 2) All of the following: a) disease is platinum-resistant recurrent, b) patient has received no more than 2 prior chemotherapy regimens, and c) used in combination with paclitaxel, pegylated liposomal doxorubicin, or topotecan, OR 3) All of the following: a) disease is platinum-sensitive recurrent, b) patient has been treated with bevacizumab as a single agent, and c) used in combination with one of the following: i) carboplatin and paclitaxel or ii) carboplatin and gemcitabine.</p>
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months

Last Updated: June 2021

<b>Other Criteria</b>	Approve for continuation of prior therapy.
-----------------------	--------------------------------------------

## MYALEPT (S)

---

### Products Affected

- Myalept

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Lipodystrophy (initial): Diagnosis of congenital or acquired generalized lipodystrophy AND one of the following: 1) Diabetes mellitus or insulin resistance despite optimized insulin therapy at maximum tolerated doses OR 2) Hypertriglyceridemia despite optimized therapy with at least two triglyceride-lowering agents from different classes (e.g., fibrates, statins).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Initial: Prescribed by or in consultation with an endocrinologist
<b>Coverage Duration</b>	Initial, reauth: 12 months
<b>Other Criteria</b>	Lipodystrophy (reauth): Patient has experienced an objective response to therapy, such as A) Sustained reduction in hemoglobin A1c (HbA1c) level from baseline OR B) Sustained reduction in triglyceride (TG) levels from baseline.

Last Updated: June 2021

Formulary ID: 21381, Version 14

198

## MYCAPSSA (S)

---

### Products Affected

- Mycapssa

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Acromegaly (initial): Diagnosis of acromegaly. One of the following: 1) Inadequate response to surgical resection and/or pituitary irradiation, or 2) Patient is not a candidate for surgical resection or pituitary irradiation. Patient has responded to and tolerated treatment with octreotide or lanreotide.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Acromegaly (initial, reauth): 12 months
<b>Other Criteria</b>	Acromegaly (reauth): Documentation of positive clinical response to therapy (e.g., reduction or normalization of IGF-1/GH level for same age and sex, reduction in tumor size)

Last Updated: June 2021

Formulary ID: 21381, Version 14

199

# NAGLAZYME (S)

---

## Products Affected

- Naglazyme

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Mucopolysaccharidosis (MPS VI): Diagnosis of MPS VI (Maroteaux-Lamy Syndrome).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	MPS VI: 12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

200



## NATPARA (S)

---

### Products Affected

- Natpara

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Hypocalcemia (Initial): Diagnosis of hypocalcemia due to chronic hypoparathyroidism. Not used in the setting of acute post-surgical hypoparathyroidism. Patient does not have a known calcium-sensing receptor mutation. Patient has a documented parathyroid hormone concentration that is inappropriately low for the level of calcium, recorded on at least two occasions within the previous 12 months. Patient has normal thyroid-stimulating hormone concentrations if not on thyroid hormone replacement therapy (or if on therapy, the dose had to have been stable for greater than or equal to 3 months). Patient has normal magnesium and serum 25-hydroxyvitamin D concentrations. Will be used as an adjunct treatment.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Hypocalcemia (initial): Prescribed by or in consultation with an endocrinologist.
<b>Coverage Duration</b>	Initial: 6 months. Reauth: 12 months
<b>Other Criteria</b>	Hypocalcemia (Reauth): One of the following: A) Patient has achieved and maintained serum calcium levels in the ideal range (7.5 - 10.6 mg/dL ), OR B) Patient has experienced a 50% or greater reduction in oral calcium intake, OR C) Patient has experienced a 50% or greater reduction in oral vitamin D intake.

Last Updated: June 2021

## NERLYNX (S)

---

### Products Affected

- Nerlynx

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Early Stage Breast cancer: Diagnosis (dx) of early stage breast cancer. Disease is human epidermal growth factor receptor 2 (HER2)-positive. Patient has received adjuvant trastuzumab-based therapy. Advanced or Metastatic Breast Cancer: Dx of advanced or metastatic breast cancer. Disease is human epidermal growth factor receptor 2 (HER2)-positive. Patient has received two or more prior anti-HER2 based regimens (e.g., trastuzumab + pertuzumab + docetaxel, ado-trastuzumab emtansine, etc.). Used in combination with capecitabine.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

202

# NEULASTA (S)

---

## **Products Affected**

- Neulasta
- Neulasta Onpro Kit

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	<p>Febrile neutropenia (FN) prophylaxis: Patient will be receiving prophylaxis for FN due to one of the following: 1) Patient is receiving National Cancer Institute’s Breast Intergroup, INT C9741 dose dense chemotherapy protocol for primary breast cancer, 2) patient is receiving a dose-dense chemotherapy regimen for which the incidence of FN is unknown, 3) patient is receiving chemotherapy regimen(s) associated with greater than 20% incidence of FN, 4) both of the following: a) patient is receiving chemotherapy regimen(s) associated with 10-20% incidence of FN, AND b) patient has one or more risk factors associated with chemotherapy-induced infection, FN, or neutropenia, OR 5) Both of the following: a) patient is receiving myelosuppressive anticancer drugs associated with neutropenia, AND b) patient has a history of FN or dose-limiting event during a previous course of chemotherapy (secondary prophylaxis). Acute radiation syndrome (ARS): Patient was/will be acutely exposed to myelosuppressive doses of radiation (hematopoietic subsyndrome of ARS). Treatment of FN: Patient has received or is receiving myelosuppressive anticancer drugs associated with neutropenia. Diagnosis of FN. Patient is at high risk for infection-associated complications.</p>
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses: Prescribed by or in consultation with a hematologist/oncologist
<b>Coverage Duration</b>	ARS: 1 mo. FN (prophylaxis, treatment): 3 mo or duration of tx.
<b>Other Criteria</b>	N/A

Last Updated: June 2021

# NEXAVAR (S)

---

## Products Affected

- Nexavar

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Renal cell carcinoma (RCC): Diagnosis of RCC. Hepatocellular carcinoma (HCC): Diagnosis of HCC. One of the following: patient has metastatic disease, or patient has extensive liver tumor burden, or patient is inoperable by performance status or comorbidity (local disease or local disease with minimal extrahepatic disease only), or disease is unresectable. Differentiated thyroid carcinoma (DTC): Diagnosis of DTC (ie, follicular carcinoma, Hurthle cell carcinoma, or papillary carcinoma). One of the following: locally recurrent disease, metastatic disease, or unresectable disease. One of the following: patient has symptomatic disease or patient has progressive disease. Disease is refractory to radioactive iodine (RAI) treatment. Medullary thyroid carcinoma (MTC): Diagnosis of MTC. One of the following: 1) Disease is progressive or 2) Disease is symptomatic with distant metastases. Trial and failure, contraindication, or intolerance to Caprelsa (vandetanib) or Cometriq (cabozantinib).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	DTC, MTC: Prescribed by or in consultation with an oncologist. RCC: Prescribed by or in consultation with an oncologist or nephrologist. HCC: Prescribed by or in consultation with an oncologist, hepatologist, or gastroenterologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

## NINLARO (S)

---

### Products Affected

- Ninlaro

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple myeloma: Diagnosis of multiple myeloma. Used in combination with Revlimid (lenalidomide) and dexamethasone. Patient has received at least one prior therapy for multiple myeloma [eg, Revlimid (lenalidomide), Thalomid (thalidomide), Velcade (bortezomib)].
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist/oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

206

## NON-PREFERRED TIRF (S)

---

### Products Affected

- Abstral SUBL 100MCG, 200MCG, 400MCG, 600MCG, 800MCG
- Fentanyl Citrate TABS
- Lazanda

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For the management of breakthrough cancer pain. Patient is currently taking a long-acting opioid around the clock for cancer pain. Patient must have at least a one week history of ONE of the following medications to demonstrate tolerance to opioids: Morphine sulfate at doses of greater than or equal to 60 mg/day, Fentanyl transdermal patch at doses greater than or equal to 25 µg/hr, Oxycodone at a dose of greater than or equal to 30 mg/day, Oral hydromorphone at a dose of greater than or equal to 8 mg/day, Oral oxymorphone at a dose of greater than or equal to 25 mg/day, or an alternative opioid at an equianalgesic dose (e.g., oral methadone greater than or equal to 20 mg/day). Trial and failure or intolerance to generic fentanyl lozenge.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with one of the following: Pain specialist, Oncologist, Hematologist, Hospice care specialist, or Palliative care specialist.
Coverage Duration	12 months
Other Criteria	N/A

Last Updated: June 2021

## NORTHERA (S)

---

### Products Affected

- Droxidopa
- Northera

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Neurogenic orthostatic hypotension (NOH) (init): Diagnosis of symptomatic NOH. NOH is caused by one of the following conditions: primary autonomic failure (eg, Parkinson's disease, multiple system atrophy, pure autonomic failure), dopamine beta-hydroxylase deficiency, non-diabetic autonomic neuropathy. Trial and failure, contraindication, or intolerance to one of the following agents: fludrocortisone acetate, midodrine.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	NOH (init): Prescribed by or in consultation with a cardiologist, neurologist, or nephrologist
<b>Coverage Duration</b>	NOH (init): 1 month (reauth): 12 months
<b>Other Criteria</b>	NOH (reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

208



# NOVANTRONE (S)

---

## Products Affected

- Mitoxantrone Hcl INJ 2MG/ML

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple Sclerosis (MS): Diagnosis (dx) of one of the following: secondary progressive MS: gradually worsening disability with or without superimposed relapses, progressive relapsing MS: progression of disability from the onset with superimposed relapses, or worsening relapsing-remitting MS: neurological status remains significantly abnormal in between MS relapses. Trial and failure, contraindication, or intolerance to one of the following disease-modifying therapies for MS: Avonex (interferon beta-1a), Aubagio (teriflunomide), Betaseron (interferon beta-1b), Copaxone/Glatopa (glatiramer acetate), Extavia (interferon beta-1b), Gilenya (fingolimod), Lemtrada (alemtuzumab), Mavenclad (cladribine), Mayzent (siponimod), Ocrevus (ocrelizumab), Plegridy (peginterferon beta-1a), Rebif (interferon beta-1a), Tecfidera (dimethyl fumarate), Tysabri (natalizumab). Left ventricular ejection fraction (LVEF) greater than or equal to 50%. Neutrophil count greater than or equal to 1500 cell/mm <sup>3</sup> . Lifetime cumulative dose less than 140 mg/m <sup>2</sup> . Prostate Cancer (PC): Dx of advanced hormone-refractory (castration-resistant) PC. Used in combination with corticosteroids (eg, prednisone, methylprednisolone). LVEF greater than or equal to 50%. Neutrophil count greater than or equal to 1500 cell/mm <sup>3</sup> . Acute Non-Lymphocytic Leukemia (ANLL): Dx of ANLL (eg, myelogenous, promyelocytic, monocytic, and erythroid). Used in combination with other medications used for the treatment of ANLL. LVEF greater than or equal to 50%.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	PC: Prescribed by or in consultation with an oncologist. ANLL: Prescribed by or in consultation with a hematologist/oncologist.
<b>Coverage Duration</b>	All Uses: 6 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

210

## NPLATE (S)

---

### Products Affected

- Nplate

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Immune thrombocytopenia (ITP): Diagnosis of one of the following: a) ITP or b) relapsed/refractory ITP. Baseline platelet count is less than 30,000/mcL. Patient's degree of thrombocytopenia and clinical condition increase the risk of bleeding. Trial and failure, contraindication, or intolerance to one of the following: corticosteroids (e.g., dexamethasone, prednisone), immune globulins (e.g., Gammaplex, Gammagard S/D), or splenectomy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	ITP (initial): Prescribed by or in consultation with a hematologist/oncologist.
<b>Coverage Duration</b>	ITP (initial, reauth): 12 months
<b>Other Criteria</b>	ITP (reauth): Documentation of positive clinical response to therapy as evidenced by an increase in platelet count to a level sufficient to avoid clinically important bleeding.

Last Updated: June 2021

Formulary ID: 21381, Version 14

211

## NUBEQA (S)

---

### Products Affected

- Nubeqa

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Non-metastatic castration-resistant or castration-recurrent prostate cancer (nmCRPC): Diagnosis of non-metastatic castration-resistant (chemical or surgical) or castration-recurrent prostate cancer.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist or urologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

212

# NUCALA (S)

---

## **Products Affected**

- Nucala

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Asthma (init): Diagnosis of severe asthma. Asthma is an eosinophilic phenotype as defined by one of the following: baseline (pre-treatment) peripheral blood eosinophil level is greater than or equal to 150 cells/microliter or peripheral blood eosinophil levels were greater than or equal to 300 cells/microliter within the past 12 months. Patient has had at least one or more asthma exacerbations requiring systemic corticosteroids within the past 12 months or Patient has had any prior intubation for an asthma exacerbation or Patient has had a prior asthma-related hospitalization within the past 12 months. Patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications: a) Both of the following: i) High-dose inhaled corticosteroid (ICS) [e.g., greater than 500 mcg fluticasone propionate equivalent/day] and ii) additional asthma controller medication [e.g., leukotriene receptor antagonist, long-acting beta-2 agonist (LABA), tiotropium], OR b) One maximally-dosed combination ICS/LABA product [e.g., Advair (fluticasone propionate/salmeterol), Dulera (mometasone/formoterol), Symbicort (budesonide/formoterol), Breo Ellipta (fluticasone/vilanterol)]. Eosinophilic Granulomatosis with Polyangiitis (EGPA) (init): Diagnosis of EGPA. Patient's disease has relapsed or is refractory to standard of care therapy (i.e., corticosteroid treatment with or without immunosuppressive therapy). Patient is currently receiving corticosteroid therapy (e.g., prednisolone, prednisone).
<b>Age Restrictions</b>	Asthma (init): Age greater than or equal to 6 years
<b>Prescriber Restrictions</b>	Asthma (init, reauth): Prescribed by or in consultation with a pulmonologist or allergist/immunologist. EGPA (init): Prescribed by or in consultation with a pulmonologist, rheumatologist or allergist/immunologist. HES (init): Prescribed by or in consultation with an allergist/immunologist or hematologist.
<b>Coverage Duration</b>	Asthma (init): 6 mo, Asthma (reauth): 12 months. EGPA, HES (init, reauth): 12 months
<b>Other Criteria</b>	Hypereosinophilic Syndrome (HES) (init): Diagnosis of HES. Patient has been diagnosed for at least 6 months. Verification that other non-

Last Updated: June 2021

hematologic secondary causes have been ruled out (e.g., drug hypersensitivity, parasitic helminth infection, HIV infection, non-hematologic malignancy). Patient is FIP1L1-PDGFR $\alpha$ -negative. Patient has uncontrolled HES defined as both of the following: a) History of 2 or more flares within the past 12 months AND b) Pre-treatment blood eosinophil count greater than or equal to 1000 cells/microliter. Trial and failure, contraindication, or intolerance to corticosteroid therapy (e.g., prednisone) or cytotoxic/immunosuppressive therapy (e.g., hydroxyurea, cyclosporine, imatinib). Asthma (reauth): Documentation of positive clinical response to therapy (eg, reduction in exacerbations, improvement in forced expiratory volume in 1 second (FEV1), decreased use of rescue medications). Patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications: a) Both of the following: i) inhaled corticosteroid (ICS) (e.g., fluticasone, budesonide) and ii) additional asthma controller medication [e.g., leukotriene receptor antagonist, long-acting beta-2 agonist (LABA), tiotropium], OR b) A combination ICS/LABA product [e.g., Advair (fluticasone propionate/salmeterol), Dulera (mometasone/formoterol), Symbicort (budesonide/formoterol), Breo Ellipta (fluticasone/vilanterol)]. EGPA (reauth): Documentation of positive clinical response to therapy (e.g., increase in remission time). HES (reauth): Documentation of positive clinical response to therapy (e.g., reduction in flares, decreased blood eosinophil count, reduction in corticosteroid dose).

# NUEDEXTA (S)

## Products Affected

- Nuedexta

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Pseudobulbar affect (PBA) (initial): Diagnosis of PBA. Patient does not have any of the following contraindications: a) Concomitant use with other drugs containing quinidine, quinine, or mefloquine, b) History of Nuedexta, quinine, mefloquine or quinidine-induced thrombocytopenia, hepatitis, bone marrow depression, or lupus-like syndrome, c) Known hypersensitivity to dextromethorphan (e.g., rash, hives), d) Taking monoamine oxidase inhibitors (MAOIs) (e.g., phenelzine, selegiline, tranylcypromine) or have taken MAOIs within the preceding 14 days, e) Has prolonged QT interval, congenital long QT syndrome or a history suggestive of torsades de pointes, or has heart failure, f) Receiving drugs that both prolong QT interval and are metabolized by CYP2D6 (e.g., thioridazine, pimozide), g) Has complete atrioventricular (AV) block without implanted pacemakers, or at high risk of complete AV block. PBA (reauth): Documentation of clinical benefit from ongoing therapy as demonstrated by a decrease in inappropriate laughing or crying episodes.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	PBA (initial): Prescribed by or in consultation with one of the following specialists: neurologist, psychiatrist.
<b>Coverage Duration</b>	PBA (initial/reauth): 12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021



## NUPLAZID (S)

---

### Products Affected

- Nuplazid

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Parkinson's disease psychosis: Diagnosis of Parkinson's disease. Patient has at least one of the following: hallucinations or delusions.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

## NURTEC (S)

### Products Affected

- Nurtec

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Initial: Diagnosis of migraine with or without aura. Will be used for the acute treatment of migraine. Will not be used for preventive treatment of migraine. Patient has fewer than 15 headache days per month. Trial and failure or intolerance to one triptan (e.g., eletriptan, rizatriptan, sumatriptan) or a contraindication to all triptans. If patient has 4 or more headache days per month, patient must meet one of the following: a) currently being treated with Elavil (amitriptyline) or Effexor (venlafaxine) unless there is a contraindication or intolerance to these medications, OR b) currently being treated with Depakote/Depakote ER (divalproex sodium) or Topamax (topiramate) unless there is a contraindication or intolerance to these medications, OR c) currently being treated with a beta blocker (i.e., atenolol, propranolol, nadolol, timolol, or metoprolol) unless there is a contraindication or intolerance to these medications. Medication will not be used in combination with another oral CGRP inhibitor.
<b>Age Restrictions</b>	Initial: 18 years of age or older.
<b>Prescriber Restrictions</b>	Initial, Reauth: Prescribed by or in consultation with a neurologist, headache specialist, or pain specialist.
<b>Coverage Duration</b>	Initial: 3 months. Reauth: 12 months.
<b>Other Criteria</b>	Reauth: Patient has experienced a positive response to therapy (e.g., reduction in pain, photophobia, phonophobia, nausea). Will not be used for preventive treatment of migraine. Medication will not be used in combination with another oral CGRP inhibitor.

Last Updated: June 2021

## **NYVEPRIA (S)**

---

### **Products Affected**

- Nyvepria

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	<p>Febrile neutropenia (FN) prophylaxis: Patient will be receiving prophylaxis for FN due to one of the following: 1) Patient is receiving National Cancer Institute’s Breast Intergroup, INT C9741 dose dense chemotherapy protocol for primary breast cancer, 2) patient is receiving a dose-dense chemotherapy regimen for which the incidence of FN is unknown, 3) patient is receiving chemotherapy regimen(s) associated with greater than 20% incidence of FN, 4) both of the following: a) patient is receiving chemotherapy regimen(s) associated with 10-20% incidence of FN, AND b) patient has one or more risk factors associated with chemotherapy-induced infection, FN, or neutropenia, OR 5) both of the following: a) patient is receiving myelosuppressive anticancer drugs associated with neutropenia, AND b) patient has a history of FN or dose-limiting event during a previous course of chemotherapy (secondary prophylaxis). Treatment of FN (off-label): Patient has received or is receiving myelosuppressive anticancer drugs associated with neutropenia. Diagnosis of FN. Patient is at high risk for infection-associated complications. Acute radiation syndrome (ARS) (off-label): Patient was/will be acutely exposed to myelosuppressive doses of radiation (hematopoietic subsyndrome of ARS).</p>
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses: Prescribed by or in consultation with a hematologist/oncologist
<b>Coverage Duration</b>	ARS: 1 mo. FN (prophylaxis, treatment): 3 mo or duration of tx.
<b>Other Criteria</b>	All Indications: Trial and failure or intolerance to both of the following: Neulasta/Neulasta Onpro AND Udenyca.

Last Updated: June 2021

## ODOMZO (S)

---

### Products Affected

- Odomzo

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Basal cell carcinoma: Diagnosis of locally advanced basal cell carcinoma AND One of the following: 1) Cancer has recurred following surgery or radiation therapy or 2) Patient is not a candidate for surgery or radiation therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist or dermatologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

221

## OFEV (S)

---

### Products Affected

- Ofev

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	<p>Idiopathic pulmonary fibrosis (IPF) (initial): Diagnosis of IPF as documented by all of the following: a) exclusion of other known causes of interstitial lung disease (ILD) (eg, domestic and occupational environmental exposures, connective tissue disease, drug toxicity) AND b) one of the following: i) in patients not subjected to surgical lung biopsy, the presence of a usual interstitial pneumonia (UIP) pattern on high-resolution computed tomography (HRCT) revealing IPF or probable IPF, OR ii) in patients subjected to a lung biopsy, both HRCT and surgical lung biopsy pattern revealing IPF or probable IPF. Systemic sclerosis-associated interstitial lung disease (SSc-ILD) (initial): Diagnosis of SSc-ILD as documented by all of the following: a) exclusion of other known causes of ILD (eg, domestic and occupational environmental exposures, connective tissue disease, drug toxicity) AND b) One of the following: i) In patients not subjected to surgical lung biopsy, the presence of idiopathic interstitial pneumonia (eg, fibrotic nonspecific interstitial pneumonia [NSIP], usual interstitial pneumonia [UIP] and centrilobular fibrosis) pattern on HRCT revealing SSc-ILD or probable SSc-ILD, OR ii) in patients subjected to a lung biopsy, both HRCT and surgical lung biopsy pattern revealing SSc-ILD or probable SSc-ILD. Chronic Fibrosing Interstitial Lung Diseases (ILDs) with a Progressive Phenotype (initial): 1) diagnosis of chronic fibrosing interstitial lung disease, AND 2) patient has a high-resolution computed tomography (HRCT) showing at least 10% of lung volume with fibrotic features, AND 3) disease has a progressive phenotype as observed by one of the following: decline of forced vital capacity (FVC), worsening of respiratory symptoms, or increased extent of fibrosis seen on imaging.</p>
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	IPF, SSc-ILD, Chronic Fibrosing ILDs with a Progressive Phenotype (initial): Prescribed by or in consultation with a pulmonologist
<b>Coverage Duration</b>	Initial, reauth: 12 months

Last Updated: June 2021

<b>Other Criteria</b>	IPF, SSc-ILD, Chronic Fibrosing ILDs with a Progressive Phenotype (reauth): Documentation of positive clinical response to therapy.
-----------------------	-------------------------------------------------------------------------------------------------------------------------------------



## ONUREG (S)

---

### Products Affected

- Onureg

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Acute Myeloid Leukemia (AML): Diagnosis of acute myeloid leukemia (AML). Patient has received previous treatment with an intensive induction chemotherapy regimen (e.g., cytarabine + daunorubicin, cytarabine + idarubicin, etc.). Patient has achieved one of the following: a) first complete remission (CR) or b) complete remission with incomplete blood count recovery (CRi). Patient is not able to complete intensive curative therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist/oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

225

## OPDIVO (S)

---

### Products Affected

- Opdivo

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

226

## OPSUMIT (S)

---

### Products Affected

- Opsumit

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist.
<b>Coverage Duration</b>	PAH: Initial: 6 months. Reauth: 12 months.
<b>Other Criteria</b>	PAH (Reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

227

## ORENCIA IV (S)

---

### Products Affected

- Orencia INJ 250MG

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. One of the following: a) Trial and failure, contraindication, or intolerance (TF/C/I) to two of the following: Enbrel (etanercept), Humira (adalimumab), Rinvoq (upadacitinib), or Xeljanz/Xeljanz XR (tofacitinib), or b) attestation demonstrating a trial may be inappropriate, OR c) For continuation of prior Orencia therapy. Juvenile Idiopathic Arthritis (JIA) (Initial): Diagnosis of moderately to severely active polyarticular JIA. Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. JIA, PsA (Initial): One of the following: Trial and failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior Orencia therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	RA, JIA (initial): Prescribed by or in consultation with a rheumatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist.
<b>Coverage Duration</b>	All uses (initial, reauth): 12 months
<b>Other Criteria</b>	All indications (Reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

228

# ORENCIA SC (S)

---

## Products Affected

- Orencia INJ 125MG/ML, 50MG/0.4ML, 87.5MG/0.7ML
- Orencia Clickject

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. One of the following: a) Either a trial and failure, contraindication, or intolerance (TF/C/I) to two of the following: Enbrel (etanercept), Humira (adalimumab), Rinvoq (upadacitinib), Xeljanz/Xeljanz XR (tofacitinib), or attestation demonstrating a trial may be inappropriate, OR b) For continuation of prior therapy. Juvenile Idiopathic Arthritis (JIA) (Initial): Diagnosis of moderately to severely active polyarticular JIA. One of the following: a) Either a trial and failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR b) for continuation of prior therapy. Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. One of the following: a) Either a trial and failure, contraindication, or intolerance to Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR b) for continuation of prior therapy.
Age Restrictions	N/A
Prescriber Restrictions	RA, JIA (initial): Prescribed by or in consultation with a rheumatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist.
Coverage Duration	All uses (initial, reauth): 12 months
Other Criteria	All indications (Reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Last Updated: June 2021

Formulary ID: 21381, Version 14

# ORENITRAM (S)

---

## Products Affected

- Orenitram

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist.
<b>Coverage Duration</b>	PAH: Initial: 6 months. Reauth: 12 months.
<b>Other Criteria</b>	PAH (Reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

231

## ORGOVYX (S)

---

### Products Affected

- Orgovyx

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Prostate Cancer: Diagnosis of advanced prostate cancer. Disease is one of the following: 1) Evidence of biochemical or clinical relapse following local primary intervention with curative intent or 2) Newly diagnosed androgen-sensitive metastatic disease or 3) Advanced localized disease unlikely to be cured by local primary intervention with curative intent.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an urologist or oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

232



## ORIAHNN (S)

---

### Products Affected

- Oriahnn

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Initial: Diagnosis of heavy menstrual bleeding associated with uterine leiomyomas (fibroids). Patient is premenopausal. One of the following: 1) History of inadequate control of bleeding following a trial of at least 3 months, or history of intolerance or contraindication to one of the following: combination (estrogen/progesterone) oral contraceptive, progestins, or tranexamic acid or 2) Patient has had a previous interventional therapy to reduce bleeding. Treatment duration of therapy has not exceeded a total of 24 months.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Reauth: Patient has improvement in bleeding associated with uterine leiomyomas (fibroids) (e.g., significant/sustained reduction in menstrual blood loss per cycle, improved quality of life, etc.). Treatment duration of therapy has not exceeded a total of 24 months.

Last Updated: June 2021

Formulary ID: 21381, Version 14

233

# ORILISSA (S)

---

## Products Affected

- Orilissa

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Endometriosis (EM) (initial - 150 mg): Diagnosis of moderate to severe pain associated with EM. One of the following: 1) History of inadequate pain control response following a trial of at least 3 months, or history of intolerance or contraindication to one of the following: danazol, combination (estrogen/progesterone) oral contraceptive, or progestins, or 2) Patient has had surgical ablation to prevent recurrence. EM (200 mg): Diagnosis of moderate to severe pain associated with EM. One of the following: 1) History of inadequate pain control response following a trial of at least 3 months, or history of intolerance or contraindication to one of the following: danazol, combination (estrogen/progesterone) oral contraceptive, or progestins, or 2) Patient has had surgical ablation to prevent recurrence.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	EM (init, reauth-150mg): 6 mo. EM (200mg): 6 mo.
<b>Other Criteria</b>	EM (reauthorization - 150 mg): Patient has improvement in pain associated with endometriosis (e.g., improvement in dysmenorrhea and nonmenstrual pelvic pain). Treatment duration has not exceeded a total of 24 months.

Last Updated: June 2021

## ORKAMBI (S)

---

### Products Affected

- Orkambi TABS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cystic Fibrosis (CF) (Initial): Diagnosis of CF. Patient is homozygous for the F508del mutation in the CF transmembrane conductance regulator (CFTR) gene as detected by a U.S. Food and Drug Administration (FDA)-cleared cystic fibrosis mutation test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
<b>Age Restrictions</b>	CF (Initial): Patient is 6 years of age or older
<b>Prescriber Restrictions</b>	CF (initial): Prescribed by or in consultation with a specialist affiliated with a CF care center or pulmonologist
<b>Coverage Duration</b>	CF (initial, reauth): 12 months
<b>Other Criteria</b>	CF (Reauth): Patient is benefiting from treatment (i.e. improvement in lung function [forced expiratory volume in one second (FEV1)], decreased number of pulmonary exacerbations).

Last Updated: June 2021

Formulary ID: 21381, Version 14

235

# ORKAMBI GRANULES (S)

---

## Products Affected

- Orkambi PACK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cystic Fibrosis (CF) (Initial): Diagnosis of CF. Patient is homozygous for the F508del mutation in the CF transmembrane conductance regulator (CFTR) gene as detected by a U.S. Food and Drug Administration (FDA)-cleared cystic fibrosis mutation test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). One of the following: A) Patient is 2 through 5 years of age, OR B) Both of the following: Patient is 6 years of age or greater AND Patient is unable to swallow oral tablets.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	CF (Initial): Prescribed by or in consultation with a specialist affiliated with a CF care center or pulmonologist
<b>Coverage Duration</b>	CF (initial, reauth): 12 months
<b>Other Criteria</b>	CF (Reauth): Patient is benefiting from treatment (i.e., improvement in lung function [forced expiratory volume in one second (FEV1)], decreased number of pulmonary exacerbations). One of the following: A) Patient is 2 through 5 years of age, OR B) Both of the following: Patient is 6 years of age or greater AND Patient is unable to swallow oral tablets.

Last Updated: June 2021

## ORLADEYO (S)

---

### Products Affected

- Orladeyo

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Prophylaxis of hereditary angioedema (HAE) attacks: Diagnosis of HAE. For prophylaxis against HAE attacks.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	HAE (prophylaxis): Prescribed by or in consultation with an immunologist or allergist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

237

# OSMOLEX ER (S)

---

## Products Affected

- Osmolex Er TB24

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Parkinson's Disease (PD) (initial): Diagnosis of Parkinson's disease. Trial and failure or intolerance to both of the following: A) amantadine immediate release AND B) one of the following: carbidopa-levodopa, MAO-B inhibitor (e.g., rasagiline, selegiline), or dopamine agonist (e.g., pramipexole, ropinirole). Drug-Induced Extrapyrarnidal Reactions (EPS) (initial): Patient is experiencing drug-induced extrapyramidal reactions. One of the following: A) Patient has persistent extrapyramidal symptoms despite a trial of dose reduction, tapering, or discontinuation of the offending medication OR B) Patient is not a candidate for a trial of dose reduction, tapering, or discontinuation of the offending medication. Trial and failure or intolerance to amantadine immediate release.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	PD (initial): Prescribed by or in consultation with a neurologist. EPS (initial): Prescribed by or in consultation with a neurologist or psychiatrist.
<b>Coverage Duration</b>	PD, EPS (initial, reauth): 12 months
<b>Other Criteria</b>	PD, EPS (Reauth): Documentation of positive clinical response to therapy

Last Updated: June 2021

Formulary ID: 21381, Version 14

238

## OSPHENA (S)

---

### Products Affected

- Ospheña

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Dyspareunia (initial): Diagnosis of moderate to severe dyspareunia due to vulvar and vaginal atrophy associated with menopause. Vaginal dryness (initial): Diagnosis of moderate to severe vaginal dryness due to vulvar and vaginal atrophy associated with menopause.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	All uses (Initial, reauth): 12 months
<b>Other Criteria</b>	Dyspareunia, Vaginal dryness (reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

239

## OXANDRIN (S)

---

### Products Affected

- Oxandrolone TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Promote weight gain (initial): Used as adjunctive therapy to promote weight gain AND Diagnosis of one of the following: Extensive surgery, Chronic infections, Severe trauma, Failure to gain or maintain at least 90% of ideal body weight without definite pathophysiologic reasons AND a nutritional consult was performed. Counterbalance protein catabolism (initial): Used to counterbalance protein catabolism associated with chronic corticosteroid administration. Bone pain: Diagnosis of bone pain associated with osteoporosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	bone pain: 1 month. Others (initial, reauth): 3 months
Other Criteria	All diagnoses except bone pain (reauth): Documentation of a positive clinical response to therapy as evidenced by an improvement in weight gain or increase in lean body mass.

Last Updated: June 2021



## OXERVATE (S)

---

### Products Affected

- Oxervate

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Neurotrophic keratitis (NK): Diagnosis of NK.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an ophthalmologist or optometrist.
<b>Coverage Duration</b>	8 weeks.
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

241

# PALFORZIA (S)

---

## **Products Affected**

- Palforzia Initial Dose Escalation
- Palforzia Level 1
- Palforzia Level 10
- Palforzia Level 11 (maintenance)
- Palforzia Level 11 (titration)
- Palforzia Level 2
- Palforzia Level 3
- Palforzia Level 4
- Palforzia Level 5
- Palforzia Level 6
- Palforzia Level 7
- Palforzia Level 8
- Palforzia Level 9

Last Updated: June 2021

Formulary ID: 21381, Version 14

242

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	Initial: Excluded if any of the following: 1) history of eosinophilic esophagitis (EoE) or eosinophilic gastrointestinal disease, OR 2) history of severe or life-threatening episode(s) of anaphylaxis or anaphylactic shock within the past 2 months, OR 3) severe or poorly controlled asthma
<b>Required Medical Information</b>	Initial: Diagnosis and clinical history of peanut allergy as documented by both of the following: a) a serum peanut-specific IgE level of greater than or equal to 0.35 kUA/L, AND b) a mean wheal diameter that is at least 3mm larger than the negative control on skin-prick testing for peanut. One of the following: 1) patient is 4 to 17 years of age and is in the initial dose escalation phase of therapy, OR 2) patient is 4 years of age and older and is in the up-dosing or maintenance phase of therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Initial, Reauth: Prescribed by or in consultation with an allergist/immunologist.
<b>Coverage Duration</b>	Initial, Reauth: 12 months.
<b>Other Criteria</b>	N/A

Last Updated: June 2021

## PEGASYS (S)

---

### Products Affected

- Pegasys
- Pegasys Proclick INJ  
180MCG/0.5ML

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chronic hepatitis B: Diagnosis of chronic hepatitis B infection, and patient is without decompensated liver disease. Chronic Hepatitis C: Criteria will be applied consistent with current AASLD-IDSA guidance.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	HepB: 48 wks. HepC: Initial: 28 wks. Reauth: 20 wks.
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

244

## PEMAZYRE (S)

---

### Products Affected

- Pemazyre

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of cholangiocarcinoma. Disease is one of the following: unresectable locally advanced or metastatic. Disease has presence of a fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangement. Patient has been previously treated.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hepatologist, gastroenterologist, or oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

245

## PENNSAID (S)

---

### Products Affected

- Diclofenac Sodium EXTERNAL SOLN 1.5%

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Osteoarthritis of the knees (initial): Diagnosis of osteoarthritis of the knees. Patient meets one of the following: 1) Treatment failure with at least two prescription strength non-steroidal anti-inflammatory drugs (NSAIDs) OR 2) History of peptic ulcer disease/gastrointestinal bleed OR 3) Patient is older than 65 years of age with one additional risk factor for gastrointestinal adverse events (e.g. use of anticoagulants, chronic corticosteroids).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Initial, reauth: 12 months
<b>Other Criteria</b>	Osteoarthritis of the knees (reauth): Documentation of positive clinical response to therapy (e.g., improvement in pain symptoms of osteoarthritis).

Last Updated: June 2021

Formulary ID: 21381, Version 14

246

## PIQRAY (S)

---

### Products Affected

- Piqray 200mg Daily Dose
- Piqray 250mg Daily Dose
- Piqray 300mg Daily Dose

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Breast Cancer (BC): Diagnosis of advanced or metastatic BC. Disease is hormone receptor (HR)-positive, and human epidermal growth factor receptor 2 (HER2)-negative. Cancer is PIK3CA-mutated as detected by an FDA-approved test (therascreen PIK3CA RGQ PCR Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Patient is a postmenopausal woman or male. Used in combination with fulvestrant. Disease has progressed on or after an endocrine-based regimen.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

247

## POLIVY (S)

---

### Products Affected

- Polivy

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diffuse large B-cell lymphoma (DLBCL): Diagnosis of diffuse large B-cell lymphoma (DLBCL). Disease is relapsed or refractory. Used in combination with bendamustine and a rituximab product. Patient has received at least two prior therapies for DLBCL (e.g., RCHOP [rituximab, cyclophosphamide, doxorubicin, vincristine, prednisone], HSCT [hematopoietic stem cell transplantation], CAR T [chimeric antigen receptor T-cell] therapy, RCEPP [rituximab, cyclophosphamide, etoposide, prednisone, procarbazine], GemOx [gemcitabine, oxaliplatin] with or without rituximab).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist/oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

248



## POMALYST (S)

---

### Products Affected

- Pomalyst

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple Myeloma (MM): Diagnosis of MM. Used in combination with dexamethasone. Patient has received two prior therapies, including Revlimid (lenalidomide) and a proteasome inhibitor [eg, Velcade (bortezomib) or Kyprolis (carfilzomib)] or has a contraindication or intolerance to Revlimid and proteasome inhibitors. Patient has experienced disease progression on or within 60 days of completion of last therapy. Kaposi sarcoma (KS): One of the following: 1) Both of the following: a) Diagnosis of AIDS-related KS and b) Patient has failed highly active antiretroviral therapy (HAART), OR 2) Both of the following: a) Diagnosis of KS and b) Patient is HIV-negative.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All indications: Prescribed by or in consultation with an oncologist or hematologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

## PROCYSBI (S)

---

### Products Affected

- Procysbi CPDR

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Nephropathic cystinosis: Diagnosis of nephropathic cystinosis, confirmed by elevated leukocyte cystine levels (LCL) or genetic analysis of the CTNS gene or demonstration of cysteine corneal crystals by slit lamp examination AND Trial and failure or intolerance to therapy with Cystagon (immediate-release cysteamine bitartrate).
<b>Age Restrictions</b>	1 year of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

250

## **PROMACTA (S)**

---

### **Products Affected**

- Promacta

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chronic immune (idiopathic) thrombocytopenic purpura (ITP) (initial): Diagnosis of one of the following: relapsed/refractory ITP or chronic ITP. Baseline platelet count is less than 30,000/mcL. Patient's degree of thrombocytopenia and clinical condition increase the risk of bleeding. Trial and failure, intolerance, contraindication to corticosteroids or immunoglobulins or splenectomy. Chronic hepatitis C (initial): Diagnosis of chronic hepatitis C-associated thrombocytopenia. One of the following: 1) Planning to initiate and maintain interferon-based treatment, or 2) currently receiving interferon-based treatment. First-line for severe aplastic anemia (SAA): Diagnosis of SAA. Used for first-line treatment (i.e., patient has not received prior immunosuppressive therapy). Used in combination with standard immunosuppressive therapy. Patient meets at least two of the following: 1) absolute neutrophil count less than 500/mcL, 2) platelet count less than 20,000/mcL, 3) absolute reticulocyte count less than 60,000/mcL. Refractory SAA (initial): Diagnosis of refractory severe aplastic anemia. Patient has a platelet count less than 30,000/mcL. Insufficient response to immunosuppressive therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Chronic ITP and SAA: Prescribed by or in consultation with a hematologist/oncologist. Chronic hepatitis C associated thrombocytopenia: Prescribed by or in consultation with a hematologist/oncologist, gastroenterologist, hepatologist, infectious disease specialist, or HIV specialist certified through the American Academy of HIV Medicine.
<b>Coverage Duration</b>	ITP(init, reauth): 12mo. HepC: 3mo(init), 12mo(reauth). 1stline SAA: 6mo. RefractSAA: 16wk-init, 12mo-reauth
<b>Other Criteria</b>	ITP (reauth): Documentation of positive clinical response to therapy as evidenced by an increase in platelet count to a level sufficient to avoid clinically important bleeding. Hepatitis C (reauth): One of the following: 1) For patients that started treatment with eltrombopag prior to initiation of treatment with interferon, eltrombopag will be approved when both of the following are met: a) Currently on antiviral interferon therapy for

Last Updated: June 2021

---

<p>treatment of chronic hepatitis C and b) Documentation that the patient reached a threshold platelet count that allows initiation of antiviral interferon therapy with eltrombopag treatment by week 9, OR 2) For patients that started treatment with Promacta while on concomitant treatment with interferon, eltrombopag will be approved based on the following: Currently on antiviral interferon therapy for treatment of chronic hepatitis C. Refractory SAA (reauth): Documentation of positive clinical response to therapy as evidenced by an increase in platelet count.</p>
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

---

## **PROVIGIL (S)**

---

### **Products Affected**

- Modafinil

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	<p>Obstructive sleep apnea (OSA) (Initial): Diagnosis (dx) of OSA defined by one of the following: 15 or more obstructive respiratory events per hour of sleep confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible), or both of the following: 5 or more obstructive respiratory events per hour of sleep confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible), and 1 of the following symptoms: unintentional sleep episodes during wakefulness, daytime sleepiness, unrefreshing sleep, fatigue, insomnia, waking up breath holding/gasping/choking, loud snoring, or breathing interruptions during sleep. Shift-work disorder (SWD) (Initial):Dx of SWD confirmed by one of the following: 1) Symptoms of excessive sleepiness or insomnia for at least 3 months, which is associated with a work period (usually night work) that occurs during the normal sleep period, OR 2) A sleep study demonstrating loss of a normal sleep-wake pattern (ie, disturbed chronobiologic rhythmicity). Confirmation that no other medical conditions or medications are causing the symptoms of excessive sleepiness or insomnia. Narcolepsy (initial): Dx of narcolepsy as confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible). MS Fatigue (initial): Dx of multiple sclerosis (MS). Patient is experiencing fatigue. Depression (initial): Treatment-resistant depression defined as diagnosis of major depressive disorder (MDD) or bipolar depression, AND trial and failure, contraindication, or intolerance to at least two antidepressants from different classes (eg, SSRIs, SNRIs, bupropion). Used as adjunctive therapy. Idiopathic Hypersomnia (Initial): Diagnosis of idiopathic hypersomnia as confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible).</p>
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Narcolepsy: Init, Reauth: 12 mo. All other indications: Init, Reauth: 6 mo.

Last Updated: June 2021

<b>Other Criteria</b>	OSA, Narcolepsy, Idiopathic Hypersomnia (Reauth): Documentation of positive clinical response to modafinil therapy. SWD (Reauth): Documentation of positive clinical response to modafinil therapy. MS Fatigue (reauth): Patient is experiencing relief of fatigue with modafinil therapy. Depression (reauth): Documentation of positive clinical response to modafinil therapy. Used as adjunctive therapy.
-----------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------



## PULMOZYME (S)

---

### Products Affected

- Pulmozyme

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cystic Fibrosis (CF) (Initial, Reauth): Diagnosis of CF.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	CF (initial, reauth): 12 months
<b>Other Criteria</b>	Part B vs D determination applies. CF (reauth): Patient is benefiting from treatment (i.e. improvement in lung function [forced expiratory volume in one second (FEV1)], decreased number of pulmonary exacerbations).

Last Updated: June 2021

Formulary ID: 21381, Version 14

257

## QINLOCK (S)

---

### Products Affected

- Qinlock

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Gastrointestinal Stromal Tumor (GIST): Diagnosis of gastrointestinal stromal tumor (GIST). Disease is advanced. Patient has received prior treatment with three or more kinase inhibitors (e.g., sunitinib, regorafenib), one of which must include imatinib.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

258

## QUALAQUIN (S)

---

### Products Affected

- Quinine Sulfate CAPS 324MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	Excluded if used solely for the treatment or prevention of nocturnal leg cramps.
<b>Required Medical Information</b>	Malaria: Diagnosis of uncomplicated malaria. One of the following: 1) Treatment in areas of chloroquine-sensitive malaria, and trial and failure, contraindication, or intolerance to chloroquine or hydroxychloroquine, OR 2) Treatment in areas of chloroquine-resistant malaria.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	7 days
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

259

## RADICAVA (S)

---

### Products Affected

- Radicava

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Amyotrophic lateral sclerosis (ALS) (initial): Submission of medical records (e.g., chart notes, previous medical history, diagnostic testing including: imaging, nerve conduction studies, laboratory values) to support a diagnosis of “definite” or “probable” ALS per the revised El Escorial diagnostic criteria. Patient has scores of greater than or equal to 2 in all items of the ALS Functional Rating Scale-Revised (ALSFRRS-R) criteria at the start of treatment. Patient has a percent forced vital capacity (%FVC) of greater than or equal to 80% at the start of treatment.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	ALS (initial): Prescribed by or in consultation with a neurologist.
<b>Coverage Duration</b>	Initial, reauth: 6 months
<b>Other Criteria</b>	ALS (reauthorization): Documentation of a benefit from therapy (e.g., slowing in the decline of functional abilities), and Patient is not dependent on invasive ventilation or tracheostomy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

260

## RAVICTI (S)

---

### Products Affected

- Ravicti

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Urea cycle disorders (UCDs) (Initial): Diagnosis of UCDs.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	UCDs (Initial, reauth): 12 months
<b>Other Criteria</b>	UCDs (reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

261

## REBLOZYL (S)

---

### Products Affected

- Reblozyl

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Beta Thalassemia (initial): One of the following: a) Diagnosis of beta thalassemia major AND patient requires regular red blood cell (RBC) transfusions, OR b) Diagnosis of transfusion-dependent beta thalassemia.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Beta Thalassemia (initial): Prescribed by or in consultation with a hematologist.
<b>Coverage Duration</b>	Beta Thalassemia (initial, reauth): 12 months
<b>Other Criteria</b>	Beta Thalassemia (reauth): Documentation of a positive clinical response to therapy (eg., reduction in RBC transfusion burden).

Last Updated: June 2021

Formulary ID: 21381, Version 14

262

## **RENFLEXIS (S)**

---

### **Products Affected**

- Renflexis

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Crohn's Disease (CD) and Fistulizing Crohn's Disease (FCD) (initial): Dx of moderately to severely active CD or FCD. Ulcerative colitis (UC) (initial): Dx of moderately to severely active UC. Rheumatoid arthritis (RA) (initial): Diagnosis (Dx) of moderately to severely active RA. Ankylosing spondylitis (AS) (initial): Dx of active AS. Psoriatic arthritis (PsA) (initial): Dx of active PsA. Plaque psoriasis (initial): Dx of chronic severe (ie, extensive and/or disabling) plaque psoriasis. CD, FCD (initial): Trial and failure, contraindication or intolerance (TF/C/I) to one of the following conventional therapies: 6-mercaptopurine (Purinethol), azathioprine (Imuran), corticosteroids (eg, prednisone, methylprednisolone), methotrexate (Rheumatrex, Trexall). UC (initial): TF/C/I to one of the following conventional therapies: corticosteroids, aminosalicylate [eg, mesalamine (Asacol/Pentasa/Rowasa), olsalazine (Dipentum), sulfasalazine (Azulfidine/Sulfazine)], azathioprine (Imuran), 6-mercaptopurine (Purinethol). RA (initial): Receiving concurrent therapy with methotrexate (Rheumatrex/Trexall), or TF/C/I to methotrexate. AS (initial): TF/C/I to two or more NSAIDs (e.g., diclofenac, ibuprofen, meloxicam, naproxen). All indications (initial): One of the following: 1) Trial and failure or intolerance to Remicade or 2) For continuation of prior Renflexis therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Initial: RA, AS: Prescribed by or in consultation with a rheumatologist. PsA: Prescribed or in consultation with a rheumatologist or dermatologist. Crohn's Disease, Fistulizing Crohn's Disease, UC: Prescribed by or in consultation with a gastroenterologist. Plaque Psoriasis: Prescribed by or in consultation with a dermatologist.
<b>Coverage Duration</b>	All indications (initial, reauth): 12 months
<b>Other Criteria</b>	Reauthorization for CD, UC, AS, PsA, RA: Documentation of positive clinical response to therapy. Reauthorization (plaque psoriasis): Documentation of positive clinical response to therapy as evidenced by one of the following: reduction in the body surface area (BSA)

Last Updated: June 2021



	involvement from baseline, OR improvement in symptoms (eg, pruritus, inflammation) from baseline.
--	---------------------------------------------------------------------------------------------------

# REPATHA (S)

---

## **Products Affected**

- Repatha
- Repatha Pushtronex System
- Repatha Sureclick

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	<p>HeFH/ASCVD/Primary HLD (init): One of the following dx: A)HeFH as confirmed by one of the following: 1)Both of the following: a)Untreated/pre-treatment LDL greater than 190 mg/dL in an adult, AND b)One of the following: i) Family hx of tendinous xanthomas and/or arcus cornealis in 1st degree relative, or 2nd degree relative, ii)Hx of myocardial infarction (MI) in 1st-degree relative less than 60 years of age, iii)Family hx of MI in 2nd-degree relative less than 50 years of age, iv)Family hx of LDL-C greater than 190 mg/dL in 1st- or 2nd-degree relative, v)Family hx of FH in 1st- or 2nd-degree relative, or 2)Untreated/pre-treatment LDL-C greater than 190 mg/dL in an adult AND submission of MR (eg chart notes, lab values) documenting one of the following: presence of tendinous xanthoma in pt, arcus cornealis before age 45, or functional mutation in the LDL receptor, ApoB, or PCSK9 gene. OR B)ASCVD as confirmed by ACS, hx of MI, stable or unstable angina, coronary or other arterial revascularization, stroke,TIA, or peripheral arterial disease presumed to be of atherosclerotic origin. OR C)Primary hyperlipidemia (HLD). HoFH (init): Sub of MR (eg, chart notes, lab values) documenting dx of HoFH as confirmed by one of the following: 1)Gen confirmation of 2 mutations in LDL receptor, ApoB, PCSK9, or LDLRAP1 or ARH, or 2)either untreated LDL greater than 500 or treated LDL greater than 300, AND either xanthoma before 10 yo or evidence of HeFH in both parents. HeFH/ASCVD/Primary HLD (init): One of the following: set A)Both of the following: a)One of the following LDL values while on max tolerated lipid lowering regimen w/in the last 120 days: 1)LDL greater than or equal to 100 mg/dL w/ASCVD, or 2)LDL greater than or equal to 130 mg/dL w/o ASCVD. AND b)One of the following: 1)Pt has been receiving at least 12 wks of one high-intensity (HI) statin therapy (tx) and will continue to receive a HI statin [ie, atorvastatin 40-80 mg, rosuvastatin 20-40 mg] at max tolerated dose.</p>
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Initial/Reauth: Prescribed by or in consultation with a cardiologist, endocrinologist, or lipid specialist

Last Updated: June 2021

Formulary ID: 21381, Version 14

267

<b>Coverage Duration</b>	Initial: 6 months. Reauth: 12 months
<b>Other Criteria</b>	<p>Set A (cont, initial): OR (2) Both of the following: A) Pt unable to tolerate HI statin as evidenced by intolerable and persistent (ie, more than 2 wks) myalgia (muscle symptoms w/o CK elevations) or myositis (muscle symptoms w/ CK elevations less than 10 times ULN) AND B) One of the following: a) Pt has been receiving at least 12 wks of one moderate-intensity (MI) or low-intensity (LI) statin tx and will continue to receive a MI or LI statin [ie, atorvastatin 10-20 mg, rosuvastatin 5-10 mg, simvastatin 10-40 mg, pravastatin 10-80 mg, lovastatin 20-40 mg, Lescol XL (fluvastatin XL) 80 mg, fluvastatin 20-40 mg, or Livalo (pitavastatin) 1-4 mg] at max tolerated dose, OR b) Pt is unable to tolerate MI or LI statin as evidenced by intolerable and persistent (ie, more than 2 weeks) myalgia (muscle symptoms w/o CK elevations) or myositis (muscle symptoms w/ CK elevations less than 10 times ULN), OR (3) Submission of MR documenting pt has a labeled contraindication to all statins, OR (4) Pt has experienced rhabdomyolysis or muscle symptoms w/ statin treatment w/ CK elevations greater than 10 times ULN on one statin tx.</p> <p>OR set B) Both of the following: a) One of the following LDL values while on max tolerated statin tx w/in the last 120 days: (1) LDL b/t 70 and 99 mg/dL w/ ASCVD. (2) LDL b/t 100 and 129 mg/dL w/o ASCVD. AND b) Both of the following: (1) One of the following: i) Pt has been receiving at least 12 wks of one max-tolerated statin tx and will continue to receive a statin at max tolerated dose, ii) pt is unable to tolerate statin tx as evidenced by intolerable and persistent (ie, more than 2 wks) myalgia (muscle symptoms w/o CK elevations) or myositis (muscle symptoms w/ CK elevations less than 10 times ULN, iii) Submission of MR documenting patient has a labeled contraindication to all statins, or iv) Pt has experienced rhabdomyolysis or muscle symptoms w/ statin tx w/ CK elevations greater than 10 times ULN on one statin tx and (2) Pt has been receiving at least 12 weeks of ezetimibe (Zetia) tx as adjunct to max tolerated statin tx OR Pt has a hx of contraindication or intolerance to ezetimibe. HoFH (init): Pt is receiving other lipid-lowering tx (eg statin, ezetimibe). HeFH/ASCVD/Primary HLD (reauth): Pt continues to receive other lipid-lowering tx (eg statins, ezetimibe) at max tolerated dose (unless pt has documented inability to take these medications). HoFH (reauth): Pt continues to receive other lipid-lowering tx (eg statin, ezetimibe). HeFH/ASCVD/Primary HLD/HoFH (reauth): Sub of MR (eg, lab values) documenting LDL reduction while on Repatha tx. HeFH/ASCVD/Primary HLD/HoFH (Init, reauth): Prescriber attests that the info provided is true and accurate to the best of their knowledge and they understand that OptumRx may perform a routine audit and request the medical info necessary to verify the accuracy of the info provided. HoFH (Init, reauth): Not used in combo w/ Juxtapid.</p>

Last Updated: June 2021

Last Updated: June 2021

Formulary ID: 21381, Version 14

# RETACRIT (S)

---

## Products Affected

- Retacrit

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	<p>Anemia with Chronic Kidney Disease (CKD) (Initial): Diagnosis (Dx) of CKD. Anemia by lab values (Hct less than 30% or Hgb less than 10 g/dL) collected within 30 days of request. One of the following: a) both of the following: Patient is on dialysis, patient is without ESRD OR b) all of the following: patient is not on dialysis, the rate of hemoglobin decline indicates the likelihood of requiring a red blood cell (RBC) transfusion, and reducing the risk of alloimmunization and/or other RBC transfusion-related risks is a goal. Anemia with chemo (Initial): Other causes of anemia have been ruled out. Anemia by lab values (Hct less than 30%, Hgb less than 10 g/dL) collected within the prior 2 weeks of request. Cancer is a non-myeloid malignancy. Patient is receiving chemo.</p> <p>Preoperative for reduction of allogeneic blood transfusion: Patient is scheduled to undergo elective, non-cardiac, non-vascular surgery. Hgb is greater than 10 to less than or equal to 13 g/dL. Patient is at high risk for perioperative transfusions. Patient is unwilling or unable to donate autologous blood pre-operatively. Anemia in hepatitis C virus (HCV)-infected pts due to ribavirin in combination with interferon/peg-interferon (Initial): Dx of HCV infection. Anemia by labs (Hct less than 36% or Hgb less than 12 g/dL) collected within 30 days of request. Patient is receiving ribavirin and one of the following: interferon alfa or peginterferon alfa.</p> <p>Anemia with HIV (Initial): Anemia by lab values (Hgb less than 12 g/dL or Hct less than 36%) collected within 30 days of request. Serum erythropoietin level less than or equal to 500 mU/mL. Receiving zidovudine therapy or dx of HIV. Anemia in Myelodysplastic Syndrome (MDS) (Initial): Dx of MDS. Serum erythropoietin level is 500 mU/mL or less, or dx of transfusion-dependent MDS.</p>
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	CKD,HIV(Init):6mo. CKD,HIV(reauth):12mo. Chemo,HCV(all):3mo. MDS:(init) 3mo,(reauth)12mo. Preop:1mo.

Last Updated: June 2021

<p><b>Other Criteria</b></p>	<p>Subject to ESRD review. CKD (Reauth): Dx of CKD. One of the following: 1) Most recent or average (avg) Hct over 3 months is 33% or less (Hgb is 11 g/dL or less) for patients on dialysis, without ESRD, 2) Most recent or avg Hct over 3 mo is 30% or less (Hgb 10 g/dL or less) for patients not on dialysis, OR 3) Most recent or avg Hct over 3 mo is 36% or less (Hgb 12 g/dL or less) for pediatric patients. Decrease in the need for blood transfusion or Hgb increased by 1 g/dL or more from pre-treatment level. HIV (Reauth): Most recent or avg Hct over 3 months is below 36% or most recent or avg Hgb over 3 months is below 12 g/dl. Decrease in the need for blood transfusion or Hgb increased by 1 g/dL or more from pre-treatment level. Chemo (Reauth): Anemia by lab values (Hgb less than 10 g/dl or Hct less than 30%) collected within the prior 2 weeks of request. Decrease in the need for blood transfusion or Hgb increased by 1 g/dL or more from pre-treatment level. Patient is receiving chemo. HCV (Reauth): Most recent or avg Hct over 3 months is 36% or less, OR most recent or avg Hgb over 3 months is 12 g/dl or less. Decrease in the need for blood transfusion or Hgb increased by 1 g/dL or more from pre-treatment level. If patient has demonstrated response to therapy, authorization will be issued for the full course of ribavirin therapy. MDS (Reauth): Most recent or avg Hct over 3 months is 36% or less, OR most recent or avg Hgb over 3 months is 12 g/dl or less. Decrease in the need for blood transfusion or Hgb increased by 1 g/dL or more from pre-treatment level. Other Off-label uses (except MDS, HCV): Will not be approved if patient has Hgb greater than 10 g/dL or Hct greater than 30%. CKD (init, reauth), HIV (init), Chemo (init), Preop, MDS (init), HCV (init): Verify iron evaluation for adequate iron stores.</p>
------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------



## RETEVMO (S)

---

### Products Affected

- Retevmo

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Lung Cancer: Diagnosis of metastatic non-small cell lung cancer (NSCLC). Disease has presence of RET gene fusion-positive tumor(s). Medullary Thyroid Cancer (MTC): Diagnosis of medullary thyroid cancer (MTC). Disease is advanced or metastatic. Disease has presence of RET gene mutation tumor(s). Disease requires treatment with systemic therapy. Thyroid Cancer: Diagnosis of thyroid cancer. Disease is advanced or metastatic. Disease has presence of RET gene fusion-positive tumor(s). Disease requires treatment with systemic therapy. Patient is radioactive iodine-refractory or radioactive iodine therapy is not appropriate.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Lung Cancer, MTC: Prescribed by or in consultation with an oncologist. Thyroid Cancer: Prescribed by or in consultation with an endocrinologist or an oncologist.
<b>Coverage Duration</b>	Lung Cancer, MTC, Thyroid Cancer: 12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

273

## REVATIO (S)

---

### Products Affected

- Sildenafil Citrate TABS 20MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist.
<b>Coverage Duration</b>	PAH: Initial: 6 months. Reauth: 12 months.
<b>Other Criteria</b>	PAH (Reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

274

## REVCOVI (S)

---

### Products Affected

- Revcovi

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of adenosine deaminase deficiency (ADA) with severe combined immunodeficiency (SCID).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

275

## REVLIMID (S)

---

### Products Affected

- Revlimid

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple myeloma (MM): Diagnosis of MM. Either used as 1) combination therapy with dexamethasone, or 2) maintenance therapy following autologous hematopoietic stem cell transplantation (auto-HSCT). Myelodysplastic syndromes (MDS): Diagnosis of transfusion-dependent anemia due to low- or intermediate-1-risk MDS associated with a deletion 5q. Mantle cell lymphoma (MCL): Diagnosis of MCL. Disease has relapsed after, is refractory to, or progressed after at least one prior therapy (eg, bortezomib, bendamustine, cladribine, rituximab). Follicular Lymphoma (FL): Diagnosis of FL that has been previously treated. Used in combination with a rituximab product. Marginal Zone Lymphoma (MZL): Diagnosis of MZL that has been previously treated. Used in combination with a rituximab product.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist/hematologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

# RILUTEK (S)

---

## Products Affected

- Riluzole

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Amyotrophic lateral sclerosis (ALS): Diagnosis of amyotrophic lateral sclerosis (ALS).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	ALS: 12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

277

## RINVOQ (S)

---

### Products Affected

- Rinvoq

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Rheumatoid arthritis (RA) (initial): Diagnosis of moderately to severely active RA. One of the following: Trial and failure, contraindication, or intolerance (TF/C/I) to one disease modifying antirheumatic drug (DMARD) [eg, methotrexate (Rheumatrex/Trexall), Arava (leflunomide), Azulfidine (sulfasalazine)], OR for continuation of prior therapy. Not used in combination with a potent immunosuppressant (e.g., azathioprine, cyclosporine).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	RA (initial): Prescribed by or in consultation with a rheumatologist.
<b>Coverage Duration</b>	RA (initial, reauth): 12 months.
<b>Other Criteria</b>	RA (reauth): Documentation of positive clinical response to therapy. Not used in combination with a potent immunosuppressant (e.g., azathioprine, cyclosporine).

Last Updated: June 2021

Formulary ID: 21381, Version 14

278

# RITUXAN (S)

---

## Products Affected

- Rituxan

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	<p>Non-Hodgkin's Lymphoma (NHL): As first-line treatment of diffuse large B-cell, CD20-positive, NHL in combination with CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone) or other anthracycline-based chemotherapy regimens, or as first-line treatment of follicular, CD20-positive, B-cell NHL in combination with chemotherapy, or for the treatment of follicular, CD20-positive, B-cell NHL in patients achieving a complete or partial response to a rituximab product in combination with chemotherapy, used as monotherapy for maintenance therapy, or for the treatment of low-grade, CD20-positive, B-cell NHL in patients with stable disease following first-line treatment with CVP (cyclophosphamide, vincristine, prednisolone/prednisone) chemotherapy or who achieve a partial or complete response following first-line treatment with CVP (cyclophosphamide, vincristine, prednisolone/prednisone) chemotherapy, or diagnosis of relapsed or refractory, low grade or follicular CD20-positive, B-cell non-Hodgkin's lymphoma. Rheumatoid Arthritis (RA): Diagnosis of moderately to severely active RA. Concurrently on or contraindication, or intolerance to methotrexate. Trial and failure, contraindication, or intolerance (TF/C/I) to a TNF antagonist (eg, adalimumab, etanercept, infliximab). Not received in combination with a biologic DMARD [e.g., Enbrel (etanercept), Orencia (abatacept), Kineret (anakinra)]. Wegener's Granulomatosis (WG) and Microscopic Polyangiitis (MPA): Diagnosis of WG or MPA. Patient is concurrently on glucocorticoids (eg, prednisone) OR contraindication or intolerance to glucocorticoids (eg, prednisone). Immune or idiopathic thrombocytopenic purpura (ITP): Diagnosis of ITP. TF/C/I to one of the following: glucocorticoids (e.g., prednisone, methylprednisolone), immune globulins (e.g., IVIG), or splenectomy. Documented platelet count of less than <math>50 \times 10^9 /L</math>.</p>
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	ITP, CLL, NHL: Prescribed by or in consultation with a hematologist or oncologist. RA: Prescribed by or in consultation with a rheumatologist. WG, MPA: Prescribed by or in consultation with a nephrologist, pulmonologist, or rheumatologist. PV: Prescribed by or in consultation with a dermatologist

Last Updated: June 2021



<b>Coverage Duration</b>	All uses except RA, WG, MPA: 12 mos. RA: 3 months. WG, MPA: 3 months only.
<b>Other Criteria</b>	Chronic Lymphocytic Leukemia (CLL): Diagnosis of CLL. Used in combination with fludarabine and cyclophosphamide. Pemphigus Vulgaris (PV): Diagnosis of moderate to severe PV. All uses: Approve for continuation of prior therapy.

# RITUXAN HYCELA (S)

## Products Affected

- Rituxan Hycela

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Follicular Lymphoma: Diagnosis of follicular lymphoma. One of the following: 1) Disease is relapsed or refractory OR 2) Patient exhibited complete or partial response to prior treatment with rituximab in combination with chemotherapy OR 3) Disease is non-progressing or stable following prior treatment with first-line cyclophosphamide, vincristine, and prednisone (CVP) chemotherapy OR 4) Both of the following: a) Disease is previously untreated AND b) Medication is used in combination with first-line chemotherapy. Diffuse Large B-Cell Lymphoma: 1) Diagnosis of diffuse large B-cell lymphoma AND 2) Disease is previously untreated AND 3) Medication is being used in combination with cyclophosphamide, doxorubicin, vincristine, prednisone (CHOP) or other anthracycline-based chemotherapy. Chronic Lymphocytic Leukemia: 1) Diagnosis of chronic lymphocytic leukemia AND 2) Medication is being used in combination with fludarabine and cyclophosphamide (FC) therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses: Prescribed by or in consultation with a hematologist or oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

# ROZLYTREK (S)

---

## Products Affected

- Rozlytrek

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Non-small cell lung cancer (NSCLC): Diagnosis of metastatic non-small cell lung cancer (NSCLC). Patient has ROS1 rearrangement positive tumor(s). Solid Tumors: Patient has solid tumors with a neurotrophic tyrosine receptor kinase (NTRK) gene fusion (e.g., ETV6-NTRK3, TPM3-NTRK1, TPR-NTRK1, etc.). Disease is without a known acquired resistance mutation (e.g., TRKA G595R, TRKA G667C or TRKC G623R substitutions). Disease is one of the following: metastatic or unresectable (including cases where surgical resection is likely to result in severe morbidity). One of the following: disease has progressed following previous treatment (e.g., surgery, radiation therapy, or systemic therapy) or disease has no satisfactory alternative treatments.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

## RUBRACA (S)

### Products Affected

- Rubraca

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Ovarian cancer: Diagnosis of epithelial ovarian cancer, fallopian tube cancer, or primary peritoneal cancer. One of the following: 1) Both of the following: a) Presence of deleterious BRCA mutation as detected by a U.S. Food and Drug Administration (FDA)-approved diagnostic test (e.g., FoundationFocus CDxBRCA Assay) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) and b) Trial and failure, contraindication, or intolerance to two or more chemotherapies (e.g., cisplatin, carboplatin), OR 2) Both of the following: a) Disease is recurrent and b) Used for maintenance treatment in patients who are in a complete or partial response to platinum-based chemotherapy (e.g., cisplatin, carboplatin). Prostate cancer: Diagnosis of metastatic castration-resistant prostate cancer. Presence of deleterious BRCA mutation as detected by an FDA-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Patient has received previous treatment with both of the following: 1) Androgen receptor-directed therapy [e.g., Erleada (apalutamide), Xtandi (enzalutamide), Zytiga (abiraterone)], AND 2) A taxane-based chemotherapy [e.g., docetaxel, Jevtana (cabazitaxel)].
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Ovarian cancer: Prescribed by or in consultation with an oncologist. Prostate cancer: Prescribed by or in consultation with an oncologist or urologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy

Last Updated: June 2021

Last Updated: June 2021

Formulary ID: 21381, Version 14

## RUCONEST (S)

---

### Products Affected

- Ruconest

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Treatment of hereditary angioedema (HAE) attacks: Diagnosis of HAE. For the treatment of acute HAE attacks. Not used in combination with other approved treatments for acute HAE attacks.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	HAE: Prescribed by or in consultation with an immunologist, allergist, or rheumatologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

286

# RUXIENCE (S)

---

## Products Affected

- Ruxience

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	<p>Non-Hodgkin's Lymphoma (NHL): One of the following: 1) Diagnosis of follicular, CD20-positive, B-cell non-Hodgkin's lymphoma. Used as first-line treatment in combination with chemotherapy, 2) Diagnosis of follicular, CD20-positive, B-cell non-Hodgkin's lymphoma. Patient achieved a complete or partial response to a rituximab product in combination with chemotherapy. Used as monotherapy for maintenance therapy, 3) Diagnosis of low-grade, CD20-positive, B-cell non-Hodgkin's lymphoma. One of the following: a) Patient has stable disease following first-line treatment with CVP (cyclophosphamide, vincristine, prednisolone/ prednisone) chemotherapy or, b) Patient achieved a partial or complete response following first-line treatment with CVP (cyclophosphamide, vincristine, prednisolone/ prednisone) chemotherapy, 4) Diagnosis of relapsed or refractory, low grade or follicular CD20-positive, B-cell non-Hodgkin's lymphoma OR, 5) Diagnosis of diffuse large B-cell, CD20-positive, non-Hodgkin's lymphoma. Used as first-line treatment in combination with CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone) or other anthracycline-based chemotherapy regimens. Chronic Lymphocytic Leukemia (CLL): Diagnosis of chronic lymphocytic leukemia. Used in combination with fludarabine and cyclophosphamide. Wegener's Granulomatosis (WG) and Microscopic Polyangiitis (MPA): Diagnosis of WG or MPA. Patient is concurrently on glucocorticoids (eg, prednisone) OR contraindication or intolerance to glucocorticoids (eg, prednisone).</p>
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	NHL, CLL: Prescribed by or in consultation with a hematologist/oncologist. WG, MPA: Prescribed by or in consultation with a nephrologist, pulmonologist, or rheumatologist.
<b>Coverage Duration</b>	NHL, CLL: 12 months. WG, MPA: 3 months.
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021



## RUZURGI (S)

---

### Products Affected

- Ruzurgi

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Lambert-Eaton Myasthenic Syndrome (LEMS) (initial): Diagnosis of LEMS.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	LEMS (initial): Prescribed by or in consultation with a neurologist.
<b>Coverage Duration</b>	LEMS (initial): 3 months. LEMS (reauth): 12 months.
<b>Other Criteria</b>	LEMS (reauth): Documentation of positive clinical response to therapy (e.g., improvement in dynamometry, Timed 25-Foot Walk Test, Timed Up and Go Test).

Last Updated: June 2021

Formulary ID: 21381, Version 14

289

## RYDAPT (S)

---

### Products Affected

- Rydapt

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acute Myeloid Leukemia (AML): Newly diagnosed acute myeloid leukemia (AML), FMS-like tyrosine kinase 3 (FLT3) mutation-positive as detected by a U.S. Food and Drug Administration (FDA)-approved test (e.g., LeukoStrat CDx FLT3 Mutation Assay) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA), used in combination with standard cytarabine and daunorubicin induction and cytarabine consolidation. Aggressive Systemic Mastocytosis (ASM), Systemic Mastocytosis with Associated Hematological Neoplasm (SM-AHN), Mast Cell Leukemia (MCL): Diagnosis of one of the following: aggressive systemic mastocytosis (ASM), systemic mastocytosis with associated hematological neoplasm (SM-AHN), or mast cell leukemia (MCL).
Age Restrictions	N/A
Prescriber Restrictions	All uses: Prescribed by or in consultation with a hematologist or oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

290

## SABRIL (S)

---

### Products Affected

- Vigabatrin
- Vigadrone

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Complex Partial Seizures (CPS): For use as adjunctive therapy. Failure, contraindication, or intolerance to two formulary anticonvulsants [eg, Lamictal (lamotrigine), Depakene (valproic acid), Dilantin (phenytoin)]. Infantile Spasms (IS): Diagnosis of infantile spasms.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

291

# SANDOSTATIN (S)

## Products Affected

- Octreotide Acetate INJ  
1000MCG/ML, 100MCG/ML,  
200MCG/ML, 500MCG/ML,  
50MCG/ML

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Acromegaly (initial): Diagnosis of acromegaly. One of the following: A) Inadequate response to surgical resection and/or pituitary irradiation OR B) Patient is not a candidate for surgical resection or pituitary irradiation. Trial and failure, contraindication or intolerance to a dopamine agonist (e.g., bromocriptine or cabergoline) at maximally tolerated doses. Carcinoid tumor (initial): Diagnosis of metastatic carcinoid tumor requiring symptomatic treatment of severe diarrhea or flushing episodes. Vasoactive intestinal peptide tumor (initial): Diagnosis of vasoactive intestinal peptide tumor requiring treatment of profuse watery diarrhea.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	All uses (initial, reauth): 12 months
<b>Other Criteria</b>	Acromegaly (reauth): Documentation of positive clinical response to therapy (e.g., reduction or normalization of IGF-1/GH level for same age and sex, reduction in tumor size). Carcinoid tumor (reauth): Patient has improvement in number of diarrhea or flushing episodes. Vasoactive intestinal peptide tumor (reauth): Patient has improvement in number of diarrhea episodes.

Last Updated: June 2021

## SCIG (S)

---

### **Products Affected**

- Cuvitru
- Hizentra
- Xembify

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	All uses (initial, reauth): Contraindications to immune globulin therapy (i.e., IgA deficiency with antibodies to IgA and a history of hypersensitivity or product specific contraindication).
<b>Required Medical Information</b>	Initial: Immune globulin will be administered at the minimum effective dose and appropriate frequency for the prescribed diagnosis. Medication is being used subcutaneously. Diagnosis of chronic inflammatory demyelinating polyneuropathy (CIDP) OR one of the following FDA-approved or literature supported diagnoses: 1) Common variable immunodeficiency (CVID), OR 2) Congenital agammaglobulinemia (X-linked or autosomal recessive), OR 3) Severe combined immunodeficiencies (SCID), OR 4) Wiskott-Aldrich syndrome, OR 5) Other primary immunodeficiency with an immunologic evaluation including IgG levels below the normal laboratory value for the patient's age at the time of diagnosis and the patient lacks an adequate response to protein and polysaccharide antigens (i.e., tetanus toxoid or diphtheria toxoid and pneumovax or HiB vaccine).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses (initial, reauth): Prescribed by or in consultation with a physician who has specialized expertise in managing patients on SCIG therapy (e.g., immunologist, hematologist, neurologist).
<b>Coverage Duration</b>	Initial, reauth: 12 months
<b>Other Criteria</b>	Subject to Part B vs. Part D review. All uses (reauth): Patient has experienced an objective improvement on immune globulin therapy and the immune globulin will be administered at the minimum effective dose (by decreasing the dose, increasing the frequency, or implementing both strategies) for maintenance therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

294

## SECUADO (S)

---

### Products Affected

- Secuado

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of schizophrenia. Both of the following: 1) Trial and failure of Saphris (asenapine) and 2) Trial and failure, contraindication, or intolerance to one of the following generic formulary atypical antipsychotic agents: aripiprazole, olanzapine, paliperidone, quetiapine, risperidone, or ziprasidone.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

295

## SIGNIFOR (S)

---

### Products Affected

- Signifor

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cushing's disease (initial): Diagnosis of Cushing's disease. One of the following: a) Pituitary surgery has not been curative for the patient or b) Patient is not a candidate for pituitary surgery.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Cushing's disease (initial): Prescribed by or in consultation with an endocrinologist.
<b>Coverage Duration</b>	Cushing's disease (initial, reauth): 12 months
<b>Other Criteria</b>	Cushing's disease (reauth): Documentation of positive clinical response to therapy (e.g., a clinically meaningful reduction in 24-hour urinary free cortisol levels, improvement in signs or symptoms of the disease).

Last Updated: June 2021

Formulary ID: 21381, Version 14

296



## SIKLOS (S)

---

### Products Affected

- Siklos

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Sickle Cell Anemia: Diagnosis of sickle cell anemia. Patient has moderate to severe painful crises. One of the following: 1) Patient is less than 18 years of age or 2) Trial and failure, or intolerance to Droxia.
<b>Age Restrictions</b>	Patient is 2 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

297

## SKYRIZI (S)

---

### Products Affected

- Skyrizi
- Skyrizi Pen

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Plaque psoriasis (Initial): Diagnosis of moderate to severe plaque psoriasis.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Plaque psoriasis (initial): Prescribed by or in consultation with a dermatologist.
<b>Coverage Duration</b>	Plaque psoriasis (Initial, reauth): 12 months
<b>Other Criteria</b>	Plaque psoriasis (Reauth): Documentation of positive clinical response to therapy as evidenced by one of the following: reduction in the body surface area (BSA) involvement from baseline, OR improvement in symptoms (eg, pruritus, inflammation) from baseline.

Last Updated: June 2021

Formulary ID: 21381, Version 14

298

# SOLIRIS (S)

## Products Affected

- Soliris

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Paroxysmal Nocturnal Hemoglobinuria (PNH), Atypical Hemolytic Uremic Syndrome (aHUS) (initial): Diagnosis of PNH or aHUS. Trial and failure, contraindication, or intolerance to Ultomiris (ravulizumab). Generalized Myasthenia Gravis (gMG) (initial): Diagnosis of gMG. Patient is anti-acetylcholine (AChR) antibody positive. One of the following: 1) Trial and failure, contraindication, or intolerance (TF/C/I) to two immunosuppressive therapies (e.g., glucocorticoids, azathioprine, cyclosporine, mycophenolate mofetil, methotrexate, tacrolimus), or 2) TF/C/I to one immunosuppressive therapy (e.g., glucocorticoids, azathioprine, cyclosporine, mycophenolate mofetil, methotrexate, tacrolimus), and TF/C/I to chronic plasmapheresis/plasma exchange (PE) or intravenous immunoglobulin (IVIg). Neuromyelitis Optica Spectrum Disorder (NMOSD) (initial): Diagnosis of NMOSD. Patient is anti-aquaporin-4 (AQP4) antibody positive.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	gMG, NMOSD (initial): Prescribed by or in consultation with a neurologist.
<b>Coverage Duration</b>	All uses (initial, reauth): 12 months
<b>Other Criteria</b>	PNH (reauth): Documentation of positive clinical response (e.g., hemoglobin stabilization, decrease in the number of red blood cell transfusions) to therapy. aHUS (reauth): Documentation of positive clinical response (e.g., increase in mean platelet counts, hematologic normalization) to therapy. gMG, NMOSD (reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021



## SOMATULINE DEPOT (S)

---

### Products Affected

- Somatuline Depot

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Acromegaly: Diagnosis of acromegaly. One of the following: A) Inadequate response to one of the following: surgery or radiotherapy, OR B) Not a candidate for one of the following: surgery or radiotherapy. Gastroenteropancreatic neuroendocrine tumors (GEP-NETs) (120mg/0.5mL strength only): Diagnosis of GEP-NETs. Disease is one of the following: (a) unresectable, locally advanced or (b) metastatic. Carcinoid syndrome (120mg/0.5mL strength only): Diagnosis of carcinoid syndrome. Used to reduce the frequency of short-acting somatostatin analog rescue therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	All uses: 12 months
<b>Other Criteria</b>	All Indications: Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

301

## SOMAVERT (S)

---

### Products Affected

- Somavert

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Acromegaly (initial): Diagnosis of acromegaly AND Failure to surgery and/or radiation therapy and/or other medical therapies (such as dopamine agonists [e.g., bromocriptine, cabergoline]) unless patient is not a candidate for these treatment options AND trial and failure or intolerance to generic octreotide (a somatostatin analogue)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an endocrinologist
<b>Coverage Duration</b>	Initial and reauth: 12 months
<b>Other Criteria</b>	Acromegaly (reauth): Patient has experienced a positive clinical response to therapy (biochemical control, decrease or normalization of IGF-1 levels).

Last Updated: June 2021

Formulary ID: 21381, Version 14

302

# SPORANOX (S)

---

## Products Affected

- Itraconazole CAPS
- Itraconazole SOLN

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	One of the following: 1) patient has a systemic fungal infection (e.g., aspergillosis, histoplasmosis, blastomycosis), OR 2) all of the following: a) patient has a diagnosis of onychomycosis confirmed by one of the following (CAPSULE ONLY): i) positive potassium hydroxide (KOH) preparation, OR ii) fungal culture, OR iii) nail biopsy, AND b) patient has had a trial and failure, contraindication, or intolerance to oral terbinafine, OR 3) patient has a diagnosis of candidiasis (esophageal or oropharyngeal) that is refractory to treatment with fluconazole (ORAL SOLUTION ONLY).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Systemic fungal infxn:6mo.Candidiasis:1mo.Fingernail onycho:5wks.Toenail onycho, other:3mo.
<b>Other Criteria</b>	N/A

Last Updated: June 2021

## SPRAVATO (S)

---

### Products Affected

- Spravato 56mg Dose
- Spravato 84mg Dose

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Treatment-resistant depression (TRD): Diagnosis of major depressive disorder (treatment-resistant). Patient has not experienced a clinical meaningful improvement after treatment with at least two antidepressants from different classes for an adequate duration (at least 4 weeks each) in the current depressive episode. Used in combination with an oral antidepressant (e.g., duloxetine, escitalopram, sertraline).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	TRD: Prescribed by or in consultation with a psychiatrist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

304



## SPRYCEL (S)

---

### Products Affected

- Sprycel

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Philadelphia chromosome positive (Ph+)/BCR ABL chronic myelogenous leukemia (CML): Diagnosis of Ph+/BCR ABL CML. Ph+/BCR ABL acute lymphoblastic leukemia (ALL): Diagnosis of Ph+/BCR ABL ALL.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All Uses: Prescribed by or in consultation with an oncologist or hematologist
<b>Coverage Duration</b>	All Uses: 12 months
<b>Other Criteria</b>	All Uses: Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

305

## STELARA (IV) (S)

### Products Affected

- Stelara INJ 130MG/26ML

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Crohn's Disease (CD): Diagnosis of moderately to severely active CD. One of the following: a) trial and failure, contraindication, or intolerance to Humira (adalimumab), or (b) trial and failure, contraindication, or intolerance to treatment with at least one immunomodulator or corticosteroid [e.g., Purinethol (6-mercaptopurine), Imuran (azathioprine), Sandimmune (cyclosporine A), Prograf (tacrolimus), MTX (methotrexate)]. Ulcerative Colitis (UC): Diagnosis of moderately to severely active UC. One of the following: a) trial and failure, contraindication, or intolerance to Humira (adalimumab), or (b) trial and failure, contraindication, or intolerance to treatment with at least one immunomodulator or corticosteroid (e.g., Purinethol [6-mercaptopurine], Imuran [azathioprine], aminosalicylates [e.g., mesalamine {Asacol, Pentasa, Rowasa}, olsalazine {Dipentum}, sulfasalazine {Azulfidine, Sulfazine}]).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a gastroenterologist.
<b>Coverage Duration</b>	One time
<b>Other Criteria</b>	Stelara is to be administered as an intravenous induction dose. Stelara induction dosing is in accordance with the United States Food and Drug Administration approved labeled dosing for Crohn's Disease/ulcerative colitis: 260 mg for patients weighing 55 kg or less, 390 mg for patients weighing more than 55 kg to 85 kg, or 520 mg for patients weighing more than 85 kg.

Last Updated: June 2021



## STELARA (S)

---

### **Products Affected**

- Stelara INJ 45MG/0.5ML, 90MG/ML

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	<p>Plaque psoriasis (Initial - 45mg/0.5mL): Diagnosis of moderate to severe plaque psoriasis. Plaque psoriasis (Initial - 90mg/1mL): Diagnosis of moderate to severe plaque psoriasis. Patient's weight is greater than 100 kg (220 lbs). Plaque psoriasis (Initial): One of the following: a) Trial and failure, contraindication, or intolerance (TF/C/I) to Cosentyx (secukinumab) AND one of the following: Enbrel (etanercept), Humira (adalimumab), or Skyrizi (risankizumab), OR b) for continuation of prior therapy. Psoriatic arthritis (PsA) (Initial - 45mg/0.5mL): Diagnosis of active PsA. PsA (Initial - 90mg/1mL): Diagnosis of active PsA. Patient's weight is greater than 100 kg (220 lbs). Diagnosis of co-existent moderate to severe psoriasis. PsA (Initial): One of the following: a) TF/C/I to Cosentyx (secukinumab) AND one of the following: Enbrel (etanercept) or Humira (adalimumab) OR b) for continuation of prior therapy. Crohn's disease (CD) (Initial): Diagnosis of moderately to severely active Crohn's disease. One of the following: a) TF/C/I to Humira (adalimumab) OR b) TF/C/I to treatment with at least one immunomodulator or corticosteroid [e.g., Purinethol (6-mercaptopurine), Imuran (azathioprine), Sandimmune (cyclosporine A), Prograf (tacrolimus), MTX (methotrexate)], OR c) for continuation of prior therapy. Ulcerative colitis (UC) (Initial): Diagnosis of moderately to severely active UC. One of the following: a) TF/C/I to Humira (adalimumab) OR b) TF/C/I to treatment with at least one immunomodulator or corticosteroid (e.g., Purinethol [6-mercaptopurine], Imuran [azathioprine], aminosaliclates [e.g., mesalamine {Asacol, Pentasa, Rowasa}, olsalazine {Dipentum}, sulfasalazine {Azulfidine, Sulfazine}]), OR c) for continuation of prior therapy.</p>
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Plaque psoriasis (initial): Prescribed by or in consultation with a dermatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist. CD and UC (initial): Prescribed by or in consultation with a gastroenterologist.
<b>Coverage Duration</b>	All uses (Initial, reauth): 12 months

Last Updated: June 2021

<b>Other Criteria</b>	Reauth (PsA, CD, UC): Documentation of positive clinical response to therapy. Reauth (plaque psoriasis): Documentation of positive clinical response to therapy as evidenced by one of the following: reduction in the body surface area (BSA) involvement from baseline, OR improvement in symptoms (eg, pruritus, inflammation) from baseline.
-----------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

# STIVARGA (S)

## Products Affected

- Stivarga

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Metastatic colorectal cancer (mCRC): All of the following: 1) diagnosis of mCRC, AND 2) trial and failure, contraindication or intolerance to fluoropyrimidine-, oxaliplatin- and irinotecan-based chemotherapy (e.g., FOLFOX, FOLFIRI, FOLFOXIRI), AND 3) trial and failure, contraindication or intolerance to an anti-VEGF therapy (e.g., Avastin [bevacizumab]), AND 4) one of the following: a) RAS mutation, OR b) both of the following: RAS wild-type (RAS mutation negative tumor) and trial and failure, contraindication or intolerance to an anti-EGFR therapy [e.g., Vectibix (panitumumab), Erbitux (cetuximab)]. Gastrointestinal stromal tumor (GIST): All of the following: 1) diagnosis of locally advanced, unresectable or metastatic GIST, AND 2) trial and failure, contraindication or intolerance to both of the following: a) Gleevec (imatinib mesylate), AND b) Sutent (sunitinib malate). Hepatocellular Carcinoma (HCC): Diagnosis of HCC. Trial and failure or intolerance to Nexavar (sorafenib tosylate).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	mCRC, GIST: Prescribed by or in consultation with an oncologist. HCC: Prescribed by or in consultation with an oncologist, hepatologist, or gastroenterologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

## STRENSIQ (S)

---

### Products Affected

- Strensiq

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Hypophosphatasia: Diagnosis of perinatal/infantile or juvenile-onset hypophosphatasia AND for patients requesting the 80 mg/0.8 mL vial only: Patient's weight is greater than or equal to 40 kg.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Hypophosphatasia: Prescribed by or in consultation with a specialist experienced in the treatment of inborn errors of metabolism or endocrinologist
<b>Coverage Duration</b>	Hypophosphatasia: 12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

312



## SUPPRELIN LA (S)

---

### Products Affected

- Supprelin La

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Central Precocious Puberty (CPP) (initial): Diagnosis of CPP (idiopathic or neurogenic). Early onset of secondary sexual characteristics in females less than age 8 or males less than age 9. Advanced bone age of at least one year compared with chronologic age. One of the following: a) patient has undergone gonadotropin-releasing hormone agonist (GnRHa) testing AND Peak luteinizing hormone (LH) level above pre-pubertal range, or b) patient has a random LH level in the pubertal range.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	CPP (initial, reauth): Prescribed by or in consultation with a pediatric endocrinologist.
<b>Coverage Duration</b>	CPP (init, reauth): 12 months
<b>Other Criteria</b>	CPP (reauthorization): LH levels have been suppressed to pre-pubertal levels.

Last Updated: June 2021

Formulary ID: 21381, Version 14

313

## SUTENT (S)

---

### Products Affected

- Sutent

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Renal cell carcinoma: Diagnosis of advanced or metastatic renal cell carcinoma. Gastrointestinal stromal tumor (GIST): Diagnosis of GIST after disease progression on, or contraindication or intolerance to Gleevec (imatinib). Pancreatic neuroendocrine tumors: Diagnosis of progressive, well-differentiated pancreatic neuroendocrine tumor that is unresectable locally advanced or metastatic disease. Adjuvant treatment of renal cell carcinoma: Diagnosis of renal cell carcinoma (RCC). Used as adjuvant therapy. Patient is at high risk of recurrent RCC following nephrectomy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses: Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	All uses: 12 months
<b>Other Criteria</b>	All Indications: Approve for continuation of prior therapy.

Last Updated: June 2021

## SYLATRON (S)

---

### Products Affected

- Sylatron

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Melanoma: Diagnosis of melanoma with microscopic or gross nodal involvement AND The prescribed medication will be used as adjuvant therapy within 84 days of definitive surgical resection, including complete lymphadenectomy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist or dermatologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

315

## SYLVANT (S)

---

### Products Affected

- Sylvant

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multicentric Castleman's disease (MCD) (Initial): Diagnosis of MCD. Patient is human immunodeficiency virus (HIV) negative and human herpesvirus-8 (HHV-8) negative.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	MCD (initial): Prescribed by or in consultation with hematologist/oncologist or rheumatologist.
<b>Coverage Duration</b>	MCD (initial, reauth): 6 months
<b>Other Criteria</b>	MCD (reauth): Documentation of positive clinical response to therapy. Patient is HIV negative and HHV-8 negative.

Last Updated: June 2021

Formulary ID: 21381, Version 14

316

## SYMDEKO (S)

---

### Products Affected

- Symdeko TBPK 75MG; 50MG

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Initial: Diagnosis of cystic fibrosis. One of the following: 1) Patient is homozygous for the F508del mutation in the CF transmembrane conductance regulator (CFTR) gene as detected by a U.S. Food and Drug Administration (FDA)-cleared cystic fibrosis mutation test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) OR 2) Patient has at least one mutation in the CFTR gene that is responsive to tezacaftor/ivacaftor based on in vitro data and/or clinical evidence as detected by a U.S. Food and Drug Administration (FDA)-cleared cystic fibrosis mutation test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
<b>Age Restrictions</b>	Initial: Patient is 6 years of age or older
<b>Prescriber Restrictions</b>	Initial: Prescribed by or in consultation with a pulmonologist or specialist affiliated with a CF care center
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Reauth: Documentation of a positive clinical response to therapy (e.g., improvement in lung function or decreased number of pulmonary exacerbations).

Last Updated: June 2021

Formulary ID: 21381, Version 14

317

# SYNAGIS (S)

---

## **Products Affected**

- Synagis INJ 100MG/ML,  
50MG/0.5ML

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	<p>Patient will use palivizumab for immunoprophylaxis of respiratory syncytial virus (RSV) during the peak months of infection in the patient's geographic region AND Patient meets one of the following criteria: 1) Infants born at 28 weeks, six days gestation or earlier and who are younger than 12 months of age at the start of the RSV season OR 2) Diagnosis of chronic lung disease of prematurity, born before 32 weeks, 0 days gestation, received greater than 21% oxygen for at least the first 28 days after birth, and one of the following: a) 12 months of age or younger at the start of the RSV season OR b) greater than 12 months of age to 24 months of age at the start of the RSV season and received medical support (i.e., chronic corticosteroid therapy, diuretic therapy, or supplemental oxygen) within 6 months before the start of the second RSV season. OR 3) Patient is 12 months of age or younger at the start of the RSV season and has one of the following: a) acyanotic heart failure that will require a cardiac surgical procedure and the patient is receiving medication to control congestive heart failure, OR b) moderate to severe pulmonary hypertension OR c) cyanotic heart defect. OR 4) patient is younger than 24 months of age and will or has undergone a cardiac transplantation during the RSV season. OR 5) Patient is 12 months of age or younger at the start of the RSV season with a congenital abnormality or neuromuscular disorder and has an impaired ability to clear secretions from the upper airway due to an ineffective cough. OR 6) Patient is younger than 24 months of age with a lymphocyte count below the normal range for patient's age and has received or will receive a solid organ transplant, hematopoietic stem cell transplant recipient, or chemotherapy during the RSV season.</p>
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a pediatric specialist (i.e., pulmonologist, neonatologist, neurologist, cardiologist, pediatric intensivist, or infectious disease specialist).
<b>Coverage Duration</b>	5 months (5 doses) during RSV season.

Last Updated: June 2021

<b>Other Criteria</b>	N/A
-----------------------	-----



## SYNDROS (S)

---

### Products Affected

- Syndros

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Nausea and Vomiting Associated with Cancer Chemotherapy (CINV): Patient is receiving cancer chemotherapy. Trial and failure, contraindication, or intolerance (TF/C/I) to a 5HT-3 receptor antagonist (eg, Anzemet [dolasetron], Kytril [granisetron], or Zofran [ondansetron]). TF/C/I to one of the following: Compazine (prochlorperazine), Decadron (dexamethasone), Haldol (haloperidol), Zyprexa (olanzapine). AIDS anorexia: Diagnosis of anorexia with weight loss in patients with AIDS. Patient is on antiretroviral therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	CINV: 6 months. AIDS anorexia: 3 months.
Other Criteria	Subject to Part B vs. Part D review. CINV: Approve for continuation of therapy for treatment covered under Part B when patient is receiving chemotherapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

321

## SYNRIBO (S)

---

### Products Affected

- Synribo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic myelogenous leukemia (CML): Diagnosis of CML in the chronic or accelerated phase AND Patient has tried and has had resistance, relapse, inadequate response, intolerance or is contraindicated to TWO tyrosine kinase inhibitors (i.e., Gleevec [imatinib], Sprycel, Tasigna, and Bosulif, Iclusig).
Age Restrictions	CML: 18 years of age or older
Prescriber Restrictions	CML: Prescribed by or in consultation with a hematologist/oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

322

## SYPRINE (S)

---

### Products Affected

- Clovique
- Trientine Hydrochloride

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Initial: Diagnosis of Wilson's disease (i.e., hepatolenticular degeneration). Trial and failure, contraindication, or intolerance to a penicillamine product (e.g., Depen, Cuprimine)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Reauth: Documentation of a positive clinical response to therapy

Last Updated: June 2021

Formulary ID: 21381, Version 14

323

# TABRECTA (S)

---

## Products Affected

- Tabrecta

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of non-small cell lung cancer (NSCLC). Disease is one of the following: recurrent, advanced, metastatic. Presence of mesenchymal-epithelial transition (MET) exon 14 skipping positive tumors as detected with an FDA-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

324

# TAFAMIDIS (S)

## Products Affected

- Vyndamax
- Vyndaqel

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Transthyretin-mediated amyloidosis with cardiomyopathy (ATTR-CM) (initial): Diagnosis of transthyretin-mediated amyloidosis with cardiomyopathy (ATTR-CM). One of the following: 1) Patient has a transthyretin (TTR) mutation (e.g., V122I), 2) Cardiac or noncardiac tissue biopsy demonstrating histologic confirmation of TTR amyloid deposits, OR 3) All of the following: i) echocardiogram or cardiac magnetic resonance imaging suggestive of amyloidosis, ii) scintigraphy scan suggestive of cardiac TTR amyloidosis, and iii) absence of light-chain amyloidosis. One of the following: 1) History of heart failure (HF), with at least one prior hospitalization for HF, OR 2) Presence of clinical signs and symptoms of HF (e.g., dyspnea, edema). Patient has New York Heart Association (NYHA) Functional Class I, II, or III heart failure.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	ATTR-CM (initial, reauth): Prescribed by or in consultation with a cardiologist
<b>Coverage Duration</b>	ATTR-CM (initial, reauth): 12 months
<b>Other Criteria</b>	ATTR-CM (reauth): Documentation of positive clinical response to therapy. Patient continues to have New York Heart Association (NYHA) Functional Class I, II, or III heart failure.

Last Updated: June 2021

## TAFINLAR (S)

---

### Products Affected

- Tafinlar

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	<p>Melanoma: Diagnosis of unresectable or metastatic melanoma AND cancer is BRAF V600E mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) OR both of the following: cancer is BRAF V600E or V600K mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) and medication is used in combination with Mekinist (trametinib). Adjuvant Treatment for Melanoma: Diagnosis of melanoma. Cancer is BRAF V600E or V600K mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Involvement of lymph nodes following complete resection. Used as adjunctive therapy. Medication is used in combination with Mekinist (trametinib). Non-small Cell Lung Cancer (NSCLC): Diagnosis of metastatic non-small cell lung cancer AND cancer is BRAF V600E mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) AND medication is used in combination with Mekinist (trametinib). Anaplastic Thyroid Cancer (ATC): Diagnosis of locally advanced or metastatic anaplastic thyroid cancer. Cancer is BRAF V600E mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Cancer may not be treated with standard locoregional treatment options. Medication is used in combination with Mekinist (trametinib) .</p>
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Last Updated: June 2021

Formulary ID: 21381, Version 14



# TAGRISO (S)

---

## Products Affected

- Tagrisso

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Non-small cell lung cancer (NSCLC): One of the following: A) All of the following: Diagnosis of metastatic NSCLC. One of the following: 1) Patient has known active epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R mutations as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA), OR 2) Both of the following: a) Patient has known active EGFR T790M mutation as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) and b) Patient has experienced disease progression on or after one of the following EGFR Tyrosine Kinase Inhibitors (TKIs): Gilotrif (afatinib), Iressa (gefitinib), Tarceva (erlotinib). OR B) All of the following: Diagnosis of NSCLC. Patient has known active epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R mutations as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Both of the following: 1) Patient is receiving as adjuvant therapy, and 2) Patient has had a complete surgical resection of the primary NSCLC tumor.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

# TAKHZYRO (S)

---

## Products Affected

- Takhzyro

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Prophylaxis of hereditary angioedema (HAE) attacks: Diagnosis of HAE. For prophylaxis against HAE attacks.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	HAE (prophylaxis): Prescribed by or in consultation with an immunologist, allergist, or rheumatologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

331

# TALZENNA (S)

---

## Products Affected

- Talzenna

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Breast cancer: Diagnosis of breast cancer. Disease is one of the following: a) locally advanced or b) metastatic. Presence of a deleterious or suspected deleterious germline BRCA-mutation (gBRCAm) as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Disease is human epidermal growth factor receptor 2 (HER2)-negative.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

332

# TARCEVA (S)

---

## Products Affected

- Erlotinib Hydrochloride

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Non-small cell lung cancer (NSCLC): Diagnosis of locally advanced or metastatic (Stage III or IV) NSCLC AND Patient has known active epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutation as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Pancreatic Cancer: Diagnosis of locally advanced, unresectable, or metastatic pancreatic cancer AND erlotinib will be used in combination with gemcitabine.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses: Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	All uses: 12 months
<b>Other Criteria</b>	All Indications: Approve for continuation of prior therapy.

Last Updated: June 2021

# TARGRETIN (S)

---

## Products Affected

- Bexarotene
- Targretin GEL

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cutaneous T-Cell Lymphoma (CTCL): Diagnosis of CTCL. Trial and failure, contraindication, or intolerance to at least one prior therapy (including skin-directed therapies [eg, corticosteroids {ie, clobetasol, diflorasone, halobetasol, augmented betamethasone dipropionate}] or systemic therapies [eg, interferons]).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist or dermatologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

334

# TASIGNA (S)

---

## Products Affected

- Tassigna

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chronic myelogenous leukemia (CML): Diagnosis of Ph+/BCR ABL CML
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist or hematologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

335

## TAZVERIK (S)

---

### Products Affected

- Tazverik

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Epithelioid sarcoma: Diagnosis of epithelioid sarcoma. Disease is one of the following: metastatic or locally advanced. Patient is not eligible for complete resection. Follicular lymphoma: Diagnosis of follicular lymphoma. Disease is one of the following: relapsed or refractory.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Epithelioid sarcoma: Prescribed by or in consultation with an oncologist. Follicular lymphoma: Prescribed by or in consultation with an oncologist or hematologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

336



## TECENTRIQ (S)

---

### Products Affected

- Tecentriq

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 Months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

## TECFIDERA (S)

---

### Products Affected

- Dimethyl Fumarate CPDR
- Dimethyl Fumarate Starterpack  
MISC 0
- Tecfidera
- Tecfidera Starter Pack

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple Sclerosis (MS): Diagnosis of a relapsing form of MS (eg, clinically isolated syndrome, relapsing-remitting disease, secondary progressive disease, including active disease with new brain lesions).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

338

## TEGSEDI (S)

---

### Products Affected

- Tegsedi

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Hereditary transthyretin-mediated amyloidosis (hATTR amyloidosis) (initial): Diagnosis of hATTR amyloidosis with polyneuropathy. Patient has a transthyretin (TTR) mutation (e.g., V30M). One of the following: 1) Patient has a baseline polyneuropathy disability (PND) score less than or equal to IIIb, 2) Patient has baseline familial amyloidotic polyneuropathy (FAP) stage of 1 or 2, OR 3) Patient has a baseline neuropathy impairment score (NIS) between 10 and 130. Presence of clinical signs and symptoms of the disease (e.g., peripheral/autonomic neuropathy).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	hATTR amyloidosis (initial): Prescribed by or in consultation with a neurologist
<b>Coverage Duration</b>	hATTR amyloidosis (initial, reauth): 12 months
<b>Other Criteria</b>	hATTR amyloidosis (reauth): Patient has demonstrated a benefit from therapy (e.g., improved neurologic impairment, slowing of disease progression, quality of life assessment). One of the following: 1) Patient continues to have a PND score less than or equal to IIIb, 2) Patient continues to have a FAP stage of 1 or 2, OR 3) Patient continues to have a NIS between 10 and 130.

Last Updated: June 2021

Formulary ID: 21381, Version 14

339

## TEPMETKO (S)

---

### Products Affected

- Tepmetko

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Non-small cell lung cancer (NSCLC): Diagnosis of NSCLC. Disease is metastatic. Presence of mesenchymal-epithelial transition (MET) exon 14 skipping alterations.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

340

# TERIPARATIDE (S)

---

## **Products Affected**

- Forteo INJ 620MCG/2.48ML
- Teriparatide

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Postmenopausal osteoporosis or osteopenia or men with primary or hypogonadal osteoporosis or osteopenia (initial): Diagnosis of one of the following: a) postmenopausal osteoporosis or osteopenia or b) primary or hypogonadal osteoporosis or osteopenia. One of the following: Set I) Both of the following: A) Bone mineral density (BMD) T-score of -2.5 or lower in the lumbar spine, femoral neck, total hip, or radius (one-third radius site) AND B) One of the following: 1) history of low-trauma fracture of the hip, spine, proximal humerus, pelvis, or distal forearm, or 2) trial and failure, contraindication, or intolerance (TF/C/I) to one osteoporosis treatment (e.g., alendronate, risedronate, zoledronic acid, Prolia [denosumab]), or Set II) Both of the following: A) BMD T-score between -1.0 and -2.5 in the lumbar spine, femoral neck, total hip, or radius (one-third radius site) AND B) One of the following: 1) history of low-trauma fracture of the hip, spine, proximal humerus, pelvis, or distal forearm, or 2) both of the following: i) TF/C/I to one osteoporosis treatment (e.g., alendronate, risedronate, zoledronic acid, Prolia [denosumab]) and ii) One of the following FRAX 10-year probabilities: a) Major osteoporotic fracture at 20% or more in the U.S., or the country-specific threshold in other countries or regions, or b) Hip fracture at 3% or more in the U.S., or the country-specific threshold in other countries or regions. Glucocorticoid-Induced Osteoporosis: See Other Criteria section.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	All uses (initial): 24 months. All uses (reauth): 12 months.
<b>Other Criteria</b>	Glucocorticoid-Induced Osteoporosis (initial): Diagnosis of glucocorticoid-induced osteoporosis. History of prednisone or its equivalent at a dose greater than or equal to 5mg/day for greater than or equal to 3 months. One of the following: 1) BMD T-score less than or equal to -2.5 based on BMD measurements from lumbar spine, femoral neck, total hip, or radius (one-third radius site), or 2) One of the following

Last Updated: June 2021

---

FRAX 10-year probabilities: a) Major osteoporotic fracture at 20% or more in the U.S., or the country-specific threshold in other countries or regions, or b) Hip fracture at 3% or more in the U.S., or the country-specific threshold in other countries or regions, or 3) History of one of the following fractures resulting from minimal trauma: vertebral compression fx, fx of the hip, fx of the distal radius, fx of the pelvis, or fx of the proximal humerus. TF/C/I to one bisphosphonate (e.g., alendronate). All uses (initial, reauth): One of the following: 1) Treatment duration of parathyroid hormones [e.g., teriparatide, Tymlos (abaloparatide)] has not exceeded a total of 24 months during the patient's lifetime, or 2) Patient remains at or has returned to having a high risk for fracture despite a total of 24 months of use of parathyroid hormones [e.g., teriparatide, Tymlos (abaloparatide)].

---

# TESTOSTERONE (S)

---

## Products Affected

- Androderm PT24 2MG/24HR,  
4MG/24HR
- Testosterone GEL 1.62%,  
25MG/2.5GM, 50MG/5GM
- Testosterone Cypionate INJ  
100MG/ML, 200MG/ML
- Testosterone Pump



<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Hypogonadism (HG) (Initial): Diagnosis (dx) of HG AND male patient at birth AND one of the following: 1) Two pre-treatment serum total testosterone (T) levels less than 300 ng/dL (10.4 nmol/L) or less than the reference range for the lab OR 2) Both of the following: a) Has a condition that may cause altered sex-hormone binding globulin (SHBG) (eg, thyroid disorder, HIV disease, liver disorder, diabetes, obesity), and b) one pre-treatment calculated free or bioavailable T level less than 5 ng/dL (0.17 nmol/L) or less than reference range for the lab, OR 3) History of bilateral orchiectomy, panhypopituitarism, or a genetic disorder known to cause HG (eg, congenital anorchia, Klinefelter's syndrome). Gender Dysphoria (GD) (off-label): Dx of GD. Patient is a female-to-male transsexual.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	HG(init): (New to T tx:6 mo. New to plan and cont T tx:12 mo), (reauth): 12 mo. GD: 12 mo.
<b>Other Criteria</b>	HG (Reauth): 1) Follow-up total serum T level within or below the normal limits of the reporting lab, or 2) Follow-up total serum T level outside of upper limits of normal for the reporting lab and the dose is adjusted, OR 3) Has a condition that may cause altered SHBG (eg, thyroid disorder, HIV disease, liver disorder, diabetes, obesity), and one of the following: Follow-up calculated free or bioavailable T level within or below the normal limits of the reporting lab, or follow-up calculated free or bioavailable T level outside of upper limits of normal for the reporting lab and the dose is adjusted.

Last Updated: June 2021

Formulary ID: 21381, Version 14

345

# TESTOSTERONE ENANTHATE (S)

---

## Products Affected

- Testosterone Enanthate INJ

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Hypogonadism (HG) (Initial): Diagnosis (dx) of HG AND male patient at birth AND one of the following: 1) Two pre-treatment serum total testosterone (T) levels less than 300 ng/dL (10.4 nmol/L) or less than the reference range for the lab OR 2) Both of the following: a) Has a condition that may cause altered sex-hormone binding globulin (SHBG) (eg, thyroid disorder, HIV disease, liver disorder, diabetes, obesity), and b) one pre-treatment calculated free or bioavailable T level less than 5 ng/dL (0.17 nmol/L) or less than reference range for the lab, OR 3) History of bilateral orchiectomy, panhypopituitarism, or a genetic disorder known to cause HG (eg, congenital anorchia, Klinefelter's syndrome). Delayed puberty (DP): Dx of DP AND male patient at birth. Breast cancer (BC): Dx of inoperable BC AND used for palliative treatment AND female patient at birth. Gender Dysphoria (GD) (off-label): Dx of GD. Patient is a female-to-male transsexual.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	HG(init): (New to T tx:6 mo. Cont T tx:12 mo), (reauth): 12 mo. BC, GD: 12 mo. DP: 6 mo.
<b>Other Criteria</b>	HG (Reauth): 1) Follow-up total serum T level within or below the normal limits of the reporting lab, or 2) Follow-up total serum T level outside of upper limits of normal for the reporting lab and the dose is adjusted, OR 3) Has a condition that may cause altered SHBG (eg, thyroid disorder, HIV disease, liver disorder, diabetes, obesity), and one of the following: Follow-up calculated free or bioavailable T level within or below the normal limits of the reporting lab, or follow-up calculated free or bioavailable T level outside of upper limits of normal for the reporting lab and the dose is adjusted.

Last Updated: June 2021

Formulary ID: 21381, Version 14

347

# THALOMID (S)

---

## Products Affected

- Thalomid

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple myeloma (MM): Diagnosis of MM. Used in combination with dexamethasone, unless the patient has an intolerance to steroids. Erythema nodosum leprosum (ENL): Diagnosis of moderate to severe ENL with cutaneous manifestations. Thalomid is not used as monotherapy if moderate to severe neuritis is present.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	MM: Prescribed by or in consultation with an oncologist/hematologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

348

## TIBSOVO (S)

---

### Products Affected

- Tibsovo

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Relapsed or refractory Acute Myeloid Leukemia (AML): Diagnosis of AML. Disease is relapsed or refractory. Patient has an isocitrate dehydrogenase-1 (IDH1) mutation as detected by a U.S. Food and Drug Administration (FDA)-approved test (e.g., Abbott RealTime IDH1 assay) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Newly-Diagnosed AML: Diagnosis of newly-diagnosed AML. Patient has an isocitrate dehydrogenase-1 (IDH1) mutation as detected by an FDA-approved test (e.g., Abbott RealTime IDH1 assay) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). One of the following: 1) patient is greater than or equal to 75 years old OR 2) patient has comorbidities that preclude use of intensive induction chemotherapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist/oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

# TIGLUTIK (S)

---

## Products Affected

- Tiglutik

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Amyotrophic Lateral Sclerosis (ALS): Diagnosis of ALS. Trial and failure or intolerance to generic riluzole tablets.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

350

## TOPICAL RETINOID (S)

---

### Products Affected

- Avita
- Tretinoin CREA
- Tretinoin GEL

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Acne vulgaris: Diagnosis of acne vulgaris (i.e., acne).
<b>Age Restrictions</b>	PA applies to members 26 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

351

# TRAZIMERA (S)

---

## Products Affected

- Trazimera

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Breast cancer: Diagnosis of human epidermal growth factor receptor 2 (HER2)-overexpressing breast cancer. One of the following treatment regimens: a) As adjuvant treatment, b) metastatic disease and one of the following: 1) used in combination with a taxane (eg, docetaxel, paclitaxel), or 2) used as a single agent in a patient who has received one or more chemotherapy regimens for metastatic disease, or c) used in combination with Perjeta (pertuzumab). Gastric Cancer: Diagnosis of HER2-overexpressing gastric or gastroesophageal junction adenocarcinoma (locally advanced, recurrent, or metastatic). Used in combination with one of the following treatment regimens: a) Platinol (cisplatin) and Adrucil (5-fluorouracil), or b) Platinol (cisplatin) and Xeloda (capecitabine).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses: Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

352



## TRELSTAR (S)

---

### Products Affected

- Trelstar Mixject INJ 11.25MG, 22.5MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Prostate Cancer: Diagnosis of advanced or metastatic prostate cancer. Trial and failure, contraindication, or intolerance to any brand Lupron formulation.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

353

# TRIKAFTA (S)

---

## Products Affected

- Trikafta

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cystic Fibrosis (CF) (initial): Diagnosis of CF. Patient has at least one of the following mutations in the cystic fibrosis transmembrane conductance regulator (CFTR) gene as detected by an FDA-cleared cystic fibrosis mutation test or a test performed at a Clinical Laboratory Improvement Amendments (CLIA)-approved facility: F508del mutation OR a mutation in the CFTR gene that is responsive based on in vitro data.
<b>Age Restrictions</b>	CF (initial): 12 years of age or older.
<b>Prescriber Restrictions</b>	CF (initial): Prescribed by or in consultation with a pulmonologist or specialist affiliated with a CF care center.
<b>Coverage Duration</b>	CF (initial, reauth): 12 months
<b>Other Criteria</b>	CF (reauth): Documentation of positive clinical response to therapy (e.g., improvement in lung function [percent predicted forced expiratory volume in one second {PPFEV1}] or decreased number of pulmonary exacerbations).

Last Updated: June 2021

Formulary ID: 21381, Version 14

354

## TRIPTODUR (S)

---

### Products Affected

- Triptodur

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Central Precocious Puberty (CPP) (initial): Diagnosis of CPP (idiopathic or neurogenic). Early onset of secondary sexual characteristics in females less than age 8 or males less than age 9. Advanced bone age of at least one year compared with chronologic age. One of the following: a) patient has undergone gonadotropin-releasing hormone agonist (GnRHa) testing AND Peak luteinizing hormone (LH) level above pre-pubertal range, or b) patient has a random LH level in the pubertal range. Trial and failure or intolerance to Lupron Depot-Ped.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	CPP (initial): Prescribed by or in consultation with a pediatric endocrinologist.
<b>Coverage Duration</b>	CPP (Initial, reauth): 12 months
<b>Other Criteria</b>	CPP (reauthorization): LH levels have been suppressed to pre-pubertal levels.

Last Updated: June 2021

Formulary ID: 21381, Version 14

355

# TUKYSA (S)

---

## Products Affected

- Tukysa

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Breast cancer: Diagnosis of breast cancer. Disease is one of the following: a) advanced unresectable or b) metastatic. Disease is human epidermal growth factor receptor 2 (HER2)-positive. Used in combination with trastuzumab and capecitabine. Patient has received one or more prior anti-HER2 based regimens (e.g., trastuzumab, pertuzumab, ado-trastuzumab emtansine).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

356

## TURALIO (S)

---

### Products Affected

- Turalio

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Tenosynovial Giant Cell Tumor (TGCT): Diagnosis of TGCT. Patient is symptomatic. Patient is not a candidate for surgery due to worsening functional limitation or severe morbidity with surgical removal.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist/oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

357

# TYKERB (S)

---

## Products Affected

- Lapatinib Ditosylate
- Tykerb

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Breast Cancer: Diagnosis of human epidermal growth factor receptor 2 (HER2)-positive metastatic or recurrent breast cancer. Used in combination with one of the following: Trastuzumab, Xeloda (capecitabine), or aromatase inhibitors [eg, Aromasin (exemestane), Femara (letrozole), Arimidex (anastrozole)].
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

358

# **TYMLOS (S)**

---

## **Products Affected**

- Tymlos

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of postmenopausal osteoporosis or osteopenia. One of the following: Set I) Both of the following: A) Bone mineral density (BMD) T-score of -2.5 or lower in the lumbar spine, femoral neck, total hip, or radius (one-third radius site) AND B) One of the following: 1) history of low-trauma fracture of the hip, spine, proximal humerus, pelvis, or distal forearm, or 2) trial and failure, contraindication, or intolerance (TF/C/I) to one osteoporosis treatment (e.g., alendronate, risedronate, zoledronic acid, Prolia [denosumab]), or Set II) Both of the following: A) BMD T-score between -1.0 and -2.5 in the lumbar spine, femoral neck, total hip, or radius (one-third radius site) AND B) One of the following: 1) history of low-trauma fracture of the hip, spine, proximal humerus, pelvis, or distal forearm, or 2) both of the following: i) TF/C/I to one osteoporosis treatment (e.g., alendronate, risedronate, zoledronic acid, Prolia [denosumab]) and ii) one of the following FRAX (Fracture Risk Assessment Tool) 10-year probabilities: a) major osteoporotic fracture at 20% or more in the U.S., or the country-specific threshold in other countries or regions, or b) hip fracture at 3% or more in the U.S., or the country-specific threshold in other countries or regions. Treatment duration of parathyroid hormones (e.g., teriparatide, Tymlos [abaloparatide]) has not exceeded a total of 24 months during the patient's lifetime.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	24 months (max 24 months of therapy per lifetime)
<b>Other Criteria</b>	N/A

Last Updated: June 2021



# **TYSABRI (S)**

---

## **Products Affected**

- Tysabri

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple Sclerosis (MS): Diagnosis of a relapsing form of MS (eg, relapsing-remitting MS, secondary-progressive MS with relapses). One of the following: 1) Trial and failure, contraindication, or intolerance (TF/C/I) to one of the following disease-modifying therapies for MS: Aubagio (teriflunomide), Avonex (interferon beta-1a), Betaseron (interferon beta-1b), Copaxone/Glatopa (glatiramer acetate), Extavia (interferon beta-1b), Gilenya (fingolimod), Lemtrada (alemtuzumab), Mavenclad (cladribine), Mayzent (siponimod), Ocrevus (ocrelizumab), Plegridy (peginterferon beta-1a), Rebif (interferon beta-1a), Tecfidera (dimethyl fumarate), 2) Patient is not a candidate for any of the drugs listed as prerequisites due to the severity of their MS, or 3) for continuation of prior therapy. Not used in combination with another disease-modifying therapy for MS. Crohn's Disease (CD) (initial): Diagnosis of moderate to severe CD with evidence of inflammation (eg, elevated C-reactive protein [CRP], elevated erythrocyte sedimentation rate, presence of fecal leukocytes). TF/C/I to one of the following conventional therapies: corticosteroids, 6-mercaptopurine (6MP [Purinethol], azathioprine (Imuran), methotrexate, aminosalicylates (eg, sulfasalazine, mesalamine, olsalazine). TF/C/I to a TNF-inhibitor (eg, Cimzia [certolizumab pegol], Humira [adalimumab], Remicade [infliximab]). CD (initial and reauth): Not used in combination with an immunosuppressant (eg, 6-MP, azathioprine, cyclosporine, or methotrexate). Not used in combination with a TNF-inhibitor (eg, Enbrel [etanercept], Humira [adalimumab], or Remicade [infliximab]).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	MS: 12mo. CD (Init): 3 mo. CD (Reauth): 6 mo if not on steroids. Otherwise, 3 mo.
<b>Other Criteria</b>	CD (reauth): Documentation of positive clinical response (eg, improved disease activity index) to therapy.

Last Updated: June 2021

## UDENYCA (S)

### Products Affected

- Udenyca

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	<p>Febrile neutropenia (FN) prophylaxis: Patient will be receiving prophylaxis for FN due to one of the following: 1) Patient is receiving National Cancer Institute’s Breast Intergroup, INT C9741 dose dense chemotherapy protocol for primary breast cancer, 2) patient is receiving a dose-dense chemotherapy regimen for which the incidence of FN is unknown, 3) patient is receiving chemotherapy regimen(s) associated with greater than 20% incidence of FN, 4) both of the following: a) patient is receiving chemotherapy regimen(s) associated with 10-20% incidence of FN, AND b) patient has one or more risk factors associated with chemotherapy-induced infection, FN, or neutropenia, OR 5) Both of the following: a) patient is receiving myelosuppressive anticancer drugs associated with neutropenia, AND b) patient has a history of FN or dose-limiting event during a previous course of chemotherapy (secondary prophylaxis). Treatment of FN (off-label): Patient has received or is receiving myelosuppressive anticancer drugs associated with neutropenia. Diagnosis of FN. Patient is at high risk for infection-associated complications. Acute radiation syndrome (ARS) (off-label): Patient was/will be acutely exposed to myelosuppressive doses of radiation (hematopoietic subsyndrome of ARS).</p>
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses: Prescribed by or in consultation with a hematologist/oncologist
<b>Coverage Duration</b>	ARS: 1 mo. FN (prophylaxis, treatment): 3 mo or duration of tx.
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Last Updated: June 2021

Formulary ID: 21381, Version 14

364

## UKONIQ (S)

---

### Products Affected

- Ukoniq

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Marginal zone lymphoma (MZL): Diagnosis of MZL. Disease is one of the following: relapsed or refractory. Patient has received at least one prior anti-CD20-based regimen (e.g., bendamustine + rituximab, bendamustine + obinutuzumab, etc.). Follicular lymphoma (FL): Diagnosis of FL. Disease is one of the following: relapsed or refractory. Patient has received at least three prior lines of systemic therapy (e.g., bendamustine + rituximab, bendamustine + obinutuzumab, etc.).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	MZL/FL: Prescribed by or in consultation with a hematologist/oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

365

# ULTOMIRIS (S)

---

## Products Affected

- Ultomiris

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Paroxysmal nocturnal hemoglobinuria (PNH) (initial): Diagnosis of PNH. Atypical hemolytic uremic syndrome (aHUS) (initial): Diagnosis of aHUS.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	PNH, aHUS (initial, reauth): 12 months
<b>Other Criteria</b>	PNH, aHUS (reauth): Documentation of positive clinical response (e.g., hemoglobin stabilization, decrease in the number of red blood cell transfusions) to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

366

## VALCHLOR (S)

---

### Products Affected

- Valchlor

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Mycosis fungoides-type cutaneous T-cell lymphoma (MF-CTCL) (initial): All of the following: 1) diagnosis of Stage IA MF-CTCL, OR diagnosis of Stage IB MF-CTCL, AND 2) patient has received at least one prior skin-directed therapy [e.g., topical corticosteroids, bexarotene topical gel (Targretin topical gel), etc.].
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist or dermatologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

367

## VARIZIG (S)

---

### Products Affected

- Varizig INJ 125UNIT/1.2ML

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	Presence of contraindications to immune globulin therapy (i.e., IgA deficiency with antibodies to IgA and a history of hypersensitivity or product specific contraindication).
<b>Required Medical Information</b>	Immune globulin is being used intramuscularly. The immune globulin is being used for passive immunization or post exposure-prophylaxis of varicella. Patient is considered a high risk individual (i.e., immune compromised, pregnant woman, newborn of mother with varicella, premature infant, and infant less than 1 year old).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	3 months (approve one dose only)
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

368



## VASCEPA (S)

---

### Products Affected

- Icosapent Ethyl
- Vascepa

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Severe Hypertriglyceridemia (init): Diagnosis (dx) of hypertriglyceridemia and patient has a pre-treatment triglyceride (TG) level greater than or equal to 500 mg/dL. Prevention of CV Events (init): Dx of hypertriglyceridemia and patient has a pre-treatment TG level of 150 to 499 mg/dL. One of the following: 1) Patient has established cardiovascular disease (CVD) (e.g., coronary artery disease, cerebrovascular or carotid disease, peripheral artery disease, etc.) OR 2) Both of the following: a) Dx of diabetes mellitus AND b) Patient has two or more risk factors for developing CVD. Medication will be used as an adjunct to maximally tolerated statin therapy unless there is a contraindication or intolerance to statin therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Initial/Reauth: 12 months
<b>Other Criteria</b>	Severe Hypertriglyceridemia (reauth): Documentation of positive clinical response to therapy. Prevention of CV Events (Reauth): Documentation of positive clinical response to therapy. Medication continues to be used as an adjunct to maximally tolerated statin therapy unless there is a contraindication or intolerance to statin therapy.

Last Updated: June 2021

## VENCLEXTA (S)

---

### Products Affected

- Venclexta
- Venclexta Starting Pack

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL): Diagnosis of CLL or SLL. Acute Myeloid Leukemia (AML): Diagnosis of newly diagnosed AML. Used in combination with azacitidine, or decitabine, or low-dose cytarabine. One of the following: 1) age 75 years or older OR 2) comorbidities that preclude use of intensive induction chemotherapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist or oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

370

## VENTAVIS (S)

---

### Products Affected

- Ventavis

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist.
<b>Coverage Duration</b>	PAH (Initial): 6 months. (Reauth): 12 months
<b>Other Criteria</b>	Subject to Part B vs D review. PAH (Reauth): Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

371

## VERZENIO (S)

---

### Products Affected

- Verzenio

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Breast Cancer: Diagnosis of advanced or metastatic breast cancer. Disease is hormone receptor (HR)-positive and human epidermal growth factor receptor 2 (HER2)-negative. One of the following: a) used in combination with an aromatase inhibitor (e.g., Arimidex [anastrozole], Aromasin [exemestane], Femara [letrozole]) and patient is a postmenopausal woman, OR b) used in combination with Faslodex (fulvestrant) and disease has progressed following endocrine therapy, OR c) used as monotherapy and disease has progressed following endocrine therapy and patient has already received at least one prior chemotherapy regimen.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

372

## VIMIZIM (S)

---

### Products Affected

- Vimizim

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Mucopolysaccharidosis (initial): Diagnosis of Mucopolysaccharidosis type IVA (MPS IVA, Morquio A syndrome) confirmed by both of the following: a) documented clinical signs and symptoms of the disease (e.g., kyphoscoliosis, genu valgum, pectus carinatum, gait disturbance, growth deficiency, etc.) and b) documented reduced fibroblast or leukocyte GALNS enzyme activity or molecular genetic testing of GALNS.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Initial, reauth: 12 months
<b>Other Criteria</b>	Reauth: Documentation of positive clinical response to therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

373

## VITRAKVI (S)

---

### Products Affected

- Vitrakvi

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Presence of solid tumors (e.g., salivary gland, soft tissue sarcoma, infantile fibrosarcoma, thyroid cancer, lung, melanoma, colon, etc.). Disease is positive for neurotrophic receptor tyrosine kinase (NTRK) gene fusion (e.g. ETV6-NTRK3, TPM3-NTRK1, LMNA-NTRK1, etc.). Disease is without a known acquired resistance mutation [e.g., TRKA G595R substitution, TRKA G667C substitution, or other recurrent kinase domain (solvent front and xDFG) mutations]. Disease is one of the following: metastatic or unresectable (including cases where surgical resection is likely to result in severe morbidity). One of the following: Disease has progressed on previous treatment (e.g., surgery, radiotherapy, or systemic therapy) OR Disease has no satisfactory alternative treatments.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

374

## VIZIMPRO (S)

---

### Products Affected

- Vizimpro

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Non-small cell lung cancer (NSCLC): Diagnosis of NSCLC. Disease is metastatic. Disease is positive for one of the following epidermal growth factor receptor (EGFR) mutations: exon 19 deletion or exon 21 L858R substitution.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

375

## VOSEVI (S)

---

### Products Affected

- Vosevi

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Criteria will be applied consistent with current AASLD/IDSA guideline. All patients: Diagnosis of chronic hepatitis C, patient is without decompensated liver disease (defined as Child-Pugh Class B or C), and not used in combination with another HCV direct acting antiviral agent [e.g., Harvoni, Zepatier].
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with one of the following: Hepatologist, Gastroenterologist, Infectious disease specialist, HIV specialist certified through the American Academy of HIV Medicine.
<b>Coverage Duration</b>	12 weeks. Criteria will be applied consistent with current AASLD/IDSA guideline.
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

376



## VOTRIENT (S)

---

### Products Affected

- Votrient

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Renal cell carcinoma (RCC): Diagnosis of advanced/metastatic RCC. Soft tissue sarcoma: Diagnosis of advanced soft tissue sarcoma and patient received at least one prior chemotherapy (e.g., ifosfamide, doxorubicin, cisplatin, dacarbazine, docetaxel, oxaliplatin, etc.).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses: Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

377

## VPRIV (S)

---

### Products Affected

- Vpriv

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Gaucher disease: Diagnosis of type 1 Gaucher disease. Patient has evidence of symptomatic disease (e.g., moderate to severe anemia, thrombocytopenia, bone disease, hepatomegaly, or splenomegaly).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Gaucher disease: 12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

378

## XALKORI (S)

---

### Products Affected

- Xalkori

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Non-small cell lung cancer (NSCLC): Diagnosis of advanced or metastatic NSCLC AND One of the following: A) Patient has an anaplastic lymphoma kinase (ALK)-positive tumor as detected with a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) or B) Patient has MET amplification- or ROS1 rearrangement-positive tumor as detected with an FDA-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Anaplastic Large Cell Lymphoma (ALCL): Diagnosis of systemic ALCL. Disease is relapsed or refractory. Patient has an anaplastic lymphoma kinase (ALK)-positive tumor as detected with a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	NSCLC: Prescribed by or in consultation with an oncologist. ALCL: Prescribed by or in consultation with an oncologist or hematologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

379

## XCOPRI (S)

---

### Products Affected

- Xcopri

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of partial onset seizures.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

380

# XELJANZ (S)

---

## Products Affected

- Xeljanz
- Xeljanz Xr

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Xeljanz tab/Xeljanz XR tab: Rheumatoid arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. Trial and failure, contraindication, or intolerance (TF/C/I) to one disease modifying antirheumatic drug (DMARD) [eg, methotrexate (Rheumatrex/Trexall), Arava (leflunomide), Azulfidine (sulfasalazine)]. Psoriatic arthritis (PsA) (Initial): Diagnosis of active PsA. PsA (initial): One of the following: TF/C/I to Enbrel (etanercept) or Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR patient has a documented needle-phobia to the degree that the patient has previously refused any injectable therapy or medical procedure (refer to DSM-IV-TR 300.29/F40.2 for specific phobia diagnostic criteria), OR for continuation of prior tofacitinib therapy. Ulcerative colitis (UC) (Initial): Diagnosis of moderately to severely active UC. Trial and failure, contraindication or intolerance to one of the following conventional therapies: 6-mercaptopurine (Purinethol), aminosalicylate [e.g., mesalamine (Asacol, Pentasa, Rowasa), olsalazine (Dipentum), sulfasalazine (Azulfidine, Sulfazine)], azathioprine (Imuran), or corticosteroids (e.g., prednisone, methylprednisolone). Trial and failure, contraindication, or intolerance to Humira (adalimumab), OR patient has a documented needle-phobia to the degree that the patient has previously refused any injectable therapy or medical procedure (refer to DSM-V-TR F40.2 for specific phobia diagnostic criteria), OR for continuation of prior tofacitinib therapy. All indications: Patient is not receiving tofacitinib in combination with a potent immunosuppressant (eg, azathioprine, cyclosporine).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	RA, PJIA (initial): Prescribed by or in consultation with a rheumatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist. UC (initial): Prescribed by or in consultation with a gastroenterologist.
<b>Coverage Duration</b>	RA/PJIA/PsA (initial, reauth): 12 months. UC (init): 4 mo. UC (reauth): 12 mo.

Last Updated: June 2021

<b>Other Criteria</b>	Xeljanz: Polyarticular course juvenile idiopathic arthritis (PJIA) (Initial): Diagnosis of active polyarticular course juvenile idiopathic arthritis. One of the following: TF/C/I to Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR patient has a documented needle-phobia to the degree that the patient has previously refused any injectable therapy or medical procedure (refer to DSM-V-TR 300.29/F40.2 for specific phobia diagnostic criteria), OR for continuation of prior tofacitinib therapy. All Indications (Reauth): Documentation of positive clinical response to tofacitinib therapy. Patient is not receiving tofacitinib in combination with a potent immunosuppressant (eg, azathioprine, cyclosporine).
-----------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

## XENAZINE (S)

---

### Products Affected

- Tetrabenazine

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chorea associated with Huntington's Disease (HD) (Initial): Diagnosis of chorea in patients with Huntington's disease. Tardive dyskinesia (Initial): Diagnosis of tardive dyskinesia. One of the following: 1) Patient has persistent symptoms of tardive dyskinesia despite a trial of dose reduction, tapering, or discontinuation of the offending medication, OR 2) Patient is not a candidate for a trial of dose reduction, tapering, or discontinuation of the offending medication. Tourette's syndrome (Initial): Patient has tics associated with Tourette's syndrome. Failure, contraindication, or intolerance to Haldol (haloperidol).
Age Restrictions	N/A
Prescriber Restrictions	HD (Initial): Prescribed by or in consultation with a neurologist. Tardive dyskinesia, Tourette's syndrome (Initial): Prescribed by or in consultation with neurologist or psychiatrist.
Coverage Duration	All uses: (initial) 3 months. (Reauth) 12 months.
Other Criteria	All indications (Reauth): Documentation of clinical response and benefit from therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

384



## XGEVA (S)

---

### Products Affected

- Xgeva

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple Myeloma (MM)/Bone metastasis from solid tumors (BMST): One of the following: 1) Diagnosis of multiple myeloma OR 2) diagnosis of solid tumors (eg, breast cancer, kidney cancer, lung cancer, prostate cancer, thyroid cancer), AND documented evidence of one or more metastatic bone lesions. Giant cell tumor of bone (GCTB): Both of the following: 1) diagnosis of giant cell tumor of bone AND 2) One of the following: a) tumor is unresectable, OR b) surgical resection is likely to result in severe morbidity. Hypercalcemia of malignancy (HCM): Both of the following: 1) diagnosis of hypercalcemia of malignancy, AND 2) Trial and failure, contraindication, or intolerance to one intravenous bisphosphonate (eg, pamidronate, Zometa (zoledronic acid)).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	GCTB, HCM: Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	MM/BMST, GCTB: 12 mo. HCM: 2 mo.
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

385

## XIFAXAN (S)

---

### Products Affected

- Xifaxan

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Travelers' diarrhea (TD) (200 mg strength only): Diagnosis of travelers' diarrhea, AND one of the following: a) Trial and failure, contraindication, or intolerance to one of the following: Cipro (ciprofloxacin), Levaquin (levofloxacin), ofloxacin, Zithromax (azithromycin) OR b) resistance to all of the following: Cipro (ciprofloxacin), Levaquin (levofloxacin), ofloxacin, Zithromax (azithromycin). Prophylaxis of hepatic encephalopathy (HE) recurrence (550mg strength only): Used for the prophylaxis of hepatic encephalopathy recurrence, AND trial and failure, contraindication or intolerance to lactulose. Treatment of HE: Used for the treatment of HE. Trial and failure, contraindication, or intolerance to lactulose. Irritable bowel syndrome with diarrhea (IBS-D) (550mg strength only) (initial): Diagnosis of IBS-D, AND trial and failure, contraindication or intolerance to an antidiarrheal agent [eg, loperamide].
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	TD: 14 days. HE (prophylaxis, treatment): 12 months. IBS-D (initial, reauth): 2 weeks.
Other Criteria	IBS-D (reauth): Patient experiences IBS-D symptom recurrence.

Last Updated: June 2021

Formulary ID: 21381, Version 14

386

# **XOLAIR (S)**

---

## **Products Affected**

- Xolair

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	<p>Asthma (init): Diagnosis of moderate to severe persistent allergic asthma. Positive skin test or in vitro reactivity to a perennial aeroallergen. Pretreatment serum immunoglobulin (Ig)E level between 30 to 700 IU/mL for patients 12 years of age and older OR 30 to 1300 IU/mL for patients 6 years to less than 12 years of age. Patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications: a) Both of the following: i) High-dose inhaled corticosteroid (ICS) [e.g., greater than 500 mcg fluticasone propionate equivalent/day] and ii) additional asthma controller medication [e.g., leukotriene receptor antagonist, long-acting beta-2 agonist (LABA), tiotropium], OR b) One maximally-dosed combination ICS/LABA product [e.g., Advair (fluticasone propionate/salmeterol), Dulera (mometasone/formoterol), Symbicort (budesonide/formoterol), Breo Ellipta (fluticasone/vilanterol)].</p> <p>Chronic Idiopathic Urticaria (CIU) (init): Diagnosis of CIU. Persistent symptoms (itching and hives) for at least 4 consecutive weeks despite titrating to an optimal dose with a second generation H1 antihistamine (e.g., cetirizine, fexofenadine), unless there is a contraindication or intolerance to H1 antihistamines. Patient has tried and had an inadequate response or intolerance or contraindication to at least one of the following additional therapies: H2 antagonist (e.g., famotidine, cimetidine), leukotriene receptor antagonist (e.g., montelukast), H1 antihistamine, hydroxyzine, doxepin. Used concurrently with an H1 antihistamine, unless there is a contraindication or intolerance to H1 antihistamines.</p> <p>Nasal polyps (NP) (init): Diagnosis of NP. Unless contraindicated, the patient has had an inadequate response to 2 months of treatment with an intranasal corticosteroid (e.g., fluticasone, mometasone). Used in combination with another agent for nasal polyps (e.g., intranasal corticosteroid).</p>
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	<p>Asthma (init/reauth): Prescribed by or in consultation with an allergist/immunologist, or pulmonologist. CIU (init): Prescribed by or in consultation with an allergist/immunologist, or dermatologist. NP (init/reauth): Prescribed by or in consultation with an allergist/immunologist, otolaryngologist, or pulmonologist.</p>

Last Updated: June 2021

<b>Coverage Duration</b>	Asthma, init: 6 mo, reauth: 12 mo. CIU, init: 3 mo, reauth: 6 mo. NP, init/reauth: 12 mo.
<b>Other Criteria</b>	<p>Asthma (reauth): Documentation of positive clinical response to therapy (e.g., Reduction in number of asthma exacerbations, improvement in forced expiratory volume in 1 second (FEV1), or decreased use of rescue medications). Patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications: a) Both of the following: i) inhaled corticosteroid (ICS) (e.g., fluticasone, budesonide) and ii) additional asthma controller medication [e.g., leukotriene receptor antagonist, long-acting beta-2 agonist (LABA), tiotropium], OR b) A combination ICS/LABA product [e.g., Advair (fluticasone propionate/salmeterol), Dulera (mometasone/formoterol), Symbicort (budesonide/formoterol), Breo Ellipta (fluticasone/vilanterol)].</p> <p>CIU (reauth): Patient's disease status has been re-evaluated since the last authorization to confirm the patient's condition warrants continued treatment. Patient has experienced one or both of the following: Reduction in itching severity from baseline or Reduction in the number of hives from baseline. NP (reauth): Documentation of a positive clinical response to therapy (e.g., reduction in nasal polyps score [NPS: 0-8 scale], improvement in nasal congestion/obstruction score [NCS: 0-3 scale]). Used in combination with another agent for nasal polyps (e.g., intranasal corticosteroid).</p>

## XOSPATA (S)

---

### Products Affected

- Xospata

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of acute myeloid leukemia (AML). Disease is relapsed or refractory. Patient has a FMS-like tyrosine kinase (FLT3) mutation as determined by a U.S. Food and Drug Administration (FDA)-approved test (e.g., LeukoStrat CDx FLT3 Mutation Assay) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist or oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

390

## **XPOVIO (S)**

---

### **Products Affected**

- Xpovio
- Xpovio 100 Mg Once Weekly
- Xpovio 40 Mg Once Weekly
- Xpovio 40 Mg Twice Weekly
- Xpovio 60 Mg Once Weekly
- Xpovio 60 Mg Twice Weekly
- Xpovio 80 Mg Once Weekly
- Xpovio 80 Mg Twice Weekly

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple Myeloma (MM): Diagnosis of multiple myeloma. Patient has received at least one prior therapy. Used in combination with bortezomib and dexamethasone. Relapsed/Refractory Multiple Myeloma (RRMM): Diagnosis of relapsed or refractory multiple myeloma (RRMM). Patient has received at least four prior therapies. Disease is refractory to all of the following: 1) Two proteasome inhibitors (e.g., bortezomib, carfilzomib), 2) Two immunomodulatory agents (e.g., lenalidomide, thalidomide), and 3) An anti-CD38 monoclonal antibody (e.g. daratumumab). Used in combination with dexamethasone. Diffuse large B-cell lymphoma (DLBCL): Diagnosis of one of the following: 1) Relapsed or refractory DLBCL not otherwise specified OR 2) Relapsed or refractory DLBCL arising from follicular lymphoma. Patient has previously received at least two lines of systemic therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist/hematologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021



## XTANDI (S)

---

### Products Affected

- Xtandi

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Castration-resistant or castration-recurrent prostate cancer (CRPC): Diagnosis of castration-resistant (chemical or surgical) or recurrent prostate cancer. Metastatic castration-sensitive prostate cancer (M-CSPC): Diagnosis of metastatic castration-sensitive prostate cancer.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	CRPC, M-CSPC: Prescribed by or in consultation with an oncologist or urologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

393

## XYREM (S)

---

### Products Affected

- Xyrem

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Narcolepsy with cataplexy (Narcolepsy Type 1)(initial): Diagnosis of narcolepsy as confirmed by sleep study (unless the prescriber provides justification confirming that a sleep study would not be feasible), AND symptoms of cataplexy are present, AND symptoms of excessive daytime sleepiness (eg, irrepressible need to sleep or daytime lapses into sleep) are present. Narcolepsy without cataplexy (Narcolepsy Type 2)(initial): Diagnosis of narcolepsy as confirmed by sleep study (unless the prescriber provides justification confirming that a sleep study would not be feasible), AND symptoms of cataplexy are absent, AND symptoms of excessive daytime sleepiness (eg, irrepressible need to sleep or daytime lapses into sleep) are present, AND trial and failure, contraindication, or intolerance to one of the following: 1) amphetamine-based stimulant (eg, amphetamine, dextroamphetamine), OR 2) methylphenidate-based stimulant.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses (initial): Prescribed by or in consultation with one of the following: neurologist, psychiatrist, or sleep medicine specialist.
<b>Coverage Duration</b>	All uses (initial, reauth): 12 months
<b>Other Criteria</b>	Narcolepsy Type 1 (reauth): Documentation demonstrating a reduction in the frequency of cataplexy attacks associated with therapy, OR documentation demonstrating a reduction in symptoms of excessive daytime sleepiness associated with therapy. Narcolepsy Type 2 (reauth): Documentation demonstrating a reduction in symptoms of excessive daytime sleepiness associated with therapy.

Last Updated: June 2021

Last Updated: June 2021

Formulary ID: 21381, Version 14

395

## **XYWAV (S)**

---

### **Products Affected**

- Xywav

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Narcolepsy with cataplexy (Narcolepsy Type 1)(initial): Diagnosis of narcolepsy as confirmed by sleep study (unless the prescriber provides justification confirming that a sleep study would not be feasible), AND symptoms of cataplexy are present, AND symptoms of excessive daytime sleepiness (eg, irrepressible need to sleep or daytime lapses into sleep) are present. Narcolepsy without cataplexy (Narcolepsy Type 2)(initial): Diagnosis of narcolepsy as confirmed by sleep study (unless the prescriber provides justification confirming that a sleep study would not be feasible), AND symptoms of cataplexy are absent, AND symptoms of excessive daytime sleepiness (eg, irrepressible need to sleep or daytime lapses into sleep) are present, AND one of the following: 1) both of the following: A) patient is 18 years of age or older, and B) trial and failure, contraindication or intolerance to both of the following: i) generic modafinil, AND ii) generic armodafinil, OR 2) patient is 7 to 17 years of age.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses (initial): Prescribed by or in consultation with one of the following: neurologist, psychiatrist, or sleep medicine specialist.
<b>Coverage Duration</b>	All uses (initial, reauth): 12 months
<b>Other Criteria</b>	Narcolepsy with cataplexy (Narcolepsy Type 1)(reauth): Documentation demonstrating a reduction in the frequency of cataplexy attacks associated with therapy, OR documentation demonstrating a reduction in symptoms of excessive daytime sleepiness associated with therapy. Narcolepsy without Cataplexy (Narcolepsy Type 2)(reauth): Documentation demonstrating a reduction in symptoms of excessive daytime sleepiness associated with therapy.

Last Updated: June 2021

## YERVOY (S)

---

### Products Affected

- Yervoy

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

398

## ZAVESCA (S)

---

### Products Affected

- Miglustat

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Gaucher disease: Diagnosis of mild to moderate type 1 Gaucher disease. Patient is unable to receive enzyme replacement therapy due to one of the following conditions: allergy or hypersensitivity to enzyme replacement therapy, poor venous access, unavailability of enzyme replacement therapy (e.g., Cerezyme, VPRIV).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Gaucher disease: 12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

399

## **ZEJULA (S)**

---

### **Products Affected**

- Zejula



<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer: Diagnosis of one of the following: advanced epithelial ovarian cancer, advanced fallopian tube cancer, or advanced primary peritoneal cancer. Used for maintenance treatment in patients who are in a complete or partial response to first-line platinum-based chemotherapy (e.g., cisplatin, carboplatin). Recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer: Diagnosis of one of the following: recurrent epithelial ovarian cancer, recurrent fallopian tube cancer, or recurrent primary peritoneal cancer. Used for maintenance treatment in patients who are in a complete or partial response to platinum-based chemotherapy (e.g., cisplatin, carboplatin). Treatment of advanced ovarian cancer after three or more chemotherapies: Diagnosis of advanced ovarian cancer, advanced fallopian tube cancer, or advanced primary peritoneal cancer. Patient has been treated with three or more prior chemotherapy regimens. Patient's cancer is associated with homologous recombination deficiency (HRD) positive status defined by one of the following: (a) a deleterious or suspected deleterious BRCA mutation or (b) both of the following: (1) genomic instability and (2) cancer has progressed more than 6 months after response to the last platinum-based chemotherapy (e.g., cisplatin, carboplatin).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

## ZELBORAF (S)

---

### Products Affected

- Zelboraf

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Melanoma: Diagnosis of unresectable melanoma or metastatic melanoma. Cancer is BRAFV600 mutant type (MT) as detected by a U.S. Food and Drug Administration (FDA)-approved test (eg, cobas 4600 BRAFV600 Mutation Test) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Erdheim-Chester Disease: Diagnosis of Erdheim-Chester disease AND Disease is BRAFV600 mutant type (MT).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Melanoma: Prescribed by or in consultation with an oncologist. Erdheim-Chester Disease: Prescribed by or in consultation with a hematologist/oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	All indications: Approve for continuation of therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

402

## ZEPOSIA (S)

---

### Products Affected

- Zeposia
- Zeposia 7-day Starter Pack
- Zeposia Starter Kit

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple Sclerosis (MS): Diagnosis of a relapsing form of MS (e.g., clinically isolated syndrome, relapsing-remitting disease, secondary progressive disease, including active disease with new brain lesions). One of the following: a) Failure after a trial of at least 4 weeks, contraindication, or intolerance to two of the following disease-modifying therapies for MS: 1) Aubagio (teriflunomide), 2) Gilenya (fingolimod), or 3) Brand Tecfidera/generic dimethyl fumarate, OR b) for continuation of prior therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

403

## ZIRABEV (S)

---

### Products Affected

- Zirabev

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	Non-Small Cell Lung Cancer: Excluded if squamous cell histology.
<b>Required Medical Information</b>	<p>Colorectal Cancer: Diagnosis of metastatic colorectal cancer. One of the following: 1) Both of the following: a) used as first- or second-line treatment and b) used in combination with an intravenous 5-fluorouracil-based chemotherapy, OR 2) All of the following: a) used as second-line treatment, b) used in combination with fluoropyrimidine-irinotecan-based chemotherapy or fluoropyrimidine-oxaliplatin-based chemotherapy, and c) patient has progressed on a first-line bevacizumab-containing regimen.</p> <p>Non-Small Cell Lung Cancer (NSCLC): Diagnosis of NSCLC. Disease is unresectable, locally advanced, recurrent, or metastatic. Used as first-line treatment. Used in combination with paclitaxel and carboplatin.</p> <p>Renal Cell Cancer: Diagnosis of metastatic renal cell cancer. Used in combination with interferon-alpha.</p> <p>Cervical Cancer: Diagnosis of carcinoma of the cervix. Disease is persistent, recurrent, or metastatic. Used in combination with one of the following: a) paclitaxel and cisplatin or b) paclitaxel and topotecan.</p> <p>Glioblastoma: Diagnosis of recurrent glioblastoma.</p> <p>Epithelial Ovarian, Fallopian Tube, or Primary Peritoneal Cancer: Diagnosis of epithelial ovarian cancer, fallopian tube cancer, or primary peritoneal cancer. One of the following: 1) All of the following: a) disease is stage 3 or 4, b) patient has been treated with bevacizumab as a single agent, c) treatment is following surgical resection, and d) used in combination with carboplatin and paclitaxel, OR 2) All of the following: a) disease is platinum-resistant recurrent, b) patient has received no more than 2 prior chemotherapy regimens, and c) used in combination with paclitaxel, pegylated liposomal doxorubicin, or topotecan, OR 3) All of the following: a) disease is platinum-sensitive recurrent, b) patient has been treated with bevacizumab as a single agent, and c) used in combination with one of the following: i) carboplatin and paclitaxel or ii) carboplatin and gemcitabine.</p>
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months

Last Updated: June 2021

<b>Other Criteria</b>	Approve for continuation of prior therapy.
-----------------------	--------------------------------------------

# ZOLGENSMA (S)

---

## **Products Affected**

- Zolgensma 10.1-10.5 Kg
- Zolgensma 10.6-11.0 Kg
- Zolgensma 11.1-11.5 Kg
- Zolgensma 11.6-12.0 Kg
- Zolgensma 12.1-12.5 Kg
- Zolgensma 12.6-13.0 Kg
- Zolgensma 13.1-13.5 Kg
- Zolgensma 2.6-3.0 Kg
- Zolgensma 3.1-3.5 Kg
- Zolgensma 3.6-4.0 Kg
- Zolgensma 4.1-4.5 Kg
- Zolgensma 4.6-5.0 Kg
- Zolgensma 5.1-5.5 Kg
- Zolgensma 5.6-6.0 Kg
- Zolgensma 6.1-6.5 Kg
- Zolgensma 6.6-7.0 Kg
- Zolgensma 7.1-7.5 Kg
- Zolgensma 7.6-8.0 Kg
- Zolgensma 8.1-8.5 Kg
- Zolgensma 8.6-9.0 Kg
- Zolgensma 9.1-9.5 Kg
- Zolgensma 9.6-10.0 Kg

Last Updated: June 2021

Formulary ID: 21381, Version 14

407

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Spinal muscular atrophy (SMA): The mutation or deletion of genes in chromosome 5q resulting in one of the following: 1) Homozygous gene deletion or mutation of SMN1 gene (e.g., homozygous deletion of exon 7 at locus 5q13) or 2) Compound heterozygous mutation of SMN1 gene (e.g., deletion of Survival of Motor Neuron 1 [SMN1] exon 7 [allele 1] and mutation of SMN1 [allele 2]). One of the following: A) Both of the following: 1) Diagnosis of symptomatic Type I or Type II spinal muscular atrophy (SMA) confirmed by a neurologist with expertise in the diagnosis of SMA and 2) Patient is less than or equal to 2 years of age, OR B) All of the following: 1) Diagnosis of SMA based on the results of SMA newborn screening, 2) Patient has 3 copies or less of Survival of Motor Neuron 2 (SMN 2), and 3) Patient is less than or equal to 6 months of age. Patient is not dependent on either of the following: 1) Invasive ventilation or tracheostomy or 2) Use of non-invasive ventilation beyond use of naps and nighttime sleep. Submission of medical records (e.g., chart notes, laboratory values) documenting patient's anti-AAV9 antibody titers are less than or equal to 1:50. Patient is not to receive concomitant SMN modifying therapy (e.g. Spinraza). Patient has never received Zolgensma treatment in their lifetime.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by a neurologist with expertise in the diagnosis of SMA
<b>Coverage Duration</b>	1 Time Authorization in Lifetime
<b>Other Criteria</b>	N/A

Last Updated: June 2021



## ZOLINZA (S)

---

### Products Affected

- Zolinza

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cutaneous T-cell lymphoma (CTCL): Diagnosis of CTCL. Progressive, persistent or recurrent disease on or contraindication or intolerance to two systemic therapies (e.g., bexarotene, romidepsin, etc.).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist/oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

409

## ZORBTIVE (S)

---

### Products Affected

- Zorbtive

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Short Bowel Syndrome (SBS): Diagnosis of SBS. Patient is currently receiving specialized nutritional support (eg, intravenous parenteral nutrition, fluid, and micronutrient supplements). Patient has not previously received 4 weeks of treatment with Zorbtive (somatropin).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a gastroenterologist.
<b>Coverage Duration</b>	SBS: 4 weeks.
<b>Other Criteria</b>	N/A

Last Updated: June 2021

Formulary ID: 21381, Version 14

410

## ZYDELIG (S)

### Products Affected

- Zydelig

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chronic lymphocytic leukemia (CLL): Diagnosis of CLL. Used in combination with Rituxan (rituximab). The patient has relapsed on at least one prior therapy (eg, purine analogues [fludarabine, pentostatin, cladribine], alkylating agents [chlorambucil, cyclophosphamide], or monoclonal antibodies [rituximab]). Patient is a candidate for Rituxan (rituximab) monotherapy due to presence of other comorbidities (eg, coronary artery disease, peripheral vascular disease, diabetes mellitus, pulmonary disease [COPD]). Follicular Lymphoma (FL): Diagnosis of FL. The patient has relapsed on at least two prior systemic therapies (eg, rituximab, alkylating agents [cyclophosphamide, chlorambucil], anthracyclines [doxorubicin, daunorubicin], purine analogs [fludarabine]). Small lymphocytic lymphoma (SLL): Diagnosis of SLL. The patient has relapsed on at least two prior systemic therapies (eg, rituximab, alkylating agents [cyclophosphamide, chlorambucil], anthracyclines [doxorubicin, daunorubicin], purine analogs [fludarabine]).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses: Prescribed by or in consultation with an oncologist/hematologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

411

## ZYKADIA (S)

---

### Products Affected

- Zykadia

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Non-small cell lung cancer (NSCLC): Diagnosis of NSCLC that is metastatic or recurrent. Tumor is anaplastic lymphoma kinase (ALK)-positive as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

Last Updated: June 2021

Formulary ID: 21381, Version 14

412

## ZYTIGA (PREFERRED) (S)

---

### Products Affected

- Abiraterone Acetate

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Metastatic Castration-Resistant Prostate Cancer (mCRPC): Diagnosis of metastatic castration-resistant (chemical or surgical) or recurrent prostate cancer. Used in combination with prednisone. Metastatic Castration-Sensitive Prostate Cancer (mCSPC): Diagnosis of metastatic high-risk castration-sensitive prostate cancer. Used in combination with prednisone.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	mCRPC, mCSPC: Prescribed by or in consultation with an oncologist or urologist
<b>Coverage Duration</b>	mCRPC, mCSPC: 12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy

Last Updated: June 2021

Formulary ID: 21381, Version 14

413

# **PART B VERSUS PART D**

---

Last Updated: June 2021

Formulary ID: 21381, Version 14

## Products Affected

- Abelcet
- Acetylcysteine SOLN
- Acyclovir Sodium INJ 50MG/ML
- Adrucil INJ 2.5GM/50ML, 500MG/10ML, 5GM/100ML
- Akynzeo CAPS
- Albuterol Sulfate NEBU 0.083%, 0.63MG/3ML, 1.25MG/3ML, 2.5MG/0.5ML
- Ambisome
- Aminosyn II INJ 107.6MEQ/L; 1490MG/100ML; 1527MG/100ML; 1050MG/100ML; 1107MG/100ML; 750MG/100ML; 450MG/100ML; 990MG/100ML; 1500MG/100ML; 1575MG/100ML; 258MG/100ML; 447MG/100ML; 1083MG/100ML; 795MG/100ML; 50MEQ/L; 600MG/100ML; 300MG/100ML; 405MG/100ML; 750MG/100ML
- Amphotericin B INJ
- Anzemet TABS
- Aprepitant CAPS
- Azasan
- Azathioprine TABS
- Bleomycin
- Bleomycin Sulfate INJ
- Budesonide SUSP
- Clinimix 6/5
- Clinimix 8/10
- Clinimix 8/14
- Clinimix E 8/10
- Clinimix E 8/14
- Clinisol Sf 15%
- Cromolyn Sodium NEBU
- Cyclophosphamide CAPS
- Cyclosporine CAPS
- Cyclosporine Modified
- Cytarabine INJ 100MG/ML, 20MG/ML
- Cytarabine Aqueous
- Emend SUSR

- Engerix-b
- Everolimus TABS 0.25MG, 0.5MG, 0.75MG
- Fentanyl Citrate INJ 100MCG/2ML
- Fluorouracil INJ 1GM/20ML, 2.5GM/50ML, 500MG/10ML, 5GM/100ML
- Ganciclovir INJ 500MG, 500MG/10ML
- Gengraf CAPS 100MG, 25MG
- Gengraf SOLN
- Granisetron Hcl TABS
- Hepagam B
- Hyperhep B
- Hyperrab
- Imovax Rabies (h.d.c.v.)
- Intralipid INJ 20GM/100ML
- Ipratropium Bromide INHALATION SOLN 0.02%
- Ipratropium Bromide/albuterol Sulfate
- Morphine Sulfate INJ 10MG/ML, 1MG/ML, 4MG/ML, 5MG/ML, 8MG/ML
- Mycophenolate Mofetil CAPS
- Mycophenolate Mofetil SUSR
- Mycophenolate Mofetil TABS
- Mycophenolic Acid Dr
- Nabi-hb
- Nutrilipid
- Ondansetron Hcl SOLN
- Ondansetron Hcl TABS 24MG
- Ondansetron Hydrochloride TABS
- Ondansetron Odt
- Pentamidine Isethionate INHALATION SOLR
- Perforomist

Last Updated: June 2021

Formulary ID: 21381, Version 14

416



- Plenamine INJ 151MEQ/L;  
2170MG/100ML; 1470MG/100ML;  
434MG/100ML; 749MG/100ML;  
1040MG/100ML; 894MG/100ML;  
749MG/100ML; 1040MG/100ML;  
1180MG/100ML; 749MG/100ML;  
1040MG/100ML; 894MG/100ML;  
592MG/100ML; 749MG/100ML;  
250MG/100ML; 39MG/100ML;  
960MG/100ML
- Premasol INJ 56MEQ/L;  
320MG/100ML; 730MG/100ML;  
190MG/100ML; 3MEQ/L;  
20MG/100ML; 300MG/100ML;  
220MG/100ML; 290MG/100ML;  
490MG/100ML; 840MG/100ML;  
490MG/100ML; 200MG/100ML;  
290MG/100ML; 410MG/100ML;  
230MG/100ML; 5MEQ/L;  
15MG/100ML; 250MG/100ML;  
120MG/100ML; 140MG/100ML;  
470MG/100ML
- Prograf PACK
- Rabavert
- Recombivax Hb
- Sandimmune SOLN
- Sirolimus SOLN
- Sirolimus TABS
- Tacrolimus CAPS
- Tobramycin NEBU
- Yupelri
- Zortress TABS 1MG

## Details

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

Last Updated: June 2021

Formulary ID: 21381, Version 14

417

# INDEX

7	
7t Lido Gel .....	156
<b>A</b>	
Abelcet .....	370
Abiraterone Acetate.....	369
Abstral .....	190
Accutane.....	131
Acetylcysteine .....	370
Actemra .....	1
Actemra Actpen.....	1
Actemra Sc (s).....	1
Actimmune .....	3
Actimmune (s).....	3
Acyclovir Sodium .....	370
Adcirca (s).....	4
Adempas.....	5
Adempas (s).....	5
Adrucil.....	370
Afinitor .....	6
Afinitor (s).....	6
Afinitor Disperz.....	7
Afinitor Disperz (s) .....	7
Aimovig.....	8
Aimovig (s) .....	8
Akynzeo .....	370
Albuterol Sulfate .....	370
Aldurazyme .....	10
Aldurazyme (s).....	10
Alecensa .....	11
Alecensa (s).....	11
Alosetron Hydrochloride.....	160
Alpha-1 Proteinase Inhibitor, Non-preferred (s). 12	
Alpha-1 Proteinase Inhibitor, Prolastin (s).....	13
Alunbrig .....	14
Alunbrig (s) .....	14
Alyq.....	4
Ambisome.....	370
Ambrisentan.....	154
Aminosyn II .....	370
Amitriptyline Hcl.....	114
Amitriptyline Hydrochloride .....	114
Amnesteem .....	131
Amphotericin B.....	370
Ampyra (s) .....	15
Anadrol-50 .....	16
Anadrol-50 (s).....	16
Androderm .....	307
Anzemet .....	370
Apokyn.....	17
Apokyn (s) .....	17
Aprepitant .....	370
Aralast Np .....	12
Aranesp (s).....	18
Aranesp Albumin Free.....	18
Arcalyst.....	20
Arcalyst (s).....	20
Auryxia .....	22
Auryxia (s) .....	22
Austedo .....	23
Austedo (s).....	23
Avastin .....	24
Avastin (s).....	24
Avita.....	314
Avonex.....	179
Avonex Pen.....	179
Ayvakit.....	26
Ayvakit (s) .....	26
Azasan.....	370
Azathioprine.....	370
<b>B</b>	
Bafiertam.....	27
Bafiertam (s) .....	27

Last Updated: June 2021

Balversa.....	28	Cerezyme (s).....	44
Balversa (s).....	28	Chenodal.....	45
Bavencio.....	29	Chenodal (s).....	45
Bavencio (s).....	29	Chlorzoxazone.....	113
Benlysta.....	30	Cholbam.....	46
Benlysta (s).....	30	Cholbam (s).....	46
Berinert.....	31	Ciclodan.....	47
Berinert (s).....	31	Ciclopirox (s).....	47
Betaseron.....	179	Ciclopirox Nail Lacquer.....	47
Bexarotene.....	298	Cinryze.....	48
Bivigam.....	134	Cinryze (s).....	48
Bleomycin.....	370	Claravis.....	131
Bleomycin Sulfate.....	370	Clinimix 6/5.....	370
Bosulif.....	32	Clinimix 8/10.....	370
Bosulif (s).....	32	Clinimix 8/14.....	370
Braftovi.....	33	Clinimix E 8/10.....	370
Braftovi (s).....	33	Clinimix E 8/14.....	370
Briviact.....	34	Clinisol Sf 15%.....	370
Briviact (s).....	34	Clovique.....	289
Bronchitol.....	35	Cometriq.....	49
Bronchitol (s).....	35	Cometriq (s).....	49
Bronchitol Tolerance Test.....	35	Copiktra.....	50
Brukinsa.....	36	Copiktra (s).....	50
Brukinsa (s).....	36	Corlanor.....	51
Budesonide.....	370	Corlanor (s).....	51
Bynfezia Pen.....	37	Cosentyx.....	52
Bynfezia (s).....	37	Cosentyx (s).....	52
<b>C</b>			
Cablivi.....	38	Cosentyx Sensoready Pen.....	52
Cablivi (s).....	38	Cotellic.....	54
Cabometyx.....	39	Cotellic (s).....	54
Cabometyx (s).....	39	Cromolyn Sodium.....	370
Calquence.....	40	Cuvitru.....	264
Calquence (s).....	40	Cyclobenzaprine Hydrochloride.....	113
Caprelsa.....	41	Cyclophosphamide.....	370
Caprelsa (s).....	41	Cyclosporine.....	370
Carimune Nanofiltered.....	134	Cyclosporine Modified.....	370
Cayston.....	42	Cystadrops.....	55
Cayston (s).....	42	Cystadrops (s).....	55
Cerdelga.....	43	Cystaran.....	56
Cerdelga (s).....	43	Cystaran (s).....	56
Cerezyme.....	44	Cytarabine.....	370
		Cytarabine Aqueous.....	370

Last Updated: June 2021

**D**

D.h.e. 45 (s).....	57
Dalfampridine Er.....	15
Daliresp .....	58
Daliresp (s) .....	58
Daraprim (s) .....	59
Daurismo .....	60
Daurismo (s).....	60
Deferasirox .....	61
Deferasirox (s).....	61
Deferiprone.....	87
Diacomit.....	62
Diacomit (s).....	62
Diclofenac Sodium.....	224
Dihydroergotamine Mesylate.....	57, 176
Dimethyl Fumarate.....	302
Dimethyl Fumarate Starterpack .....	302
Dojolvi.....	63
Dojolvi (s) .....	63
Doxepin Hcl .....	114
Doxepin Hydrochloride.....	114
Dronabinol.....	170
Droxidopa.....	191
Dupixent.....	64
Dupixent (s).....	64
Dysport.....	66
Dysport (s).....	66

**E**

Elaprase .....	67
Elaprase (s).....	67
Emend.....	370
Emgality .....	68
Emgality (s).....	68
Enbrel .....	70
Enbrel (s).....	70
Enbrel Mini .....	70
Enbrel Sureclick .....	70
Endari .....	71
Endari (s).....	71
Engerix-b.....	370
Enspryng.....	72
Enspryng (s) .....	72

Last Updated: June 2021

Entyvio.....	73
Entyvio (s).....	73
Epclusa Preferred (s).....	74
Epidiolex.....	75
Epidiolex (s).....	75
Erivedge .....	76
Erivedge (s).....	76
Erleada .....	77
Erleada (s) .....	77
Erlotinib Hydrochloride .....	297
Esbriet .....	78
Esbriet (s) .....	78
Eucrisa.....	79
Eucrisa (s) .....	79
Everolimus .....	6, 370
Evrysdi .....	80
Evrysdi (s).....	80

**F**

Fabrazyme.....	82
Fabrazyme (s).....	82
Farydak .....	83
Farydak (s) .....	83
Fasenra .....	84
Fasenra (s).....	84
Fasenra Pen .....	84
Fentanyl (s) .....	86
Fentanyl Citrate.....	190, 370
Fentanyl Citrate Oral Transmucosal .....	86
Ferriprox .....	87
Ferriprox (s) .....	87
Ferriprox Twice-a-day .....	87
Fintepla .....	88
Fintepla (s) .....	88
Firazyr (s).....	89
Firdapse.....	90
Firdapse (s).....	90
Firmagon .....	91
Firmagon (s).....	91
Flebogamma Dif .....	134
Fluorouracil.....	370
Forteo .....	305
Fotivda .....	92

Fotivda (s) .....	92
<b>G</b>	
Galafold.....	93
Galafold (s).....	93
Gamastan.....	94
Gamastan S/d (s) .....	94
Gamifant.....	95
Gamifant (s).....	95
Gammagard Liquid .....	134
Gammagard S/d Iga Less Than 1mcg/ml.....	134
Gammaked .....	134
Gammaplex .....	134
Gamunex-c .....	134
Ganciclovir .....	370
Gattex .....	96
Gattex (s).....	96
Gavreto.....	97
Gavreto (s).....	97
Gengraf.....	370
Genotropin.....	103
Genotropin Miniquick.....	103
Gilenya .....	98
Gilenya (s).....	98
Gilotrif.....	99
Gilotrif (s).....	99
Givlaari.....	100
Givlaari (s).....	100
Glatiramer Acetate .....	101
Glatiramer Acetate (s).....	101
Glatopa .....	101
Gleevec (s).....	102
Glydo.....	156
Granisetron Hcl .....	370
Growth Hormone, Preferred (s) .....	103

**H**

Haegarda.....	106
Haegarda (s) .....	106
Hepagam B.....	370
Herceptin .....	107
Herceptin (s).....	107
Hetlioz .....	108

Hetlioz (s).....	108
Hetlioz Lq .....	109
Hetlioz Lq (s) .....	109
Hizentra.....	264
Hrm - Antipsychotics .....	110
Hrm - Endocrine .....	111
Hrm - Phenobarbital, Pentobarbital .....	112
Hrm - Skeletal Muscle Relaxants .....	113
Hrm - Tca.....	114
Humira .....	115
Humira (s).....	115
Humira Pediatric Crohns Disease Starter Pack..	115
Humira Pen .....	115
Humira Pen-cd/uc/hs Starter .....	115
Humira Pen-pediatric Uc Starter Pack .....	115
Humira Pen-ps/uv Starter.....	115
Hydroxyprogesterone Caproate .....	169
Hyperhep B.....	370
Hyperrab .....	370

**I**

Ibrance.....	117
Ibrance (s) .....	117
Icatibant Acetate .....	89
Iclusig.....	118
Iclusig (s) .....	118
Icosapent Ethyl.....	331
Idhifa .....	119
Idhifa (s).....	119
Ilumya .....	120
Ilumya (s) .....	120
Imatinib Mesylate .....	102
Imbruvica .....	121
Imbruvica (s).....	121
Imfinzi.....	122
Imfinzi (s).....	122
Imovax Rabies (h.d.c.v.).....	370
Increlex .....	123
Increlex (s) .....	123
Inflectra .....	124
Inflectra (s).....	124
Inlyta .....	126
Inlyta (s).....	126

Last Updated: June 2021

Inqovi .....	127
Inqovi (s) .....	127
Inrebic.....	128
Inrebic (s) .....	128
Intralipid.....	370
Intron A .....	129
Intron A (s).....	129
Ipratropium Bromide.....	370
Ipratropium Bromide/albuterol Sulfate.....	370
Iressa.....	130
Iressa (s) .....	130
Isotretinoin .....	131
Isotretinoin (s) .....	131
Istodax (s).....	132
Isturisa .....	133
Isturisa (s).....	133
Itraconazole .....	272
Ivig (s).....	134

## **J**

Jadenu Sprinkle .....	61
Jakafi .....	137
Jakafi (s) .....	137
Juxtapid .....	138
Juxtapid (s) .....	138

## **K**

Kalbitor.....	139
Kalbitor (s) .....	139
Kalydeco.....	140
Kalydeco (s) .....	140
Kanjinti.....	141
Kanjinti (s).....	141
Kanuma .....	142
Kanuma (s) .....	142
Kesimpta.....	143
Kesimpta (s) .....	143
Keytruda .....	145
Keytruda (s).....	145
Kisqali .....	146
Kisqali (s).....	146
Kisqali Femara 200 Dose .....	147
Kisqali Femara 400 Dose .....	147

Kisqali Femara 600 Dose.....	147
Kisqali-femara Pack (s).....	147
Korlym .....	148
Korlym (s).....	148
Koselugo .....	149
Koselugo (s).....	149
Kuvan.....	150
Kuvan (s).....	150
Kynmobi .....	151
Kynmobi (s) .....	151
Kynmobi Titration Kit .....	151

## **L**

Lapatinib Ditosylate.....	321
Lazanda .....	190
Lemtrada .....	152
Lemtrada (s) .....	152
Lenvima (s) .....	153
Lenvima 10 Mg Daily Dose.....	153
Lenvima 12mg Daily Dose .....	153
Lenvima 14 Mg Daily Dose.....	153
Lenvima 18 Mg Daily Dose.....	153
Lenvima 20 Mg Daily Dose.....	153
Lenvima 24 Mg Daily Dose.....	153
Lenvima 4 Mg Daily Dose.....	153
Lenvima 8 Mg Daily Dose.....	153
Letairis (s) .....	154
Leuprolide Acetate.....	164
Libtayo .....	155
Libtayo (s).....	155
Lidocaine.....	156, 157
Lidocaine Hcl.....	156
Lidocaine Hcl Jelly .....	156
Lidocaine Topical (s) .....	156
Lidocaine/prilocaine .....	156
Lidoderm (s).....	157
Lonsurf.....	158
Lonsurf (s).....	158
Lorbrena .....	159
Lorbrena (s).....	159
Lotronex (s).....	160
Lovaza (s).....	161
Lumizyme .....	162

Last Updated: June 2021

Lumizyme (s) .....	162
Lupkynis.....	163
Lupkynis (s).....	163
Lupron (s).....	164
Lupron Depot (1-month) .....	165
Lupron Depot (3-month) .....	165
Lupron Depot (4-month) .....	165
Lupron Depot (6-month) .....	165
Lupron Depot (s) .....	165
Lupron Depot Ped (s) .....	166
Lupron Depot-ped (1-month) .....	166
Lupron Depot-ped (3-month) .....	166
Lynparza.....	167
Lynparza Tablet (s) .....	167

**M**

Makena .....	169
Makena (s).....	169
Marinol (s).....	170
Mavyret .....	171
Mavyret (s) .....	171
Megestrol Acetate .....	111
Mekinist.....	172
Mekinist (s) .....	172
Mektovi .....	174
Mektovi (s) .....	174
Methotrexate Injection (s) .....	175
Miglustat.....	357
Migranal (s).....	176
Mitoxantrone Hcl .....	192
Modafinil.....	231
Morphine Sulfate.....	370
Mozobil .....	177
Mozobil (s) .....	177
Ms Interferons (non-preferred) (s) .....	178
Ms Interferons (preferred) (s).....	179
Mvasi.....	180
Mvasi (s).....	180
Myalept.....	182
Myalept (s) .....	182
Mycapssa.....	183
Mycapssa (s).....	183
Mycophenolate Mofetil.....	370

Mycophenolic Acid Dr .....	370
Myorisan .....	131

**N**

Nabi-hb .....	370
Naglazyme .....	184
Naglazyme (s) .....	184
Natpara.....	185
Natpara (s).....	185
Nerlynx .....	186
Nerlynx (s) .....	186
Neulasta.....	187
Neulasta (s) .....	187
Neulasta Onpro Kit .....	187
Nexavar .....	188
Nexavar (s).....	188
Ninlaro .....	189
Ninlaro (s) .....	189
Non-preferred Tirf (s) .....	190
Northera .....	191
Northera (s) .....	191
Novantrone (s) .....	192
Nplate.....	194
Nplate (s).....	194
Nubeqa .....	195
Nubeqa (s).....	195
Nucala .....	196
Nucala (s) .....	196
Nuedexta .....	198
Nuedexta (s) .....	198
Nuplazid.....	199
Nuplazid (s).....	199
Nurtec.....	200
Nurtec (s) .....	200
Nutrilipid.....	370
Nyvepria.....	201
Nyvepria (s) .....	201

**O**

Octagam .....	134
Octreotide Acetate .....	263
Odomzo.....	202
Odomzo (s).....	202

Last Updated: June 2021

Ofev.....	203	Palforzia Level 11 (maintenance).....	221
Ofev (s).....	203	Palforzia Level 11 (titration).....	221
Omega-3-acid Ethyl Esters.....	161	Palforzia Level 2.....	221
Ondansetron Hcl.....	370	Palforzia Level 3.....	221
Ondansetron Hydrochloride.....	370	Palforzia Level 4.....	221
Ondansetron Odt.....	370	Palforzia Level 5.....	221
Onureg.....	205	Palforzia Level 6.....	221
Onureg (s).....	205	Palforzia Level 7.....	221
Opdivo.....	206	Palforzia Level 8.....	221
Opdivo (s).....	206	Palforzia Level 9.....	221
Opsumit.....	207	Panzyga.....	134
Opsumit (s).....	207	Part B Versus Part D.....	370
Orencia.....	208, 209	Pegasys.....	222
Orencia Clickject.....	209	Pegasys (s).....	222
Orencia IV (s).....	208	Pegasys Proclick.....	222
Orencia Sc (s).....	209	Pemazyre.....	223
Orenitram.....	210	Pemazyre (s).....	223
Orenitram (s).....	210	Pennsaid (s).....	224
Orgovyx.....	211	Pentamidine Isethionate.....	370
Orgovyx (s).....	211	Perforomist.....	370
Oriahnn.....	212	Phenobarbital.....	112
Oriahnn (s).....	212	Phenobarbital Sodium.....	112
Orilissa.....	213	Piqray (s).....	225
Orilissa (s).....	213	Piqray 200mg Daily Dose.....	225
Orkambi.....	214, 215	Piqray 250mg Daily Dose.....	225
Orkambi (s).....	214	Piqray 300mg Daily Dose.....	225
Orkambi Granules (s).....	215	Plegridy.....	179
Orladeyo.....	216	Plegridy Starter Pack.....	179
Orladeyo (s).....	216	Plenammine.....	371
Osmolex Er.....	217	Polivy.....	226
Osmolex Er (s).....	217	Polivy (s).....	226
Osphena.....	218	Pomalyst.....	227
Osphena (s).....	218	Pomalyst (s).....	227
Oxandrin (s).....	219	Premasol.....	371
Oxandrolone.....	219	Privigen.....	134
Oxervate.....	220	Procysbi.....	228
Oxervate (s).....	220	Procysbi (s).....	228
<b>P</b>		Prograf.....	371
Palforzia (s).....	221	Prolastin-c.....	13
Palforzia Initial Dose Escalation.....	221	Promacta.....	229
Palforzia Level 1.....	221	Promacta (s).....	229
Palforzia Level 10.....	221	Provigil (s).....	231
		Pulmozyme.....	233

Last Updated: June 2021



Pulmozyme (s).....	233
Pyrimethamine .....	59
<b>Q</b>	
Qinlock .....	234
Qinlock (s).....	234
Qualaquin (s).....	235
Quinine Sulfate.....	235
<b>R</b>	
Rabavert .....	371
Radicava .....	236
Radicava (s).....	236
Rasuvo.....	175
Ravicti .....	237
Ravicti (s) .....	237
Rebif.....	178
Rebif Rebidose .....	178
Rebif Rebidose Titration Pack .....	178
Rebif Titration Pack .....	178
Reblozyl .....	238
Reblozyl (s) .....	238
Recombivax Hb.....	371
Renflexis.....	239
Renflexis (s) .....	239
Repatha.....	241
Repatha (s).....	241
Repatha Pushtrex System .....	241
Repatha Sureclick.....	241
Retacrit .....	244
Retacrit (s).....	244
Retevmo .....	246
Retevmo (s) .....	246
Revatio (s) .....	247
Revcovi.....	248
Revcovi (s) .....	248
Revlimid.....	249
Revlimid (s).....	249
Rilutek (s).....	250
Riluzole .....	250
Rinvoq.....	251
Rinvoq (s).....	251
Rituxan .....	252

Rituxan (s).....	252
Rituxan Hycela.....	254
Rituxan Hycela (s) .....	254
Romidepsin .....	132
Rozlytrek.....	255
Rozlytrek (s).....	255
Rubraca .....	256
Rubraca (s) .....	256
Ruconest.....	257
Ruconest (s) .....	257
Ruxience .....	258
Ruxience (s) .....	258
Ruzurgi.....	260
Ruzurgi (s) .....	260
Rydapt.....	261
Rydapt (s).....	261
<b>S</b>	
Sabril (s).....	262
Sandimmune .....	371
Sandostatin (s).....	263
Sapropterin Dihydrochloride .....	150
Scig (s) .....	264
Secuado .....	265
Secuado (s).....	265
Signifor .....	266
Signifor (s) .....	266
Siklos.....	267
Siklos (s) .....	267
Sildenafil Citrate .....	247
Sirolimus .....	371
Skyrizi .....	268
Skyrizi (s).....	268
Skyrizi Pen .....	268
Sofosbuvir/velpatasvir .....	74
Soliris .....	269
Soliris (s).....	269
Somatuline Depot.....	270
Somatuline Depot (s) .....	270
Somavert .....	271
Somavert (s) .....	271
Sporanox (s) .....	272
Spravato (s) .....	273

Last Updated: June 2021

Spravato 56mg Dose .....	273	Tarceva (s) .....	297
Spravato 84mg Dose .....	273	Targretin.....	298
Sprycel.....	274	Targretin (s) .....	298
Sprycel (s) .....	274	Tasigna.....	299
Stelara.....	275, 276	Tasigna (s).....	299
Stelara (iv) (s).....	275	Tazverik .....	300
Stelara (s).....	276	Tazverik (s).....	300
Stivarga.....	278	Tecentriq .....	301
Stivarga (s) .....	278	Tecentriq (s) .....	301
Strensiq.....	279	Tecfidera .....	302
Strensiq (s).....	279	Tecfidera (s).....	302
Supprelin La .....	280	Tecfidera Starter Pack.....	302
Supprelin La (s).....	280	Tegsedi .....	303
Sutent.....	281	Tegsedi (s).....	303
Sutent (s) .....	281	Tepmetko .....	304
Sylatron .....	282	Tepmetko (s) .....	304
Sylatron (s) .....	282	Teriparatide.....	305
Sylvant.....	283	Teriparatide (s).....	305
Sylvant (s) .....	283	Testosterone .....	307
Symdeko.....	284	Testosterone (s).....	307
Symdeko (s).....	284	Testosterone Cypionate.....	307
Synagis .....	285	Testosterone Enanthate .....	309
Synagis (s).....	285	Testosterone Enanthate (s).....	309
Syndros.....	287	Testosterone Pump.....	307
Syndros (s).....	287	Tetrabenazine.....	345
Synribo .....	288	Thalomid.....	311
Synribo (s).....	288	Thalomid (s).....	311
Syprine (s) .....	289	Thioridazine Hcl .....	110
<b>T</b>			
Tabrecta.....	290	Tibsovo .....	312
Tabrecta (s).....	290	Tibsovo (s) .....	312
Tacrolimus.....	371	Tiglutik.....	313
Tadalafil .....	4	Tiglutik (s) .....	313
Tafamidis (s) .....	291	Tobramycin.....	371
Tafinlar .....	292	Topical Retinoid (s) .....	314
Tafinlar (s).....	292	Trazimera .....	315
Tagrisso .....	294	Trazimera (s).....	315
Tagrisso (s).....	294	Trelstar (s).....	316
Takhzyro.....	295	Trelstar Mixject.....	316
Takhzyro (s) .....	295	Tretinoin.....	314
Talzenna .....	296	Trientine Hydrochloride.....	289
Talzenna (s).....	296	Trikafta.....	317
		Trikafta (s) .....	317
		Triptodur .....	318

Last Updated: June 2021

Triptodur (s) .....	318
Tukysa .....	319
Tukysa (s) .....	319
Turalio .....	320
Turalio (s) .....	320
Tykerb .....	321
Tykerb (s) .....	321
Tymlos .....	322
Tymlos (s) .....	322
Tysabri .....	324
Tysabri (s) .....	324

## U

Udenyca .....	326
Udenyca (s) .....	326
Ukoniq .....	327
Ukoniq (s) .....	327
Ultomiris .....	328
Ultomiris (s) .....	328

## V

Valchlor .....	329
Valchlor (s) .....	329
Varizig .....	330
Varizig (s) .....	330
Vascepa .....	331
Vascepa (s) .....	331
Venclexta .....	332
Venclexta (s) .....	332
Venclexta Starting Pack .....	332
Ventavis .....	333
Ventavis (s) .....	333
Verzenio .....	334
Verzenio (s) .....	334
Vigabatrin .....	262
Vigadrone .....	262
Vimizim .....	335
Vimizim (s) .....	335
Vitrakvi .....	336
Vitrakvi (s) .....	336
Vizimpro .....	337
Vizimpro (s) .....	337
Vosevi .....	338

Vosevi (s) .....	338
Votrient .....	339
Votrient (s) .....	339
Vpriv .....	340
Vpriv (s) .....	340
Vyndamax .....	291
Vyndaquel .....	291

## X

Xalkori .....	341
Xalkori (s) .....	341
Xcopri .....	342
Xcopri (s) .....	342
Xeljanz .....	343
Xeljanz (s) .....	343
Xeljanz Xr .....	343
Xembify .....	264
Xenazine (s) .....	345
Xgeva .....	346
Xgeva (s) .....	346
Xifaxan .....	347
Xifaxan (s) .....	347
Xolair .....	348
Xolair (s) .....	348
Xospata .....	350
Xospata (s) .....	350
Xpovio .....	351
Xpovio (s) .....	351
Xpovio 100 Mg Once Weekly .....	351
Xpovio 40 Mg Once Weekly .....	351
Xpovio 40 Mg Twice Weekly .....	351
Xpovio 60 Mg Once Weekly .....	351
Xpovio 60 Mg Twice Weekly .....	351
Xpovio 80 Mg Once Weekly .....	351
Xpovio 80 Mg Twice Weekly .....	351
Xtandi .....	352
Xtandi (s) .....	352
Xyrem .....	353
Xyrem (s) .....	353
Xywav .....	354
Xywav (s) .....	354

Last Updated: June 2021

<b>Y</b>	
Yervoy.....	356
Yervoy (s).....	356
Yupelri.....	371
<b>Z</b>	
Zavesca (s).....	357
Zejula.....	358
Zejula (s).....	358
Zelboraf.....	359
Zelboraf (s).....	359
Zemaira.....	12
Zenatane.....	131
Zeposia.....	360
Zeposia (s).....	360
Zeposia 7-day Starter Pack.....	360
Zeposia Starter Kit.....	360
Zirabev.....	361
Zirabev (s).....	361
Zolgensma (s).....	363
Zolgensma 10.1-10.5 Kg.....	363
Zolgensma 10.6-11.0 Kg.....	363
Zolgensma 11.1-11.5 Kg.....	363
Zolgensma 11.6-12.0 Kg.....	363
Zolgensma 12.1-12.5 Kg.....	363
Zolgensma 12.6-13.0 Kg.....	363
Zolgensma 13.1-13.5 Kg.....	363
Zolgensma 2.6-3.0 Kg.....	363
Zolgensma 3.1-3.5 Kg.....	363
Zolgensma 3.6-4.0 Kg.....	363
Zolgensma 4.1-4.5 Kg.....	363
Zolgensma 4.6-5.0 Kg.....	363
Zolgensma 5.1-5.5 Kg.....	363
Zolgensma 5.6-6.0 Kg.....	363
Zolgensma 6.1-6.5 Kg.....	363
Zolgensma 6.6-7.0 Kg.....	363
Zolgensma 7.1-7.5 Kg.....	363
Zolgensma 7.6-8.0 Kg.....	363
Zolgensma 8.1-8.5 Kg.....	363
Zolgensma 8.6-9.0 Kg.....	363
Zolgensma 9.1-9.5 Kg.....	363
Zolgensma 9.6-10.0 Kg.....	363
Zolinza.....	365
Zolinza (s).....	365
Zorbtive.....	366
Zorbtive (s).....	366
Zortress.....	371
Zydelig.....	367
Zydelig (s).....	367
Zykadia.....	368
Zykadia (s).....	368
Zytiga (preferred) (s).....	369